



156

(51) International Patent Classification ⁶ : C12N 15/12, C07K 14/705, C12Q 1/68		A1	(11) International Publication Number: WO 96/05302
			(43) International Publication Date: 22 February 1996 (22.02.96)
(21) International Application Number: PCT/JP95/01599			
(22) International Filing Date: 10 August 1995 (10.08.95)			
(30) Priority Data:			
6/189272	11 August 1994 (11.08.94)	JP	
6/189273	11 August 1994 (11.08.94)	JP	
6/189274	11 August 1994 (11.08.94)	JP	
6/236356	30 September 1994 (30.09.94)	JP	
6/236357	30 September 1994 (30.09.94)	JP	
6/270017	2 November 1994 (02.11.94)	JP	
6/326611	28 December 1994 (28.12.94)	JP	
7/7177	20 January 1995 (20.01.95)	JP	
7/57186	16 March 1995 (16.03.95)	JP	
7/93989	19 April 1995 (19.04.95)	JP	
(71) Applicant (for all designated States except US): TAKEDA CHEMICAL INDUSTRIES, LTD. [JP/JP]; 1-1, Doshomachi 4-chome, Chuo-ku, Osaka-shi, Osaka 541 (JP).			
(72) Inventors; and			
(75) Inventors/Applicants (for US only): HINUMA, Shuji [JP/JP]; 7-9-1402, Kasuga 1-chome, Tsukuba-shi, Ibaraki 305 (JP). HOSOYA, Masaki [JP/JP]; 711-83, Itaya 1-chome, Tsuchiura-shi, Ibaraki 300 (JP). FUJII, Ryo [JP/JP];			
		7-9-303, Kasuga 1-chome, Tsukuba-shi, Ibaraki 305 (JP). OHTAKI, Tetsuya [JP/JP]; 7-9-802, Kasuga 1-chome, Tsukuba-shi, Ibaraki 305 (JP). FUKUSUMI, Shoji [JP/JP]; 17-6-302, Namiki 3-chome, Tsukuba-shi, Ibaraki 305 (JP). OHGI, Kazuhiro [JP/JP]; 16-1-206, Umezono 2-chome, Tsukuba-shi, Ibaraki 305 (JP). (74) Agent: MIZUNO, Akinobu; Gloria Miyamasuzaka III 305, 10-7, Shibuya 1-chome, Shibuya-ku, Tokyo 150 (JP). (81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, KE, KG, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TT, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG). Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.	
(54) Title: G PROTEIN COUPLED RECEPTOR PROTEIN, PRODUCTION, AND USE THEREOF			
(57) Abstract			
<p>DNA primers effective in screening G protein coupled receptor protein-encoding DNA fragments are provided. The primers which are complementary to nucleotide sequences that are in community with (homologous to) the nucleotide sequences encoding amino acid sequences corresponding to or near the first membrane-spanning domain or the sixth membrane-spanning domain each of known various G protein coupled receptor proteins were designed and synthesized. Methods of amplifying G protein coupled receptor protein-encoding DNAs using the above DNA primers, and novel target G protein coupled receptor protein-encoding DNAs are also provided. Screening of DNA libraries can be efficiently carried out. Human pituitary gland or amygdala-derived and mouse pancreas-derived G protein coupled receptor proteins, etc., or salts thereof, partial peptides thereof, DNAs coding for the above G protein coupled receptor proteins, processes for producing the above G protein coupled receptor proteins, methods of determining ligands for the above G protein coupled receptor proteins, methods of screening compounds that inhibit the binding between the ligand and the G protein coupled receptor proteins or screening kits therefor, compounds or salts thereof obtained by the above screening method or the screening kit, pharmaceutical compositions containing the above compounds or salts thereof, and antibodies against the above protein coupled receptor proteins or partial peptides thereof are provided.</p>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

DESCRIPTION

G PROTEIN COUPLED RECEPTOR PROTEIN,
PRODUCTION, AND USE THEREOFFIELD OF THE INVENTION

5 The present invention relates to novel DNAs which are
useful as DNA primers for a polymerase chain reaction (PCR);
methods for amplifying DNAs each coding for a G protein
coupled receptor protein via PCR techniques using said DNA;
screening methods for DNAs each encoding a G protein coupled
10 receptor protein via PCR techniques using said DNA;
G protein coupled receptor protein-encoding DNAs obtained by
said screening method; G protein coupled receptor proteins
which are encoded by the DNA obtained via said screening
method, peptide fragments or segments thereof, and modified
15 peptide derivatives thereof; etc.

 The present invention also relates to novel G protein
coupled receptor proteins; novel G protein coupled receptor
protein-encoding DNAs; processes for producing said G protein
coupled receptor protein; use of said receptor protein and
20 said protein-encoding DNA; etc.

 The present invention also relates to novel human
amygdaloid nucleus-derived G protein coupled receptor proteins;
novel DNAs each coding for said G protein coupled receptor
protein; processes for producing said G protein coupled
25 receptor protein; use of said receptor protein and
said protein-encoding DNA; etc.

 The present invention also relates to novel mouse
pancreatic β cell line MIN6-derived G protein coupled receptor
proteins; novel DNAs each coding for said G protein coupled
30 receptor protein; processes for producing said G protein
coupled receptor protein; use of said receptor protein and

said protein-encoding DNA; etc. Further, the present invention relates to novel human-derived G protein coupled receptor proteins (human prinoceptors); novel DNAs each coding for said G protein coupled receptor protein; processes for producing
5 said G protein coupled receptor protein; use of said receptor protein and said protein-encoding DNA; etc.

BACKGROUND OF THE INVENTION

A variety of hormones, neurotransmitters and the like control, regulate or adjust the functions of living bodies via
10 specific receptors located in cell membranes. Many of these receptors mediate the transmission of intracellular signals via activation of guanine nucleotide-binding proteins (hereinafter, sometimes referred to as G proteins) with which the receptor is coupled and possess the common (homologous) structure, i.e.
15 seven transmembranes (membrane-spanning regions (domains)). Therefore, such receptors are generically referred to as G protein coupled receptors or seven transmembrane (membrane-spanning) receptors.

G protein coupled receptor proteins have a very
20 important role as targets for molecules such as hormones, neurotransmitters and physiologically active substances, which molecules control, regulate or adjust the functions of living bodies. Each molecule has its own receptor protein which is specific thereto, whereby the specificities of individual
25 physiologically active substances, including specific target cells and organs, specific pharmacological actions, specific action strength, action time, etc., are decided. Accordingly, it has been believed that, if G protein coupled receptor genes or cDNA can be cloned, those will be helpful not only for the
30 clarification of structure, function, physiological action, etc. of the G protein coupled receptor but also for the development of pharmaceuticals by investigating the substances which act on the receptor. Until now, only several G protein coupled receptor genes or cDNAs have been cloned but it is
35 believed that there are many unknown G protein coupled receptor genes which have not been recognized yet.

The characteristic feature of the G protein coupled receptor proteins which have been known up to now is that seven clusters of hydrophobic amino acid residues are located in the primary structure and pass through (span) the cell membrane at each region thereof. It has been known that such a structure is common among all of the known G protein coupled receptor proteins and further that the amino acid sequences corresponding to the area where the protein passes through the membrane (membrane-spanning region or transmembrane region) and the amino acid sequences near the membrane-spanning region are often highly conserved among the receptors.

When an unknown protein has such a structure, it is strongly suggested that said protein is within a category of the G protein coupled receptor proteins. In addition, some amino acid residue alignments are common (homologous) and, by taking it as a characteristic feature, it is further strongly suggested that said protein is a G protein coupled receptor protein.

Libert, F, et al. (Science, 244:569-571; 1989) reported a method for cloning novel receptor genes by means of a polymerase chain reaction (hereinafter, sometimes referred to as PCR or a PCR technique) for a synthetic DNA primer which was synthesized based upon the information of common amino acid sequences obtained from a comparison among known G protein coupled receptor proteins. Libert, F. et al. used a pair of synthetic DNA primers corresponding to the portions of the third and the sixth membrane-spanning regions. However, in general, the design of primers used for the PCR regulates the molecular species of DNAs which are to be amplified.

In addition, when a similarity (homology) in the amino acid sequence level is used as a basis, the use of different codons affects on the binding (hybridization) of the primer thereby resulting in a decrease in the amplifying efficiency.

Accordingly, although various novel receptor protein DNAs have been obtained using said DNA primers, it is not possible to succeed in amplifying DNAs for all receptor proteins in the prior art.

Further, the amino acid sequence which is common to from the first to the seventh membrane-spanning regions among 74 G protein coupled receptor proteins was reported by William C. Probst, et al. (DNA and Cell Biology, Vol. 11, No. 1, 5 1992, pp. 1-20). In this report, however, there is no suggestion for a method in which DNA coding for a novel G protein coupled receptor protein is screened by means of PCR using DNA primers which are complementary to the DNA coding for those amino acid sequences.

10 It would be desirable to develop DNA primers for PCR techniques which allow selective and efficient screenings of DNAs coding for the areas (regions) more nearer the full length of novel G protein coupled receptor proteins by utilizing the common (homologous) sequence(s) of the G protein coupled 15 receptor protein or the DNA coding therefor.

It would also be desirable to develop synthetic DNA primers corresponding to the portions of the third and the sixth membrane-spanning regions, said primer being useful in screening for DNA coding for G protein coupled receptor 20 proteins in more selective and efficient manner as compared with a series of the synthetic DNA primers corresponding to the sequences of the third to the sixth membrane-spanning regions as reported by Libert, F. et al.

G protein coupled receptor proteins are important for 25 investigating substances which control the function of living organisms and proceeding developments thereof as pharmaceuticals. Finding and development of candidate compounds for new pharmaceuticals can be efficiently proceeded by using G protein coupled receptor proteins and by conducting 30 receptor binding experiments and evaluating experiments on agonists/antagonists using intracellular information transmittance systems as indexes. Especially when the presence of a novel G protein coupled receptor protein can be clarified, the presence of a substance having a specific action thereon 35 can be suggested.

If a novel DNA which codes for a novel G protein coupled receptor protein can be efficiently screened and

isolated, it will now be possible to proceed with the isolation of DNA having an entire coding region, the construction of an expression system therefor and the screening of an acting ligand.

5 A hypothalamo-hypophysial system is one of the passages for controlling, regulating or adjusting the functions of organisms relying upon interactions of hormones and neurotransmitters with G protein coupled receptors. In the hypothalamo-hypophysial system, the secretion of pituitary
10 hormones from the pituitary body (hypophysis) is regulated by hypothalamic hormones (hypophysiotropic releasing factors), and the functions of target cells and organs are controlled by pituitary hormones released into the blood. Functions which are important for the living body are regulated through this
15 system, such as maintenance of homeostasis and control of development and growth of a genital system and an individual organism. Representative examples of the hypothalamic hormones include TRH, LH-RH, CRF, GRF, somatostatin, galanin, etc. Representative examples of the pituitary hormones include TSH,
20 ACTH, FSH, LH, prolactin, growth hormone, oxytocin, vasopressin, etc. In particular, the secretion of pituitary hormones is regulated according to a positive feedback mechanism or a negative feedback mechanism relied on the hypothalamic hormones and peripheral hormones secreted from
25 the target endocrine glands. A variety of receptor proteins present in the pituitary gland play a major role for regulating the hypothalamo-hypophysial system.

 It has been widely known that these hormones, factors and receptors are widely distributed in the brain instead of
30 existing only locally in the hypothalamo-hypophysial system. This fact suggests that the substances which are called "hypothalamic hormones" are working as neurotransmitters or neuroregulators in the central nervous system. It is further considered that these substances are similarly
35 distributed even in the peripheral tissues to play the role of important functions. The pancreas plays an important role of carrying out the carbohydrate metabolism by secreting not only

a digestive fluid but also glucagon and insulin. Insulin is secreted from the β cells and its secretion is promoted chiefly by glucose. It has, however, been known that a variety of receptors exist in the β cells, and the secretion of
5 insulin is controlled by various factors such as peptide hormones (galanin, somatostatin, gastric inhibitory polypeptide, glucagon, amylin, etc.), sugars (mannose, etc.), amino acids, and neurotransmitters in addition to glucose.

It has thus been known that in the pituitary gland
10 and in the pancreas are present receptor proteins for many hormones and neurotransmitters, said receptor proteins playing important roles for regulating the functions. As for the galanin and amylin, however, there has not yet been reported any discovery concerning the structure of their receptor
15 protein cDNAs. It is not known whether there exist any unknown receptor proteins or receptor protein subtypes.

For substances regulating the functions of the pituitary gland and pancreas, there exist receptor proteins specific to said substance on the surfaces of various
20 functional cells of the pituitary gland and pancreas. The pituitary gland and the pancreas are associations of a plurality of functional cells, and the actions of the individual substances are defined by the distributions of their target receptor proteins among the functional cells.
25 Accordingly, a substance, in many cases, exhibits an extensive variety of actions. To comprehend such complex systems, it is necessary to clarify the relations between the acting substances and the specific receptor proteins. It is further necessary to efficiently screen for receptor protein agonists
30 and antagonists capable of regulating the pituitary gland and pancreas, to clarify the structures of genes of receptor proteins from the standpoint of investigating and developing pharmaceuticals, and further to express them in a suitable expression system.

35 By utilizing the fact that a G protein coupled receptor protein exhibits homology in part of the structure thereof at the amino acid sequence level, an experiment of

looking at DNAs coding for novel receptor proteins relying upon a polymerase chain reaction (hereinafter simply referred to as "PCR") has recently been made.

5 In the central nervous system, many receptor proteins such as dopamine receptor protein, LH-RH receptor protein, neurotensin receptor protein, opioid receptor protein, CRF receptor protein, CRF receptor protein, somatostatin receptor protein, galanin receptor protein, TRH receptor protein, etc. are G protein coupled receptor proteins, and it has been
10 clarified that ligands to these receptors exert a variety of effects in the central nervous system.

In the immune system, an α - or a β -chemokine receptor protein, an MIP1 α receptor protein, an IL-8 receptor protein, a C5a receptor protein, etc. have been known as such
15 G protein coupled receptor proteins, and are working as receptor proteins responsive to immunoregulating substances to play important roles for regulating the functions of the living body. There is, for example, an IL-6 receptor protein that acts both in the above-mentioned central nervous system and in
20 the immune system. IL-6 is both a β -cell differentiating factor and a biologically active factor related to the proliferation and differentiation of nerve cells.

It has been widely known that these hormones, factors and receptor proteins are usually widely distributed
25 up to the peripheral tissues instead of existing only locally in the central nervous system and in the immune system and are producing important functions, respectively. Agonists and antagonists for these receptor proteins are now being developed as various useful pharmaceuticals.

30 For substances regulating the functions of the central nervous system and the immune system, there exist receptor proteins specific to said substance on the surfaces of various functional cells of the central nervous system and the immune system. The central nervous system and the immune
35 system are associations of a plurality of functional cells, and the actions of the individual substances are defined by the distributions of their target receptor proteins among the

functional cells. Accordingly, a substance, in many cases, exhibits an extensive variety of actions. Moreover, there is an example wherein many factors play a part in a physiological phenomenon. To comprehend such complex systems, it is necessary to clarify relations between the acting substances and the specific receptor proteins.

As discussed herein above, the G protein coupled receptor protein is present on the cell surface of living body cells and organs and has a very important role as a target for molecules such as hormones, neurotransmitters and physiologically active substances, which molecules control, regulate or adjust the functions of living body cells and organs.

SUMMARY OF THE INVENTION

One object of the present invention is to provide novel DNAs which are useful as DNA primers for a polymerase chain reaction; methods for amplifying a DNA coding for a G protein coupled receptor protein using said DNA; screening methods for the DNA coding for a G protein coupled receptor protein using said DNA; DNAs obtained by said screening method; and G protein coupled receptor proteins encoded by the DNA obtained by said screening method, peptide fragments or segments thereof, modified peptide derivatives thereof or salts thereof.

Another object of the present invention is to provide processes for producing said receptor protein; transformants capable of expressing said receptor protein; cell membrane fractions obtained from said transformant; methods for determining a ligand to the receptor protein; screening methods for a compound or a salt thereof capable of inhibiting the binding of the ligand with the receptor protein; kits for said screening method, pharmaceutical compositions comprising an effective amount of the inhibitory compound; antibodies against said receptor protein; immunoassays using said receptor protein or said antibody and use of said receptor protein and encoding DNA.

Yet another object of the present invention is to provide novel G protein coupled receptor proteins which are expressed in pituitary glands or pancreatic β cells; DNAs comprising a DNA coding for said G protein coupled receptor protein; processes for producing said receptor protein; transformants capable of expressing said receptor protein; cell membrane fractions obtained from said transformant; methods for determining a ligand to the receptor protein; screening methods for a compound or a salt thereof capable of inhibiting the binding of the ligand with the receptor protein; kits for said screening method, pharmaceutical compositions comprising the inhibitory compound; antibodies against said receptor protein; immunoassays using said receptor protein or said antibody and use of said receptor protein and encoding DNA.

Still another object of the present invention is to provide novel human amygdaloid nucleus-derived G protein coupled receptor proteins; DNAs comprising a DNA coding for said G protein coupled receptor protein; processes for producing said receptor protein; transformants capable of expressing said receptor protein; cell membrane fractions obtained from said transformant; methods for determining a ligand to the receptor protein; screening methods for a compound or a salt thereof capable of inhibiting the binding of the ligand with the receptor protein; kits for said screening method, pharmaceutical compositions comprising the inhibitory compound; antibodies against said receptor protein; immunoassays using said receptor protein or said antibody and use of said receptor protein and encoding DNA.

Yet another object of the present invention is to provide novel mouse pancreatic β cell line MIN6-derived G protein coupled receptor proteins; DNAs comprising a DNA coding for said G protein coupled receptor protein; processes for producing said receptor protein; transformants capable of expressing said receptor protein; cell membrane fractions obtained from said transformant; methods for determining a ligand to the receptor protein; screening methods for a

compound or a salt thereof capable of inhibiting the binding of the ligand with the receptor protein; kits for said screening method, pharmaceutical compositions comprising the inhibitory compound; antibodies against said receptor protein; 5 immunoassays using said receptor protein or said antibody and use of said receptor protein and encoding DNA.

The present inventors have succeeded in synthesizing novel DNA primers based upon the similarity (homology) with the base sequences coding for the first membrane-spanning region 10 or the sixth membrane-spanning region each of known G protein coupled receptor proteins. It is to be particularly noted that there has been no report of a DNA primer pair which has been synthesized paying attention to the similarity with the base sequence coding for the first and the sixth membrane-spanning 15 region of the known G protein coupled receptor protein.

Next the present inventors have succeeded in synthesizing other novel DNA primers based upon the similarity (homology) with the base sequences coding for the third or the sixth membrane-spanning region each of known G protein 20 coupled receptor proteins. They have also unexpectedly succeeded in efficiently amplifying DNAs (DNA fragments) coding for G protein coupled receptor proteins by means of PCR using those DNA primers.

They have further succeeded in synthesizing novel 25 DNA primers based upon the similarity (homology) with the base sequences coding for the second or the seventh membrane-spanning region each of known G protein coupled receptor proteins; upon the similarity (homology) with the base sequences coding for first or the third membrane-spanning 30 region each of known G protein coupled receptor proteins; and upon the similarity (homology) with the base sequences coding for the second or the sixth membrane-spanning region each of known G protein coupled receptor proteins. They have furthermore and unexpectedly succeeded in efficiently 35 amplifying DNAs (DNA fragments) coding for G protein coupled receptor proteins by conducting PCR using those DNA primers.

Moreover, the present inventors have succeeded in

efficiently cloning full-length DNA coding for said G protein coupled receptor protein via using amplified DNAs (DNA fragments) coding for said G protein coupled receptor protein. Thus, they have found that novel DNA coding for novel G protein coupled receptor proteins can be isolated, characterized or prepared via conducting amplifications and analyses of various DNA using said DNA primers.

To be more specific, the present inventors have selected amino acid sequences which are each common to the portion corresponding to or near the first and the sixth membrane-spanning region of the known individual G protein coupled receptor proteins and have designed the DNA primer (SEQ ID NO: 1) coding for the amino acid sequence common (homologous) to the first membrane-spanning region and the DNA primer (SEQ ID NO: 2) which is complementary to the nucleotide sequence coding for the amino acid sequence common (homologous) to the area near the sixth membrane-spanning region. Those DNA primers have a different nucleotide sequence as compared with reported DNA primers (e.g. a set of synthetic DNA primers corresponding to the third and the sixth membrane-spanning regions (SEQ ID NO: 60 and SEQ ID NO: 61) as reported by Libert, F. et al.) and such instant primers are novel and unique.

Especially for an object of conducting an efficient elongation reaction in the PCR, the 3'-terminal region of the instant primers contains the nucleotide sequence which is common (homologous) among many receptor proteins. Even in other areas, the similarity (homology) at the nucleotide sequence level (base sequence level) is utilized for setting the mixed base (nucleotide) parts wherein their nucleotide sequences (base sequences) are matched for as many nucleotides (bases) as possible among many DNA for the receptor proteins. Then the present inventors have amplified cDNA derived from human brain amygdala, human pituitary gland and rat brain, found the amplified products as shown in Figure 17 and, from those products, obtained the G protein coupled receptor protein cDNAs having the sequence as shown in

Figure 18, Figure 19, Figure 20, Figure 21, Figure 22,
Figure 23, Figure 27, Figure 29, Figure 34, Figure 37,
Figure 40, Figure 43 or Figure 46. Among them, the G protein
coupled receptor protein cDNAs having the sequence as shown in
5 Figure 22, Figure 23, Figure 27, Figure 29, Figure 34,
Figure 37, Figure 40, Figure 43 or Figure 46 are novel.

Further, the present inventors have selected the
amino acid sequences common (homologous) to the third and the
sixth membrane-spanning region each of the known G protein
10 coupled receptor proteins and designed the DNA primers coding
for the amino acid sequence common (homologous) to the third
membrane-spanning region (SEQ ID NO: 3; SEQ ID NO: 5, SEQ ID
NO: 6 and SEQ ID NO: 7) and the DNA primers which are
complementary to the nucleotide sequence coding for the amino
15 acid sequence common (homologous) to the portion near the sixth
membrane-spanning region (SEQ ID NO: 4, SEQ ID NO: 8 and SEQ ID
NO: 9). Again, those DNA primers have different base sequences
from those of the DNA primers previously reported (e.g., a set
of synthetic DNA primers corresponding to the sequence of the
20 third and the sixth membrane-spanning regions (SEQ ID NO: 60
and SEQ ID NO: 61) as reported by Libert, F. et al.) and such
instant primers are novel and unique. The present inventors
amplified cDNA derived from the smooth muscles of gastric
pylorus of rabbits using said DNA primer and obtained G protein
25 coupled receptor protein cDNA having the sequence of Figure 49
or Figure 52. Those cDNAs are novel.

Still further, the present inventors have selected
the amino acid sequences common (homologous) to the second and
the seventh membrane-spanning region each of the known G
30 protein coupled receptor proteins and designed the DNA primer
coding for the amino acid sequence common (homologous) to the
second membrane-spanning region (SEQ ID NO: 10) and the DNA
primer which is complementary to the base sequence coding for
the amino acid sequence common (homologous) to the portions
35 near the seventh membrane-spanning region (SEQ ID NO: 11).
Those DNA primers have different base sequences from those of
DNA primers previously reported (e.g., a set of synthetic DNA

primers corresponding to the part of the third and the sixth membrane-spanning regions (SEQ ID NO: 60 and SEQ ID NO: 61) as reported by Libert, F. et al) and such instant primers are novel and unique. The present inventors amplified cDNA derived from the smooth muscles of gastric pylorus of rabbits using said DNA primer and obtained G protein coupled receptor protein cDNAs having each the sequence of Figure 55, Figure 56, Figure 72, or Figure 73. Those cDNAs are novel.

Furthermore, the present inventors have selected the amino acid sequences common (homologous) to the first and the third membrane-spanning region each of the known G protein coupled receptor proteins and designed the DNA primer coding for the amino acid sequence common (homologous) to the first membrane-spanning region (SEQ ID NO: 12) and the DNA primer which is complementary to the base sequence coding for the amino acid sequence common (homologous) to the portions near the third membrane-spanning region (SEQ ID NO: 13).

Still further, the present inventors have selected the amino acid sequences common (homologous) to the third and the sixth membrane-spanning region each of the known G protein coupled receptor proteins and designed the DNA primers coding for the amino acid sequence common (homologous) to the third membrane-spanning region (SEQ ID NO: 10 and SEQ ID NO: 18) and the DNA primers which are complementary to the base sequence coding for the amino acid sequence common (homologous) to the parts near the sixth membrane-spanning region (SEQ ID NO: 15 and SEQ ID NO: 19). Further, the present inventors have selected the amino acid sequences common (homologous) to the second and the sixth membrane-spanning region each of the known G protein coupled receptor proteins and designed the DNA primer coding for the amino acid sequence common (homologous) to the second membrane-spanning region (SEQ ID NO: 16) and the DNA primer which is complementary to the base sequence coding for the amino acid sequence common (homologous) to the parts near the sixth membrane-spanning region (SEQ ID NO: 17). Those DNA primers have different base sequences from those of DNA primers previously reported (e.g., a set of synthetic DNA primers

corresponding to the part of the third and the sixth membrane-spanning regions (SEQ ID NO: 60 and SEQ ID NO: 61) as reported by Libert, F. et al) and such instant primers are novel and unique.

5 Still another object of the present invention is to provide a G protein coupled receptor protein expressed in the pituitary gland and pancreatic β cells, a DNA comprising a DNA coding for said protein, a process for producing said protein, and use of said protein and DNA.

10 In order to achieve the above-mentioned aims, the present inventors have made extensive investigations. As a result, the present inventors have succeeded in amplifying cDNA derived from the human pituitary gland and the mouse pancreatic β -cell strain, MIN 6, with a synthetic DNA primer
15 for efficiently isolating G protein coupled receptor protein-encoding DNA, and have forwarded the analysis. Thus, the present inventors have succeeded in isolating novel human and mouse-derived G protein coupled receptor protein-encoding cDNAs, in determining the partial structure thereof, and have
20 considered that these cDNA sequences are preserved very well in the human and in the mouse, and are coding for novel receptor proteins for the same ligand. Based upon the above knowledge, the present inventors have discovered that these DNAs make it possible to obtain a cDNA having a full length
25 open reading frame (ORF) of the receptor protein, hence, to produce the receptor protein. The inventors have further discovered that the above-mentioned receptor protein obtained when the G protein coupled receptor protein-encoding cDNA is expressed by a suitable means permits screening for a ligand to
30 the receptor protein from the living body or from natural or non-natural compounds under guidance of data obtainable in receptor coupling tests or measurements of intracellular second messengers, etc. and further allows screening for a compound that inhibits the binding of the ligand and the
35 receptor protein.

 In one embodiment, the present inventors have carried out PCR amplification of novel human pituitary gland-

derived cDNA fragments as shown in Figures 22 and 23, and have subcloned them to obtain a plasmid vector (p19P2).

From analysis of the partial sequence, it has been clarified that the cDNA has been encoded a novel receptor protein.

5 The synthetic DNA primers used for amplifying the cDNA are corresponding to seven hydrophobic clusters that exist in the known G protein coupled receptor proteins in common, i.e., corresponding to the first and sixth membrane-spanning regions among the membrane-spanning domains. The nucleotide sequence
10 (SEQ ID NO: 29) has been determined from the primer region at the 5' side (first membrane-spanning domain side) and has been translated into an amino acid sequence (SEQ ID NO: 24)

[Figure 22]. As a result, the second and third membrane-spanning domains have been confirmed on the hydrophobicity
15 plotting [Figure 58]. Similarly, the nucleotide sequence (SEQ ID NO: 30) has been determined from the primer region at the 3' side (sixth membrane-spanning domain side) and has been translated into an amino acid sequence (SEQ ID NO: 25)

[Figure 23]. As a result, the presence of the sixth and fifth
20 membrane-spanning domains has been confirmed on the hydrophobicity plots [Figure 59]. The size of the amplified cDNA is about 700 bp which is nearly comparable with the number of bases between the first membrane-spanning domain and the sixth membrane-spanning domain of the known G protein coupled,
25 receptor protein.

G protein coupled receptor proteins exert common property to some extent at an amino acid sequence level, and are forming one protein family. Therefore, data base retrieval has been carried out based upon the amino acid sequence of the
30 subject novel receptor protein (protein encoded by cDNA included in p19P2). As a result, a high homology has been exhibited as compared with the known G protein coupled receptor protein (rat neuropeptide Y receptor protein encoded by S12863) that is shown in Figure 60. This fact tells that
35 the novel receptor protein of the present invention belongs to the G protein coupled receptor protein family. Moreover, the data base has been retrieved using, as a template, the amino

acid sequence encoded by the DNA of the invention. It exhibits high homology to the amino acid sequences of the known G protein coupled receptor proteins, mouse-derived ligand unknown RP-23 (B40470), human-derived ligand unknown K-opioid receptor protein (P30098) and human-derived NK-2 receptor protein (JQ1059). However, none of them are in perfect agreement, from which it is learned that a novel receptor protein had been encoded. The aforementioned abbreviations in parentheses are reference numbers that are assigned when they are registered as data to NBRF-PIR/Swiss-PROT and are, usually, each called "Accession Number".

Next, by using the novel G protein coupled receptor protein-encoding cDNA fragment (p19P2) of the present invention, a cDNA having a full-length open reading frame of the receptor protein of the present invention has been obtained from human pituitary gland cDNA libraries. The nucleotide sequence analysis of a plasmid (phGR3) carrying the cDNA having a full length open reading frame of the receptor protein shows that the nucleotide sequence of a coding region of this receptor protein is represented by SEQ ID NO: 31, and the amino acid sequence deduced therefrom is represented by SEQ ID NO: 26 [Figure 34]. Based upon the amino acid sequence, hydrophobicity plotting has been carried out. The results are shown in Figure 36. From the hydrophobicity plotting, it has been clarified that the receptor protein of the present invention possessed seven hydrophobic domains. That is, it has been confirmed that the receptor protein encoded by the cDNA obtained according to the present invention is a seven transmembrane (membrane-spanning) G protein coupled receptor protein. An expression of mRNA for receptor genes encoded by the cDNA of the present invention has been checked by northern blotting techniques at a mRNA level, and it has been confirmed that the receptor gene has been expressed in the human pituitary gland [Figure 35].

The present inventors have further succeeded in PCR amplification of a mouse pancreatic β cell strain, MIN6 derived cDNA fragment, and cloning of pG3-2 and pG1-10.

- 17 -

Then, based on the nucleotide sequence of cDNA included in these two plasmid vectors, the nucleotide sequence shown in Figure 27 has been derived. It was learned from the nucleotide sequence that the cDNA encodes a novel receptor protein. Upon translating the nucleotide sequence into an amino acid sequence, the presence of the third, fourth, fifth and sixth membrane-spanning domains has been confirmed on the hydrophobicity plots [Figure 28]. The size of the amplified cDNA is about 400 bp which is nearly comparable with the number of bases between the third membrane-spanning domain and the sixth membrane-spanning domain of the known G protein coupled receptor protein. The amino acid sequence has been compared with amino acid sequences [Figures 22 and 23] encoded by the G protein coupled receptor protein cDNA included in p19P2 cloned from the human pituitary gland. As a result, homology is more than 95% [Figure 61]. From this fact, it was estimated that the protein encoded by the cDNA included in pG3-2 is a mouse derived one encoded by the cDNA included in p19P2.

The present inventors have further amplified a mouse pancreatic β -cell strain, MIN6-derived cDNA fragment by the PCR followed by subcloning into a plasmid vector as shown in Figure 62. From the nucleotide sequence (SEQ ID NO: 33), it has been clarified that the cDNA encodes a novel receptor protein. Upon translating the nucleotide sequence into an amino acid sequence (SEQ ID NO: 28), the presence of the third, fourth, fifth and sixth membrane-spanning domains has been confirmed on the hydrophobicity plots [Figure 64]. The size of the amplified DNA is about 400 bp that is nearly comparable with the known G protein coupled receptor protein. The amino acid sequence has been compared with amino acid sequences [Figures 22 and 23] encoded by the G protein coupled receptor protein cDNA included in p19P2 cloned from the human pituitary gland and with amino acid sequences of proteins encoded by pG3-2 and pG1-10 derived from the mouse pancreatic β -cell strain. As a result, homology is more than 95% to them

[Figure 63]. This fact suggests that the protein encoded by the human-derived pituitary gland-derived p19P2, the proteins encoded by the mouse pancreatic β -cell strain-derived pG3-2 and pG1-10, and the protein encoded by the mouse pancreatic β -cell strain-derived p5S38, pertain to a receptor family that recognizes the same ligand.

Another object of the present invention is to provide a novel human amygdaloid nucleus-derived protein coupled receptor protein, a DNA containing a DNA coding for said G protein coupled receptor protein, a process for producing said G protein coupled receptor protein, and use of said protein and DNA.

The present inventors have synthesized DNA primers for efficiently isolating a DNA coding for G protein coupled receptor proteins, amplified an amygdaloid nucleus-derived cDNA with the above primer, and have analyzed it.

As a result, the present inventors have succeeded in isolating, from the human amygdaloid nucleus, a cDNA coding for a novel G protein coupled receptor protein and have determined its partial structure. The nucleotide sequence of the isolated cDNA is preserved very well as compared with that of the mouse glucocorticoid-induced receptor (hereinafter sometimes referred to as "GIR") and is considered to be encoding a receptor protein to the same ligand (Molecular Endocrinology 5:1331-1338, 1991). It is reputed that, in the mouse, the GIR is a receptor which is induced by glucocorticoid and expressed in T-cells and is working as a receptor to immunoregulating factors in the immune system on the T-cells. The present inventors have succeeded in the isolation of this human type GIR from the human amygdaloid nucleus. Accordingly, it is suggested that the isolated GIR is expressed even in the human central nervous system to carry out some function. From these facts, it is considered that the receptor protein is strongly expressed in the human brain and in the immune system and is also functioning therein. These characterized DNAs allow one to obtain a cDNA having a full length open reading frame of the receptor and production of the receptor

proteins. The receptor proteins expressed by a suitable means, furthermore, permit screening for a ligand to the receptor proteins from the living body or from natural and non-natural compounds depending on indications obtainable in receptor protein-binding experiments, measurements of intracellular second messengers, etc. It further allows one to screen for compounds capable of inhibiting the binding between the ligand and the receptor protein.

To be more specific, the present inventors have amplified, as a novel human amygdaloid nucleus-derived cDNA, one species, as shown in Figures 29 and 30, by PCR, cloned it, and clarified from the analysis of a partial sequence thereof that a novel receptor protein is encoded. The synthetic DNA primers used for amplifying the cDNA are corresponding to seven hydrophobic clusters that exist in the G protein coupled receptor proteins in common, i.e., corresponding to the first and sixth membrane-spanning regions among the membrane-spanning domains. The nucleotide sequence has been determined from the primer region at the 5' side (first membrane-spanning domain side) and has been translated into an amino acid sequence. As a result, the second and third membrane-spanning domains have been confirmed on the hydrophobicity plotting [Figure 31]. Similarly, the nucleotide sequence has been determined from the primer region at the 3' side (sixth membrane-spanning domain side) and has been translated into an amino acid sequence. As a result, the presence of the fifth and fourth membrane-spanning domains has been confirmed on the hydrophobicity plots [Figure 32]. The size of the amplified cDNA is about 700 bp which is nearly comparable with the number of bases of the known G protein coupled receptor protein.

The inventors have further retrieved the data base based on, as a template, the nucleotide sequence of the isolated DNA and observed high homology to the DNA that codes for mouse-derived glucocorticoid-induced receptor protein which is a widely known G protein coupled receptor protein [Figure 33]. This result strongly suggests that the DNA of the present invention is encoding a human-type receptor protein of GIR.

Yet another object of the present invention is to provide a novel mouse pancreatic β -cell strain, MIN6-derived protein coupled receptor protein, a DNA containing a DNA coding for said G protein coupled receptor protein, a process
5 for producing said G protein coupled receptor protein, and use of said protein and DNA. The present inventors have synthesized DNA primers for efficiently isolating a DNA coding for G protein coupled receptor proteins, amplified a mouse pancreatic β -cell strain, MIN6-derived cDNA with the above
10 primer, and have analyzed it.

As a result, the present inventors have succeeded in isolating a mouse-derived cDNA coding for a novel G protein coupled receptor protein and have determined its partial structure. The isolated cDNA is homologous to known G protein
15 coupled receptors at the nucleotide sequence level and at the amino acid sequence level and is considered to be encoding a novel receptor protein which is expressed in the mouse pancreas and is also functioning therein. These characterized DNAs allow one to obtain a cDNA having a full length open reading
20 frame of the receptor and production of the receptor proteins. Human-derived cDNAs may be cloned by using, as a probe, said mouse-derived cDNA. The receptor proteins expressed by a suitable means, furthermore, permit screening for a ligand to the receptor protein from the living body or
25 from natural and non-natural compounds relying on indications obtainable in receptor protein-binding experiments, measurements of intracellular second messengers, etc. It further allows one to screen for compounds capable of inhibiting the binding of the ligand with the receptor protein.

To be more specific, the present inventors have amplified, as a novel mouse pancreatic β -cell strain, MIN6-derived cDNA, p3H2-17, as shown in Figures 37, by PCR, cloned it, and clarified from the analysis of a partial sequence
30 thereof that a novel receptor protein is encoded. The nucleotide sequence has been translated into an amino acid sequence. As a result, the presence of the third, fourth, fifth and sixth membrane-spanning domains has been confirmed
35

- 21 -

on the hydrophobicity plots [Figure 38]. The size of the amplified cDNA is about 400 bp which is nearly comparable with that of the known G protein coupled receptor protein.

5 The inventors have retrieved the data base based on, as a template, the nucleotide sequence of the isolated DNA and observed 30% homology to chicken ATP receptor (P34996), 25% homology to human somatostatin receptor subtype 3 (A46226), 27% homology to bovine neuropeptide Y receptor (S28787), and 28% homology to human somatostatin receptor subtype 4 (JN0605), respectively (Figure 39), which are known G protein coupled
10 receptor proteins. The aforementioned abbreviations in parentheses are reference numbers that are assigned when they are registered as data to NBRF-PIR/Swiss-PROT and are, usually, each called "Accession Number".

15 An expression of receptor genes encoded by the cDNA fragment included in p3H2-17 of the present invention has been checked by northern blotting techniques at a mRNA level, and it has been confirmed that the receptor gene has been intensely expressed in the mouse thymus and spleen. It has been also confirmed that the receptor gene has been expressed in the mouse brain and pancreas (Figure 65).

20 Next, by utilizing the information on the nucleotide sequence of the fragment included in p3H2-17, cDNA encoding a full-length open reading frame of the mouse pancreatic β -cell strain, MIN6-derived G protein coupled receptor protein of the present invention has been obtained from mouse
25 thymic and splenic poly(A)⁺ RNA by 5'RACE (5' rapid amplification of cDNA ends) techniques (Frohman M.A. et al., Proc. Natl. Acad. Sci. USA, 85:8998-9002 (1988); Edwards et al., Nucleic Acids Res., 17:2919-2932 (1989); Belyavsky A. et al., Nucleic Acids Res., 19:5227-5232 (1991)) and
30 3'RACE (3' rapid amplification of cDNA ends) techniques (Frohman M.A. et al., Proc. Natl. Acad. Sci. USA, 85:8998-9002 (1988); Belyavsky A. et al., Nucleic Acids Res., 17:2919-2932 (1989)).

35 The plasmid (pMAH2-17) carrying cDNA encoding a full-length open reading frame of the receptor protein of the

present invention has been subjected to sequencing analysis. As a result, the nucleotide sequence of the region coding for the receptor protein is represented by SEQ ID NO: 41 and the amino acid sequence deduced therefrom is represented by SEQ ID
5 NO: 39 (Figure 69). Based on the amino acid sequence, hydrophobicity plotting has been carried out. The results are shown in Figure 70.

It has been clarified from the hydrophobicity plotting that the mouse pancreatic β -cell strain, MIN6-
10 derived receptor protein of the present invention has seven hydrophobic domains. Thus, it has been confirmed that the receptor protein encoded by the cDNA included in pMAH2-17 according to the present invention is a seven transmembrane G protein coupled receptor protein.

15 Data base retrieval has been carried out based on the full-length amino acid sequence encoded by the cDNA included in pMAH2-17, and it has been observed that the amino acid sequence has 44.0% homology to mouse P_{2U} purinoceptor (P35383) and 38.1% homology to chicken
20 P_{2Y} purinoceptor (P34996), respectively (Figure 71), which are known G protein coupled receptor proteins. The aforementioned abbreviations in parentheses are reference numbers that are assigned when they are registered as data to NBRF-PIR/Swiss-PROT and are, usually, each called "Accession Number".
25 Since the receptor protein encoded by pMAH2-17 is highly homologous to prinoceptors, it is considered that there are strong possibility of a subtype within prinoceptor families. Therefore, the present inventors have carried out an electrophysiological analysis of the receptor gene in Xenopus
30 oocytes and found significant inward currents elicited by Xenopus oocytes carrying the subject receptor gene in response to ATP stimulation (Figure 75). As a result, it has been determined that the receptor encoded by pMAH2-17 is one of the subtypes within prinoceptor families. It has been
35 discussed and expected that there are a variety of subtypes among purinoceptors (Pharmac. Ther., Vol. 64, pp. 445-475 (1994)).

All data are supporting that the mouse pancreatic β -cell strain, MIN6-derived receptor protein of the present invention (e.g., SEQ ID NO: 38 and SEQ ID NO: 39, or proteins encoded by pMAH2-17) is a novel purinoceptor subtype which is
5 clearly distinct from chicken P_{2y1} purinoceptor (FEBS LETTERS, Vol. 324(2), 219-225 (1993)); mouse P_{2y2} or P_{2u} purinoceptor (Proc. Natl. Acad. Sci. USA, Vol. 90, pp.5113-5117 (1993)); rat P_{2u} or P_{2y2} purinoceptor (Am. J. Respir. Cell Mol. Biol., Vol. 12, pp. 27-32 (1995)); human P_{2u} or P_{2y2} purinoceptor (Proc.
10 Natl. Acad. Sci. USA, Vol. 91, pp.3275-3279 (1994)); and rat P_{2x} purinoceptor (Nature, Vol. 371.6, pp.516-519 (1994)).

It is also strongly suggested that agonists and/or antagonists related to the receptor encoded by pMAH2-17 would be useful in therapeutic or prophylactic treatment of
15 diseases or syndromes in connection with purine ligand compounds. It is expected that the agonists of the receptor encoded by pMAH2-17 are useful as an immunomodulator or an antitumor agent, in addition they are useful in therapeutically or prophylactically treating hypertension, diabetes, cystic
20 fibrosis, etc. It is still expected that the antagonists of the receptor encoded by pMAH2-17 are useful as hypotensive agents, analgesics, agents for therapeutically or prophylactically treating incontinence of urine, etc.

Another object of the present invention is to
25 provide a novel human-derived protein coupled receptor protein of prinoceptor type, a DNA containing a DNA coding for said G protein coupled receptor protein, a process for producing said G protein coupled receptor protein, and use of said protein and DNA. The present inventors have synthesized DNA primers for
30 efficiently isolating a DNA coding for prinoceptor type G protein coupled receptor proteins on the basis of the nucleotide sequence of mouse purinoceptor, amplified a human-derived cDNA with the above primer, and have analyzed it.

As a result, the present inventors have succeeded in
35 isolating a human-derived cDNA coding for a novel G protein coupled receptor protein and have determined its full-length structure [Figure 77]. The isolated cDNA is homologous to

mouse G protein coupled receptor (purinoceptor) at the nucleotide sequence level and at the amino acid sequence level (87% homology; Figure 79) and is considered to be encoding a novel purinoceptor protein. The receptor proteins expressed by a suitable means, furthermore, permit screening for a ligand to the receptor protein from the living body or from natural and non-natural compounds relying on indications obtainable in receptor protein-binding experiments, etc. It further allows one to screen for compounds capable of inhibiting the binding of the ligand with the receptor protein.

It is also strongly suggested that agonists and/or antagonists related to the human receptor encoded by phAH2-17 would be useful in therapeutic or prophylactic treatment of diseases or syndromes in connection with purine ligand compounds. It is expected that the agonists of the human receptor are useful as an immunomodulator or an antitumor agent, in addition they are useful in therapeutically or prophylactically treating hypertension, diabetes, cystic fibrosis, etc. It is still expected that the antagonists of the human receptor are useful as hypotensive agents, analgesics, agents for therapeutically or prophylactically treating incontinence of urine, etc.

Accordingly, one aspect of the present invention is

(1) DNAs comprising a nucleotide sequence represented by a SEQ ID NO selected from the group consisting of SEQ ID NO: 1 to SEQ ID NO: 19;

(2) DNAs according to the above (1) comprising a nucleotide sequence represented by a SEQ ID NO selected from the group consisting of SEQ ID NO: 1 to SEQ ID NO: 9;

(3) DNAs according to the above (1) comprising a nucleotide sequence represented by SEQ ID NO: 1 or SEQ ID NO: 2;

(4) DNAs according to the above (1) wherein the DNA is a primer for polymerase chain reaction in order to amplify a DNA coding for a G protein coupled receptor protein;

(5) a method for amplifying a DNA coding for a G

protein coupled receptor protein by polymerase chain reaction techniques, which comprises:

(i) carrying out a polymerase chain reaction in the presence of a mixture of

- ① a DNA coding for G protein coupled receptor protein, said DNA being capable of acting as a template,
- ② at least one DNA primer selected from the group consisting of DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 1, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 3, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 5, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 6, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 7, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 10, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 14, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 16 and DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 18, and
- ③ at least one DNA primer selected from the group consisting of DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 2, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 4, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 8, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 9, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 11, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 15, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 17 and DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 19; or

(ii) carrying out a polymerase chain reaction in the presence of a mixture of

- ① a DNA coding for G protein coupled receptor protein, said DNA being capable of acting as a template,

- ② at least one DNA primer selected from the group consisting of DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 1 and DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 12, and
- ③ at least one DNA primer selected from the group consisting of DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 13;
- (6) a method for screening a DNA library for a DNA coding for a G protein coupled receptor protein, which comprises:
- (i) carrying out a polymerase chain reaction in the presence of a mixture of
- ① said DNA library,
- ② at least one DNA primer selected from the group consisting of DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 1, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 3, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 5, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 6, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 7, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 10, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 14, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 16 and DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 18, and
- ③ at least one DNA primer selected from the group consisting of DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 2, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 4, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 8, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 9, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 11, DNA primers comprising a nucleotide

sequence represented by SEQ ID NO: 15, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 17 and DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 19,

5 to amplify selectively a template DNA coding for G protein coupled receptor protein, contained in the DNA library; or
(ii) carrying out a polymerase chain reaction in the presence of a mixture of

① said DNA library

10 ② at least one DNA primer selected from the group consisting of DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 1 and DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 12, and

15 ③ at least one DNA primer selected from the group consisting of DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 13,

to amplify selectively a DNA coding for G protein coupled receptor protein, contained in the DNA library;

20 (7) a DNA coding for a G protein coupled receptor protein, which is obtained by a method according to the above (5) or (6); and

(8) G protein coupled receptor proteins encoded by a DNA according to the above (7), their peptide segments or
25 fragments and salts thereof.

Another specific aspect of the invention is:

(9) a method for amplifying a DNA coding for G protein coupled receptor protein (e.g. a region corresponding to from the first to sixth membrane-spanning domains of G
30 protein coupled receptor proteins or other domains thereof) by polymerase chain reaction techniques, which comprises carrying out a polymerase chain reaction in the presence of a mixture of

① a DNA coding for G protein coupled receptor protein, said DNA being capable of acting as a template,

35 ② at least one DNA primer selected from the group consisting of DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 1 and DNA primers

comprising a nucleotide sequence represented by SEQ ID NO: 12, and

- 5 ③ at least one DNA primer selected from the group consisting of DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 2, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 4, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 8, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 9, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 15, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 17 and DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 19;

- 15 (10) a method for amplifying a DNA coding for G protein coupled receptor protein (e.g. a region corresponding to from the first to seventh membrane-spanning domains of G protein coupled receptor proteins or other domains thereof) by polymerase chain reaction techniques, which comprises carrying out a polymerase chain reaction in the presence of a mixture of

- 20 ① a DNA coding for G protein coupled receptor protein, said DNA being capable of acting as a template,
- ② at least one DNA primer selected from the group consisting of DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 1 and DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 12, and
- 25 ③ at least one DNA primer selected from the group consisting of DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 11;

- 30 (11) a method for amplifying a DNA coding for G protein coupled receptor protein (e.g. a region corresponding to from the third to sixth membrane-spanning domains of G protein coupled receptor proteins or other domains thereof) by polymerase chain reaction techniques, which comprises carrying out a polymerase chain reaction in the presence of a mixture of

- 35 ① a DNA coding for G protein coupled receptor protein,

said DNA being capable of acting as a template,

- ② at least one DNA primer selected from the group consisting of DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 3, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 5, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 6, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 7, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 14 and DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 18, and

- ③ at least one DNA primer selected from the group consisting of DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 2, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 4, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 8, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 9, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 15, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 17 and DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 19;

(12) a method for amplifying a DNA coding for G protein coupled receptor protein (e.g. a region corresponding to from the third to seventh membrane-spanning domains of G protein coupled receptor proteins or other domains thereof) by polymerase chain reaction techniques, which comprises carrying out a polymerase chain reaction in the presence of a mixture of

- ① a DNA coding for G protein coupled receptor protein, said DNA being capable of acting as a template,
- ② at least one DNA primer selected from the group consisting of DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 3, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 5, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 6, DNA primers comprising a

nucleotide sequence represented by SEQ ID NO: 7, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 14 and DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 18, and

- 5 ③ at least one DNA primer selected from the group consisting of DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 11;

(13) a method for amplifying a DNA coding for G protein coupled receptor protein (e.g. a region corresponding to from the second to sixth membrane-spanning domains of G protein coupled receptor proteins or other domains thereof) by polymerase chain reaction techniques, which comprises carrying out a polymerase chain reaction in the presence of a mixture of

- 10 ① a DNA coding for G protein coupled receptor protein, said DNA being capable of acting as a template,
15 ② at least one DNA primer selected from the group consisting of DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 10 and DNA primers comprising a nucleotide sequence represented by SEQ ID
20 NO: 16, and
 ③ at least one DNA primer selected from the group consisting of DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 2, DNA primers comprising a nucleotide sequence represented by SEQ ID
25 NO: 4, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 8, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 9, DNA primers comprising a nucleotide sequence represented by
30 SEQ ID NO: 15, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 17 and DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 19;

(14) a method for amplifying a DNA coding for G protein coupled receptor protein (e.g. a region corresponding to from the second to seventh membrane-spanning domains of G protein coupled receptor proteins or other domains thereof) by polymerase chain reaction techniques, which comprises carrying

out a polymerase chain reaction in the presence of a mixture of

- ① a DNA coding for G protein coupled receptor protein,
said DNA being capable of acting as a template,
- ② at least one DNA primer selected from the group
consisting of DNA primers comprising a nucleotide
sequence represented by SEQ ID NO: 10 and DNA primers
comprising a nucleotide sequence represented by SEQ ID
NO: 16, and
- ③ at least one DNA primer selected from the group
consisting of DNA primers comprising a nucleotide
sequence represented by SEQ ID NO: 11;

(15) a method for amplifying a DNA coding for G
protein coupled receptor protein (e.g. a region corresponding
to from the first to third membrane-spanning domains of G
protein coupled receptor proteins or other domains thereof) by
polymerase chain reaction techniques, which comprises carrying
out a polymerase chain reaction in the presence of a mixture of

- ① a DNA coding for G protein coupled receptor protein,
said DNA being capable of acting as a template,
- ② at least one DNA primer selected from the group
consisting of DNA primers comprising a nucleotide
sequence represented by SEQ ID NO: 1 and DNA
primers comprising a nucleotide sequence
represented by SEQ ID NO: 12, and
- ③ at least one DNA primer selected from the group
consisting of DNA primers comprising a nucleotide
sequence represented by SEQ ID NO: 13;

(16) a method for amplifying a DNA coding for G
protein coupled receptor protein by polymerase chain reaction
techniques, which comprises carrying out a polymerase chain
reaction in the presence of a mixture of

- ① a DNA coding for G protein coupled receptor protein,
said DNA being capable of acting as a template,
- ② at least one DNA primer selected from the group
consisting of DNA primers comprising a nucleotide
sequence represented by SEQ ID NO: 1, and
- ③ at least one DNA primer selected from the group

consisting of DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 2;

5 (17) a method for amplifying a DNA coding for G protein coupled receptor protein by polymerase chain reaction techniques, which comprises carrying out a polymerase chain reaction in the presence of a mixture of

- ① a DNA coding for G protein coupled receptor protein, said DNA being capable of acting as a template,
- 10 ② at least one DNA primer selected from the group consisting of DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 3, and
- ③ at least one DNA primer selected from the group consisting of DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 4;

15 (18) a method for amplifying a DNA coding for G protein coupled receptor protein by polymerase chain reaction techniques, which comprises carrying out a polymerase chain reaction in the presence of a mixture of

- 20 ① a DNA coding for G protein coupled receptor protein, said DNA being capable of acting as a template,
- ② at least one DNA primer selected from the group consisting of DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 6, and
- 25 ③ at least one DNA primer selected from the group consisting of DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 8;

30 (19) a method for amplifying a DNA coding for G protein coupled receptor protein by polymerase chain reaction techniques, which comprises carrying out a polymerase chain reaction in the presence of a mixture of

- ① a DNA coding for G protein coupled receptor protein, said DNA being capable of acting as a template,
- 35 ② at least one DNA primer selected from the group consisting of DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 10, and
- ③ at least one DNA primer selected from the group consisting of DNA primers comprising a nucleotide

sequence represented by SEQ ID NO: 11;

(20) a method for amplifying DNA coding for a G protein coupled receptor protein which comprises

(i) carrying out a polymerase chain reaction in the presence
5 of a mixture of

- ① a DNA coding for G protein coupled receptor protein,
said DNA being capable of acting as a template,
- ② at least one DNA primer which is capable of
10 binding with the 3'-side nucleotide sequence of
the - chain (minus chain) of the template DNA
coding for G protein coupled receptor protein to
allow the extension of the + chain (plus chain)
in the 5' → 3' direction, said DNA primer being
selected from the group consisting of DNA primers
15 comprising a nucleotide sequence represented by SEQ ID
NO: 1, DNA primers comprising a nucleotide sequence
represented by SEQ ID NO: 3, DNA primers comprising a
nucleotide sequence represented by SEQ ID NO: 5, DNA
primers comprising a nucleotide sequence represented by
20 SEQ ID NO: 6, DNA primers comprising a nucleotide
sequence represented by SEQ ID NO: 7, DNA primers
comprising a nucleotide sequence represented by
SEQ ID NO: 10, DNA primers comprising a
nucleotide sequence represented by SEQ ID NO: 12,
25 DNA primers comprising a nucleotide sequence
represented by SEQ ID NO: 14, DNA primers
comprising a nucleotide sequence represented by
SEQ ID NO: 16 and DNA primers comprising a
nucleotide sequence represented by SEQ ID NO: 18,
30 and
- ③ at least one DNA primer which is capable of
binding with the 3'-side nucleotide sequence of
the + chain (plus chain) of the template DNA
coding for G protein coupled receptor protein to
35 allow the extension of the - chain (minus chain)
in the 5' → 3' direction, said DNA primer being
selected from the group consisting of DNA primers

comprising a nucleotide sequence represented by SEQ ID NO: 2, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 4, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 8, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 9, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 11, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 15, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 17 and DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 19, or

(ii) carrying out a polymerase chain reaction in the presence of a mixture of

- ① a DNA coding for G protein coupled receptor protein, said DNA being capable of acting as a template,
- ② at least one DNA primer which is capable of binding with the 3'-side nucleotide sequence of the - chain (minus chain) of the template DNA coding for G protein coupled receptor protein to allow the extension of the + chain (plus chain) in the 5' → 3' direction, said DNA primer being selected from the group consisting of DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 1 and DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 12, and
- ③ at least one DNA primer which is capable of binding with the 3'-side nucleotide sequence of the + chain (plus chain) of the template DNA coding for G protein coupled receptor protein to allow the extension of the - chain (minus chain) in the 5' → 3' direction, said DNA primer being selected from the group consisting of DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 13;

(21) a method for screening DNA libraries for

a DNA coding for G protein coupled receptor protein (e.g. from the first to sixth membrane-spanning domains or other domains of G protein coupled receptor protein), which comprises carrying out a polymerase chain reaction in the presence of a mixture of

- ① said DNA library,
- ② at least one DNA primer selected from the group consisting of DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 1 and DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 12, and
- ③ at least one DNA primer selected from the group consisting of DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 2, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 4, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 8, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 9, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 15, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 17 and DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 19,

to amplify selectively a template DNA coding for G protein coupled receptor protein (e.g. from the first to sixth membrane-spanning domains or other domains of G protein coupled receptor protein), contained in the DNA library;

(22) a method for screening DNA libraries for

a DNA coding for G protein coupled receptor protein (e.g. from the first to seventh membrane-spanning domains or other domains of G protein coupled receptor protein), which comprises carrying out a polymerase chain reaction in the presence of a mixture of

- ① said DNA library,
- ② at least one DNA primer selected from the group consisting of DNA primers comprising a nucleotide

sequence represented by SEQ ID NO: 1 and DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 12, and

- 5 ③ at least one DNA primer selected from the group consisting of DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 11,

to amplify selectively a template DNA coding for G protein coupled receptor protein (e.g. from the first to seventh membrane-spanning domains or other domains of G protein coupled receptor protein), contained in the DNA library;

10

(23) a method for screening DNA libraries for a DNA coding for G protein coupled receptor protein (e.g. from the third to sixth membrane-spanning domains or other domains of G protein coupled receptor protein), which comprises carrying out a polymerase chain reaction in the presence of a mixture of

15

- ① said DNA library,
- ② at least one DNA primer selected from the group consisting of DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 3, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 5, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 6, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 7, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 14 and DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 18, and
- 20
- ③ at least one DNA primer selected from the group consisting of DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 2, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 4, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 8, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 9, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 15, DNA
- 25
- 30
- 35

primers comprising a nucleotide sequence
represented by SEQ ID NO: 17 and DNA primers
comprising a nucleotide sequence represented by
SEQ ID NO: 19,

5 to amplify selectively a template DNA coding for G protein
coupled receptor protein (e.g. from the third to sixth
membrane-spanning domains or other domains of G protein coupled
receptor protein), contained in the DNA library;

(24) a method for screening DNA libraries for
10 a DNA coding for G protein coupled receptor protein (e.g. from
the third to seventh membrane-spanning domains or other domains
of G protein coupled receptor protein), which comprises
carrying out a polymerase chain reaction in the presence of a
mixture of

15 ① said DNA library,
② at least one DNA primer selected from the group
consisting of DNA primers comprising a nucleotide
sequence represented by SEQ ID NO: 3, DNA primers
comprising a nucleotide sequence represented by
20 SEQ ID NO: 5, DNA primers comprising a nucleotide
sequence represented by SEQ ID NO: 6, DNA primers
comprising a nucleotide sequence represented by
SEQ ID NO: 7, DNA primers comprising a nucleotide
sequence represented by SEQ ID NO: 14 and DNA
25 primers comprising a nucleotide sequence
represented by SEQ ID NO: 18, and

③ at least one DNA primer selected from the group
consisting of DNA primers comprising a nucleotide
sequence represented by SEQ ID NO: 11,
30 to amplify selectively a template DNA coding for G protein
coupled receptor protein (e.g. from the third to seventh
membrane-spanning domains or other domains of G protein coupled
receptor protein), contained in the DNA library;

(25) a method for screening DNA libraries for
35 a DNA coding for G protein coupled receptor protein (e.g. from
the second to sixth membrane-spanning domains or other domains
of G protein coupled receptor protein), which comprises

carrying out a polymerase chain reaction in the presence of a mixture of

- ① said DNA library,
- ② at least one DNA primer selected from the group consisting of DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 10 and DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 16, and
- ③ at least one DNA primer selected from the group consisting of DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 2, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 4, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 8, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 9, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 15, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 17 and DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 19,

to amplify selectively a template DNA coding for G protein coupled receptor protein (e.g. from the second to sixth membrane-spanning domains or other domains of G protein coupled receptor protein), contained in the DNA library;

(26) a method for screening DNA libraries for a DNA coding for G protein coupled receptor protein (e.g. from the second to seventh membrane-spanning domains or other domains of G protein coupled receptor protein), which comprises carrying out a polymerase chain reaction in the presence of a mixture of

- ① said DNA library,
- ② at least one DNA primer selected from the group consisting of DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 10 and DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 16, and

- ③ at least one DNA primer selected from the group consisting of DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 11,

to amplify selectively a template DNA coding for G protein coupled receptor protein (e.g. from the second to seventh membrane-spanning domains or other domains of G protein coupled receptor protein), contained in the DNA library;

(27) a method for screening DNA libraries for a DNA coding for G protein coupled receptor protein (e.g. from the first to third membrane-spanning domains or other domains of G protein coupled receptor protein), which comprises carrying out a polymerase chain reaction in the presence of a mixture of

- ① said DNA library,

- ② at least one DNA primer selected from the group consisting of DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 1 and DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 12, and

- ③ at least one DNA primer selected from the group consisting of DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 13,

to amplify selectively a template DNA coding for G protein coupled receptor protein (e.g. from the first to third membrane-spanning domains or other domains of G protein coupled receptor protein), contained in the DNA library;

(28) a method for screening DNA libraries for a DNA coding for G protein coupled receptor protein, which comprises carrying out a polymerase chain reaction in the presence of a mixture of

- ① said DNA library,

- ② at least one DNA primer selected from the group consisting of DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 1, and

- ③ at least one DNA primer selected from the group consisting of DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 2,

to amplify selectively the template DNA coding for G protein coupled receptor protein, contained in the DNA library;

(29) a method for screening DNA libraries to detect a DNA coding for G protein coupled receptor protein, which
5 comprises carrying out a polymerase chain reaction in the presence of a mixture of

- ① said DNA library,
- ② at least one DNA primer selected from the group consisting of DNA primers comprising a nucleotide
10 sequence represented by SEQ ID NO: 3, and
- ③ at least one DNA primer selected from the group consisting of DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 4,

to amplify selectively a template DNA coding for G protein coupled receptor protein, contained in the DNA library;
15

(30) a method for screening DNA libraries for a DNA coding for G protein coupled receptor protein, which comprises carrying out a polymerase chain reaction in the presence of a mixture of

- 20 ① said DNA library,
- ② at least one DNA primer selected from the group consisting of DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 6, and
- ③ at least one DNA primer selected from the group
25 consisting of DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 8,

to amplify selectively a template DNA coding for G protein coupled receptor protein, contained in the DNA library;

(31) a method for screening DNA libraries for a DNA coding for G protein coupled receptor protein, which
30 comprises carrying out a polymerase chain reaction in the presence of a mixture of

- ① said DNA library,
- ② at least one DNA primer selected from the group consisting of DNA primers comprising a nucleotide
35 sequence represented by SEQ ID NO: 10, and
- ③ at least one DNA primer selected from the group

consisting of DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 11,

to amplify selectively a template DNA coding for G protein coupled receptor protein, contained in the DNA library; and

5 (32) a method for screening DNA libraries according to any of the above (6), and (21) to (31) wherein said DNA library is derived from an origin selected from the group consisting of human tissues and human cells. Examples of such human tissues include adrenal, umbilical cord, brain, tongue,
10 liver, lymph gland, lung, thymus, placenta, peritoneum, retina, spleen, heart, smooth muscle, intestine, vessel, bone, kidney, skin, fetus, mammary gland, ovary, testis, pituitary gland, pancreas, submandibular gland, spine, prostate gland, stomach, thyroid gland, trachea (windpipe), skeletal muscle, uterus,
15 adipose tissue, urinary bladder, cornea, olfactory bulb, bone marrow, amnion, etc. Examples of such human cells include nerve cells, epithelial cells, endothelial cells, leukocytes, lymphocytes, gliocytes, fibroblasts, keratinized cells, osteoblasts, osteoclasts, astrocytes, melanocytes, various
20 carcinomas, various sarcomas, various cells derived from the above-mentioned human tissues.

Yet another aspect of the present invention is a degenerate deoxynucleotide which has an oligonucleotide sequence to which a SEQ ID NO selected from the group
25 consisting of SEQ ID NO: 1 to SEQ ID NO: 19 is assigned.

Another aspect of the present invention is

(33) a G protein coupled receptor protein comprising an amino acid sequence selected from the group consisting of amino acid sequences represented by SEQ ID NO: 24 and/or SEQ ID
30 NO: 25 and substantial equivalents to the amino acid sequence represented by SEQ ID NO: 24 or SEQ ID NO: 25; or a salt thereof;

(34) a G protein coupled receptor protein according to the above (33) comprising an amino acid sequence selected
35 from the group consisting of an amino acid sequence represented by SEQ ID NO: 26 and substantial equivalents to the amino

acid sequence represented by SEQ ID NO: 26; or a salt thereof;

(35) a G protein coupled receptor protein comprising an amino acid sequence selected from the group consisting of an amino acid sequence represented by SEQ ID NO: 27 and
5 substantial equivalents to the amino acid sequence represented by SEQ ID NO: 27; or a salt thereof;

(36) a G protein coupled receptor protein comprising an amino acid sequence selected from the group consisting of an amino acid sequence represented by SEQ ID NO: 28 and
10 substantial equivalents to the amino acid sequence represented by SEQ ID NO: 28; or a salt thereof;

(37) a G protein coupled receptor protein comprising an amino acid sequence selected from the group consisting of amino acid sequences represented by SEQ ID NO: 34 and/or SEQ ID NO: 35 and substantial equivalents to the amino acid
15 sequence represented by SEQ ID NO: 34 or SEQ ID NO: 35; or a salt thereof;

(38) a G protein coupled receptor protein comprising an amino acid sequence selected from the group consisting of an amino acid sequence represented by SEQ ID NO: 38 and
20 substantial equivalents to the amino acid sequence represented by SEQ ID NO: 38; or a salt thereof;

(39) a G protein coupled receptor protein according to the above (38) comprising an amino acid sequence selected
25 from the group consisting of an amino acid sequence represented by SEQ ID NO: 39 and substantial equivalents to the amino acid sequence represented by SEQ ID NO: 39; or a salt thereof;

(40) a G protein coupled receptor protein comprising an amino acid sequence represented by SEQ ID NO: 56 and
30 substantial equivalents to the amino acid sequence represented by SEQ ID NO: 56; or a salt thereof;

(41) a peptide segment or fragment of a G protein coupled receptor protein according to any of the above (33) to (40), a modified derivative thereof or a salt thereof;

35 (42) a DNA which comprises a nucleotide sequence coding for a G protein coupled receptor protein of the above (33);

(43) a DNA which comprises a nucleotide sequence coding for a G protein coupled receptor protein of the above (34);

5 (44) a DNA which comprises a nucleotide sequence coding for a G protein coupled receptor protein of the above (35);

(45) a DNA which comprises a nucleotide sequence coding for a G protein coupled receptor protein of the above (36);

10 (46) a DNA which comprises a nucleotide sequence coding for a G protein coupled receptor protein of the above (37);

(47) a DNA which comprises a nucleotide sequence coding for a G protein coupled receptor protein of the above
15 (38);

(48) a DNA which comprises a nucleotide sequence coding for a G protein coupled receptor protein of the above (39);

20 (49) a DNA which comprises a nucleotide sequence coding for a G protein coupled receptor protein of the above (40);

(50) a DNA of the above (42) comprising a nucleotide sequence represented by SEQ ID NO: 29 and/or SEQ ID NO: 30;

25 (51) a DNA of the above (43) comprising a nucleotide sequence represented by SEQ ID NO: 31;

(52) a DNA of the above (44) comprising a nucleotide sequence represented by SEQ ID NO: 32;

(53) a DNA of the above (45) comprising a nucleotide sequence represented by SEQ ID NO: 33;

30 (54) a DNA of the above (46) comprising a nucleotide sequence represented by SEQ ID NO: 36 and/or SEQ ID NO: 37;

(55) a DNA of the above (47) comprising a nucleotide sequence represented by SEQ ID NO: 40;

35 (56) a DNA of the above (48) comprising a nucleotide sequence represented by SEQ ID NO: 41;

(57) a DNA of the above (49) comprising a nucleotide sequence represented by SEQ ID NO: 57;

(58) a vector comprising a DNA according to any of the above (42) to (57);

(59) a transformant (including a transfectant) carrying a vector of the above (58);

5 (60) a process for producing a G protein coupled receptor protein or a salt thereof according to any of the above (33) to (40), which comprises culturing a transformant of the above (59) to express said G protein coupled receptor protein on the membrane of the transformant;

10 (61) a method for determining a ligand to a G protein coupled receptor protein according to any of the above (33) to (40), which comprises contacting

(i) at least one component selected from the group
15 consisting of G protein coupled receptor proteins or salts thereof according to any of the above (33) to (40), peptide fragments or segments or salts thereof according to the above (41), and mixtures thereof, with

20 (ii) at least one compound to be tested;

(62) a screening method for a compound capable of inhibiting the binding of a G protein coupled receptor protein according to any of the above (33) to (40) with a ligand, which comprises carrying out a comparison between:

25 (i) at least one case where said ligand is contacted with at least one component selected from the group consisting of G protein coupled receptor proteins or salts thereof according to any of the above (33) to (40), peptide fragments or segments or salts thereof
30 according to the above (41), and mixtures thereof,

and

(ii) at least one case where said ligand together with a compound to be tested is contacted with at least one component selected from the group consisting
35 of G protein coupled receptor proteins or salts thereof according to any of the above (33) to (40), peptide fragments or segments or salts thereof according to the

above (41), and mixtures thereof;

(63) a kit for the screening of one or more compounds capable of inhibiting the binding of a G protein coupled receptor protein according to any of the above (33) to (40), with a ligand, which comprises at least one component selected from the group consisting of G protein coupled receptor proteins or salts thereof according to any of the above (33) to (40), peptide fragments or segments or salts thereof according to the above (41), and mixtures thereof; and

(64) an antibody against at least one component selected from the group consisting of G protein coupled receptor proteins or salts thereof according to any of the above (33) to (40), peptide fragments or segments or salts thereof according to the above (41), and mixtures thereof.

Yet another aspect of the present invention is

(65) a G protein coupled receptor protein according to the above (33) comprising

(i) an amino acid sequence selected from the group consisting of an amino acid sequence represented by SEQ ID NO: 24, amino acid sequences wherein one or more amino acid residues (preferably from 2 to 30 amino acid residues, more preferably from 2 to 10 amino acid residues) are deleted from the amino acid sequence of SEQ ID NO: 24, amino acid sequences wherein one or more amino acid residues (preferably from 2 to 30 amino acid residues, more preferably from 2 to 10 amino acid residues) are added to the amino acid sequence of SEQ ID NO: 24, and amino acid sequences wherein one or more amino acid residues (preferably from 2 to 30 amino acid residues, more preferably from 2 to 10 amino acid residues) in the amino acid sequence of SEQ ID NO: 24 are substituted with one or more other amino acid residues, or/and

(ii) an amino acid sequence selected from the group consisting of an amino acid sequence represented by SEQ ID NO: 25, amino acid sequences wherein one or more amino acid residues (preferably from 2 to 30 amino acid residues, more preferably from 2 to 10 amino acid residues) are deleted from the amino acid sequence of SEQ ID NO: 25, amino acid

sequences wherein one or more amino acid residues (preferably from 2 to 30 amino acid residues, more preferably from 2 to 10 amino acid residues) are added to the amino acid sequence of SEQ ID NO: 25, and amino acid sequences wherein one or
5 more amino acid residues (preferably from 2 to 30 amino acid residues, more preferably from 2 to 10 amino acid residues) in the amino acid sequence of SEQ ID NO: 25 are substituted with one or more other amino acid residues,
or a salt thereof;

10 (66) a G protein coupled receptor protein according to the above (34) comprising an amino acid sequence selected from the group consisting of an amino acid sequence represented by SEQ ID NO: 26, amino acid sequences wherein one or more
15 amino acid residues (preferably from 2 to 30 amino acid residues, more preferably from 2 to 10 amino acid residues) are deleted from the amino acid sequence of SEQ ID NO: 26, amino acid sequences wherein one or more amino acid residues
(preferably from 2 to 30 amino acid residues, more preferably from 2 to 10 amino acid residues) are added to the amino acid
20 sequence of SEQ ID NO: 26, and amino acid sequences wherein one or more amino acid residues (preferably from 2 to 30 amino acid residues, more preferably from 2 to 10 amino acid residues) in the amino acid sequence of SEQ ID NO: 26 are substituted with one or more other amino acid residues, or a salt thereof;

25 (67) a G protein coupled receptor protein according to the above (35) comprising an amino acid sequence selected from the group consisting of an amino acid sequence represented by SEQ ID NO: 27, amino acid sequences wherein one or more
30 amino acid residues (preferably from 2 to 30 amino acid residues, more preferably from 2 to 10 amino acid residues) are deleted from the amino acid sequence of SEQ ID NO: 27, amino acid sequences wherein one or more amino acid residues
(preferably from 2 to 30 amino acid residues, more preferably from 2 to 10 amino acid residues) are added to the amino acid
35 sequence of SEQ ID NO: 27, and amino acid sequences wherein one or more amino acid residues (preferably from 2 to 30 amino acid residues, more preferably from 2 to 10 amino acid residues) in

the amino acid sequence of SEQ ID NO: 27 are substituted with one or more other amino acid residues, or a salt thereof;

(68) a G protein coupled receptor protein according to the above (36) comprising an amino acid sequence selected from the group consisting of an amino acid sequence represented by SEQ ID NO: 28, amino acid sequences wherein one or more amino acid residues (preferably from 2 to 30 amino acid residues, more preferably from 2 to 10 amino acid residues) are deleted from the amino acid sequence of SEQ ID NO: 28, amino acid sequences wherein one or more amino acid residues (preferably from 2 to 30 amino acid residues, more preferably from 2 to 10 amino acid residues) are added to the amino acid sequence of SEQ ID NO: 28, and amino acid sequences wherein one or more amino acid residues (preferably from 2 to 30 amino acid residues, more preferably from 2 to 10 amino acid residues) in the amino acid sequence of SEQ ID NO: 28 are substituted with one or more other amino acid residues, or a salt thereof;

(69) a G protein coupled receptor protein according to the above (37) comprising

(i) an amino acid sequence selected from the group consisting of an amino acid sequence represented by SEQ ID NO: 34, amino acid sequences wherein one or more amino acid residues (preferably from 2 to 30 amino acid residues, more preferably from 2 to 10 amino acid residues) are deleted from the amino acid sequence of SEQ ID NO: 34, amino acid sequences wherein one or more amino acid residues (preferably from 2 to 30 amino acid residues, more preferably from 2 to 10 amino acid residues) are added to the amino acid sequence of SEQ ID NO: 34, and amino acid sequences wherein one or more amino acid residues (preferably from 2 to 30 amino acid residues, more preferably from 2 to 10 amino acid residues) in the amino acid sequence of SEQ ID NO: 34 are substituted with one or more other amino acid residues, or/and

(ii) an amino acid sequence selected from the group consisting of an amino acid sequence represented by SEQ ID NO: 35, amino acid sequences wherein one or more amino acid residues (preferably from 2 to 30 amino acid residues, more

preferably from 2 to 10 amino acid residues) are deleted from the amino acid sequence of SEQ ID NO: 35, amino acid sequences wherein one or more amino acid residues (preferably from 2 to 30 amino acid residues, more preferably from 2 to 10 amino acid residues) are added to the amino acid sequence of SEQ ID NO: 35, and amino acid sequences wherein one or more amino acid residues (preferably from 2 to 30 amino acid residues, more preferably from 2 to 10 amino acid residues) in the amino acid sequence of SEQ ID NO: 35 are substituted with one or more other amino acid residues, or a salt thereof;

(70) a G protein coupled receptor protein according to the above (38) comprising an amino acid sequence selected from the group consisting of an amino acid sequence represented by SEQ ID NO: 38, amino acid sequences wherein one or more amino acid residues (preferably from 2 to 30 amino acid residues, more preferably from 2 to 10 amino acid residues) are deleted from the amino acid sequence of SEQ ID NO: 38, amino acid sequences wherein one or more amino acid residues (preferably from 2 to 30 amino acid residues, more preferably from 2 to 10 amino acid residues) are added to the amino acid sequence of SEQ ID NO: 38, and amino acid sequences wherein one or more amino acid residues (preferably from 2 to 30 amino acid residues, more preferably from 2 to 10 amino acid residues) in the amino acid sequence of SEQ ID NO: 38 are substituted with one or more other amino acid residues, or a salt thereof;

(71) a G protein coupled receptor protein according to the above (39) comprising an amino acid sequence selected from the group consisting of an amino acid sequence represented by SEQ ID NO: 39, amino acid sequences wherein one or more amino acid residues (preferably from 2 to 30 amino acid residues, more preferably from 2 to 10 amino acid residues) are deleted from the amino acid sequence of SEQ ID NO: 39, amino acid sequences wherein one or more amino acid residues (preferably from 2 to 30 amino acid residues, more preferably from 2 to 10 amino acid residues) are added to the amino acid sequence of SEQ ID NO: 39, and amino acid sequences wherein one

or more amino acid residues (preferably from 2 to 30 amino acid residues, more preferably from 2 to 10 amino acid residues) in the amino acid sequence of SEQ ID NO: 39 are substituted with one or more other amino acid residues, or a salt thereof;

5 (72) a G protein coupled receptor protein according to the above (40) comprising an amino acid sequence selected from the group consisting of an amino acid sequence represented by SEQ ID NO: 56, amino acid sequences wherein one or more amino acid residues (preferably from 2 to 30 amino acid
10 residues, more preferably from 2 to 10 amino acid residues) are deleted from the amino acid sequence of SEQ ID NO: 56, amino acid sequences wherein one or more amino acid residues (preferably from 2 to 30 amino acid residues, more preferably from 2 to 10 amino acid residues) are added to the amino acid
15 sequence of SEQ ID NO: 56, and amino acid sequences wherein one or more amino acid residues (preferably from 2 to 30 amino acid residues, more preferably from 2 to 10 amino acid residues) in the amino acid sequence of SEQ ID NO: 56 are substituted with one or more other amino acid residues, or a salt thereof;

20 (73) a method for determining a ligand according to the above (61) wherein said ligand is selected from the group consisting of angiotensin, bombesin, canavaninoid, cholecystokinin, glutamine, serotonin, melatonin, neuropeptide Y, opioid, purine, vasopressin, oxytocin, VIP (vasoactive
25 intestinal and related peptides), somatostatin, dopamine, motilin, amylin, bradykinin, CGRP (calcitonin gene related peptides), adrenomedullin, leukotriene, pancreastatin, prostaglandin, thromboxanes, adenosine, adrenaline, α - and β -chemokine (IL-8, GRO α , GRO β , GRO γ , NAP-2, ENA-78, PF4, IP10, GCP-2, MCP-1, HC14, MCP-3, I-309, MIP1 α ,
30 MIP-1 β , RANTES, etc.), endothelin, enterogastrin, histamine, neurotensin, TRH, pancreatic polypeptides and galanin;

 (74) a method for the screening of a compound or a salt thereof capable of inhibiting the binding of a ligand with
35 a G protein coupled receptor protein according to any of the above (33) to (40), which comprises measuring amounts of a labeled ligand bound to the said G protein coupled receptor

protein in at least two cases:

(i) where the labeled ligand is contacted with at least one component selected from the group consisting of G protein coupled receptor proteins or salts thereof according to any of the above (33) to (40), peptide fragments or segments or salts thereof according to the above (41), and mixtures thereof, and

(ii) where the labeled ligand together with a compound to be tested is contacted with at least one component elected from the group consisting of G protein coupled receptor proteins or salts thereof according to any of the above (33) to (40), peptide fragments or segments or salts thereof according to the above (41), and mixtures thereof,

and comparing the measured amounts of the labeled ligand;

(75) a method for the screening of a compound or a salt thereof capable of inhibiting the binding of a ligand with a G protein coupled receptor protein according to any of the above (33) to (40), which comprises measuring amounts of a labeled ligand bound to a cell comprising the said G protein coupled receptor protein in at least two cases:

(i) where the labeled ligand is contacted with the said cell, and

(ii) where the labeled ligand together with a compound to be tested is contacted with the said cell,

and comparing the measured amounts of the labeled ligand;

(76) a method for the screening of a compound or a salt thereof capable of inhibiting the binding of a ligand with a G protein coupled receptor protein according to any of the above (33) to (40), which comprises measuring amounts of a labeled ligand bound to a membrane fraction of a cell comprising the said G protein coupled receptor protein in at least two cases:

(i) where the labeled ligand is contacted with the said membrane fraction, and

(ii) where the labeled ligand together with a compound to be tested is contacted with the membrane fraction,

- 51 -

and comparing the measured amounts of the labeled ligand;
(77) a method for the screening of a compound or a
salt thereof capable of inhibiting the binding of a ligand with
a G protein coupled receptor protein according to any of the
above (33) to (40), which comprises measuring amounts of a
labeled ligand bound to said G protein coupled receptor protein
in at least two cases:

- 5 (i) where the labeled ligand is contacted with a G protein
coupled receptor protein according to any of the above
(33) to (40) which is expressed on the membrane of a
transformant according to the above (59) during
incubation of the transformant, and
10 (ii) where the labeled ligand together with a compound to
be tested is contacted with the G protein coupled
receptor protein according to any of the above
(33) to (40) which is expressed on the membrane of a
transformant according to the above (59) during
incubation of the transformant,

15 and comparing the measured amounts of the labeled ligand;
(78) a method for the screening of a compound or a
salt thereof capable of inhibiting the binding of a ligand with
a G protein coupled receptor protein according to any of the
above (33) to (40), which comprises measuring G protein coupled
receptor protein-mediated cell-stimulating activities in at
least two cases:

- 20 (i) where a compound capable of activating the G protein
coupled receptor protein according to any of the above
(33) to (40) is contacted with a cell comprising the
said G protein coupled receptor protein, and
25 (ii) where the compound capable of activating the G protein
together with a compound to be tested is contacted with
the cell comprising the said G protein coupled
receptor protein,
30 and comparing the measured cell-stimulating activities;
(79) a method for the screening of a compound or a
salt thereof capable of inhibiting the binding of a ligand with
a G protein coupled receptor protein according to any of the
35

above (33) to (40), which comprises measuring G protein coupled receptor protein-mediated cell-stimulating activities in at least two cases:

5 (i) where a compound capable of activating the G protein coupled receptor protein according to any of the above (33) to (40) is contacted with a G protein coupled receptor protein according to any of the above (33) to (40) which is expressed on the membrane of a transformant according to the above (59) during
10 incubation of the transformant, and

(ii) where the compound capable of activating the G protein together with a compound to be tested is contacted with the G protein coupled receptor protein according to any of the above (33) to (40) which is
15 expressed on the membrane of a transformant according to the above (59) during incubation of the transformant,

and comparing the measured cell-stimulating activities;

(80) a method according to the above (78) or (79)
20 wherein said compound capable of activating the G protein coupled receptor protein according to any of the above (33) to (40) is selected from the group consisting of angiotensin, bombesin, canavanoid, cholecystokinin, glutamine, serotonin, melatonin, neuropeptide Y, opioid, purine, vasopressin,
25 oxytocin, VIP (vasoactive intestinal and related peptides), somatostatin, dopamine, motilin, amylin, bradykinin, CGRP (calcitonin gene related peptides), adrenomedullin, leukotriene, pancreastatin, prostaglandin, thromboxane, adenosine, adrenaline, α - and β -chemokine (IL-8, GRO α , GRO β , GRO γ ,
30 NAP-2, ENA-78, PF4, IP10, GCP-2, MCP-1, HC14, MCP-3, I-309, MIP1 α , MIP-1 β , RANTES, etc.), endothelin, enterogastrin, histamine, neurotensin, TRH, pancreatic polypeptides and galanin;

(81) a compound which is determined through a method
35 according to any of the above (62) and (74) to (80) or a salt thereof;

(82) a pharmaceutical composition comprising an

effective amount of a compound according to the above (81) or a salt thereof;

(83) a screening kit according to the above (63), comprising a cell comprising a G protein coupled receptor protein according to any of the above (33) to (40);

(84) a screening kit according to the above (63), comprising a membrane fraction derived from a cell comprising a G protein coupled receptor protein according to any of the above (33) to (40);

(85) a screening kit according to the above (63), comprising a cell of the (59) or (109) mentioned herein below;

(86) a screening kit according to the above (63), comprising a membrane fraction derived from a cell of the (59) or (109);

(87) a compound which is determined by means of a screening kit according to any of the above (63) and (83) to (86) or a salt thereof;

(88) a pharmaceutical composition comprising an effective amount of a compound according to the above (87) or a salt thereof; and

(89) a method for measuring at least one component selected from the group consisting of G protein coupled receptor proteins or salts thereof according to any of the above (33) to (40), peptide fragments or segments or salts thereof according to the above (41), and mixtures thereof, which comprises contacting an antibody according to the above (64) with the component selected from the group consisting of G protein coupled receptor proteins or salts thereof according to any of the above (33) to (40), peptide segments or salts thereof according to the above (41), and mixtures thereof.

Still another aspect of the present invention is

(90) a ligand to a G protein coupled receptor protein according to any of the above (33) to (40), which is determined through the following step of:

contacting (i) at least one component selected from the group consisting of G protein coupled receptor proteins or salts thereof according to any of the above

(33) to (40), peptide fragments or segments or salts thereof according to the above (41), and mixtures thereof,

with (ii) at least one compound to be examined; and

5 (91) a compound capable of inhibiting the binding of a G protein coupled receptor protein according to any of the above (33) to (40) with a ligand, which is determined through carrying out a comparison between:

- 10 (i) at least one case where said ligand is contacted with at least one component selected from the group consisting of G protein coupled receptor proteins or salts thereof according to any of the above (33) to (40), peptide fragments or segments or salts thereof according to the above (41), and mixtures thereof, and
- 15 (ii) at least one case where said ligand together with a compound to be tested is contacted with at least one component selected from the group consisting of G protein coupled receptor proteins or salts thereof according to any of the above (33) to (40), peptide
- 20 fragments or segments or salts thereof according to the above (41), and mixtures thereof.

Another aspect of the present invention is

(92) a recombinant G protein coupled receptor protein and a salt thereof which is obtained by the expression

25 of a DNA according to any of the above (42) to (57), or a modified or fragmented derivative thereof;

(93) a method for amplifying a DNA coding for G protein coupled receptor protein by polymerase chain reaction techniques, which comprises carrying out a polymerase

30 chain reaction in the presence of a mixture of

- (1) a DNA coding for G protein coupled receptor protein, said DNA being capable of acting as a template, and
- (2) at least one DNA primer selected from the group
- 35 consisting of DNA primers comprising either SEQ ID NO: 1 or SEQ ID NO: 2; and

(94) a method for screening DNA libraries for

a DNA coding for G protein coupled receptor protein, which comprises carrying out a polymerase chain reaction in the presence of a mixture of

(1) said DNA library, and

5 (2) at least one DNA primer selected from the group consisting of DNA primers comprising either SEQ ID NO: 1 or SEQ ID NO: 2,

to amplify selectively the DNA coding for G protein coupled receptor protein, contained in the DNA library.

10 Yet another aspect of the present invention is

(95) a monoclonal antibody against at least one component selected from the group consisting of G protein coupled receptor proteins or salts thereof according to any of the above (33) to (40), peptide fragments or segments or salts thereof according to the above (41), and mixtures thereof;

15 (96) a preparation of purified polyclonal antibodies against at least one component selected from the group consisting of G protein coupled receptor proteins or salts thereof according to any of the above (33) to (40), peptide fragments or segments or salts thereof according to the above (41), and mixtures thereof;

(97) an immunoassay for detecting a G protein coupled receptor protein which comprising

25 (i) incubating a sample to be tested with an antibody according to the above (64) to allow formation of an antigen-antibody complex; and

(ii) detecting an antigen-antibody complex formed in step (i); and

30 (98) an immunoassay for detecting antibodies against a G protein coupled receptor protein which comprising

(i) incubating a sample to be tested with at least one component selected from the group consisting of G protein coupled receptor proteins or salts thereof according to any of the above (33) to (40), peptide fragments or segments or salts thereof according to the above (41), and mixtures thereof to
35 allow formation of an antigen-antibody complex; and

(ii) detecting an antigen-antibody complex formed in step (a).

Still another aspect of the present invention is

(99) an antisense DNA or RNA which comprises a
5 nucleotide sequence complementary to at least a portion of a DNA according to any of the above (42) to (57), said antisense DNA or RNA being hybridizable to said DNA according to any of the above (42) to (57);

(100) an antisense DNA or RNA according to the above
10 (99) wherein said antisense DNA or RNA comprises the 5' end hairpin loop, 5' end 6-base-pair repeat, 5' end untranslated region, protein translation initiation site or codon, ORF translation initiation site or codon, 3'-untranslated region, 3' end palindrome region, or 3' end hairpin loop of a G protein
15 coupled receptor protein DNA according to any of the above (42) to (57);

(101) an antisense DNA or RNA according to the above (99) in a pharmaceutically acceptable carrier;

(102) an antisense DNA or RNA according to the above
20 (99) comprising from 2 to 50 nucleotides;

(103) a method for modulating the activity of a G protein coupled receptor protein comprising contacting cells expressing the G protein coupled receptor protein with an antisense DNA or RNA according to the above (99);

(104) a method for producing an antibody against a G
25 protein coupled receptor protein according to any of the above (33) to (40), which comprises administering to an individual at least one component selected from the group consisting of G protein coupled receptor proteins or salts thereof according to
30 any of the above (33) to (40), peptide fragments or segments or salts thereof according to the above (41), and mixtures thereof; and

(105) a method for producing a hybridoma which
35 produces a monoclonal antibody against a G protein coupled receptor protein according to any of the above (33) to (40), which comprises

(i) immunizing an individual with at least one

component selected from the group consisting of G protein coupled receptor proteins or salts thereof according to any of the above (33) to (40), peptide fragments or segments or salts thereof according to the above (41), and mixtures thereof;

5 (ii) immortalizing antibody producing cells from the immunized individual;

(iii) selecting an immortal cell which produces antibodies reactive with the G protein coupled receptor protein; and

10 (iv) growing said immortal cell.

Yet another aspect of the present invention is

(106) a PCR screening kit for a DNA (or nucleotide sequence) coding for G protein coupled receptor protein in a DNA library which comprises

- 15 (i) ① at least one DNA primer selected from the group consisting of DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 1, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 3, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 5, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 6, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 7, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 10, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 14, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 16 and DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 18, and
- 20 ② at least one DNA primer selected from the group consisting of DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 2, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 4, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 8, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 9, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 11, DNA primers comprising a nucleotide
- 25
- 30
- 35

- sequence represented by SEQ ID NO: 15, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 17 and DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 19; or
- 5 (ii)① at least one DNA primer selected from the group consisting of DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 1 and DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 12, and
- 10 ② at least one DNA primer selected from the group consisting of DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 13;
- (107) a vector comprising the DNA according to the above (7);
- 15 (108) an expression system comprising an open reading frame (ORF) of DNA derived from a G protein coupled receptor protein DNA according to any of the above (7) and (42) to (57), wherein the ORF is operably linked to a control sequence compatible with a desired host cell;
- 20 (109) a transformant (including a transfectant) carrying a vector of the above (107) or an expression system of the above (108);
- (110) a process for producing a G protein coupled receptor protein or a salt thereof, which comprises culturing
- 25 the transformant of the above (109) to express said G protein coupled receptor protein on the membrane of the transformant;
- (111) a method for expressing a polypeptide of G protein coupled receptor protein, comprising:
- (a) providing a transformant of the above (59) or
- 30 (109); and
- (b) incubating the transformant under conditions which allow expression of the polypeptide of G protein coupled receptor protein;
- (112) a method for preparing a transformant
- 35 according to the above (59) or (109), comprising:
- (a) providing a host cell capable of transformation;

(b) providing a vector according to the above (58) or (107) or an expression system according to the above (108); and

5 (c) incubating (a) with (b) under conditions which allow transformation of the host cell with the vector or the expression system;

(113) a pharmaceutical composition according to the above (82) or (88), comprising an effective amount of a compound according to the above (81) or (87) or a
10 pharmaceutically acceptable salt thereof in admixture with a pharmaceutically acceptable carrier, excipient or diluent;

(114) the pharmaceutical composition according to the above (82) or (88), for inhibiting the binding of a G protein coupled receptor protein according to the present invention
15 with a ligand;

(115) a method for inhibiting the binding of a G protein coupled receptor protein according to the present invention with a ligand in a medium which comprises contacting an effective amount of a compound according to
20 the above (81) or (87) or a salt thereof with said medium;

(116) a method for modulating the activity of a G protein coupled receptor protein comprising contacting cells expressing the G protein coupled receptor protein with a an effective amount of a compound according to the above (81)
25 or (87) or a salt thereof;

(117) the ligand according to the above (90) being labeled with a detectable reporter;

(118) the antibody according to the above (64) wherein the antibody is labeled with a detectable reporter;

30 (119) a pharmaceutical composition for controlling an expression of G protein coupled receptor protein, which comprises an effective amount of the antisense DNA according to the above (99), and

(120) a culture product produced by a transformant
35 according to the above (59) or (109).

Yet another aspect of the present invention is

(121) a DNA according to the above (1) wherein the

DNA is an oligonucleotide having from 8 to 60 base residues;

(122) a DNA according to the above (1) wherein the DNA is synthetic;

(123) a DNA (or nucleotide sequence) coding for a G protein coupled receptor protein or a fragment thereof, which is obtained through the method according to any of the above (5) to (32);

(124) a DNA (or nucleotide sequence) according to the above (123), wherein said G protein coupled receptor protein is selected from the group consisting of angiotensin receptor, bombesin receptor, canavaninoid receptor, cholecystokinin receptor, glutamine receptor, serotonin receptor, melatonin receptor, neuropeptide Y receptor, opioid receptor, purine receptor, vasopressin receptor, oxytocin receptor, VIP receptor (vasoactive intestinal and related peptide receptor), somatostatin receptor, dopamine receptor, motilin receptor, amylin receptor, bradykinin receptor, CGRP receptor (calcitonin gene related peptide receptor), adrenomedullin receptor, leukotriene receptor, pancreastatin receptor, prostaglandin receptor, thromboxane receptor, adenosine receptor, adrenaline receptor, α - and β -chemokine receptor including IL-8, GRO α , GRO β , GRO γ , NAP-2, ENA-78, PF4, IP10, GCP-2, MCP-1, HC14, MCP-3, I-309, MIP1 α , MIP-1 β , and RANTES receptors, endothelin receptor, enterogastrin receptor, histamine receptor, neurotensin receptor, TRH receptor, pancreatic polypeptide receptor, and galanin receptor; and

(125) a culture product produced by a transformant according to the above (59) or (109).

As used herein the term "substantial equivalent(s)" means that the activity of the protein, e.g., nature of the ligand binding activity, and physical characteristics are substantially the same. Substitutions, deletions or insertions of amino acids often do not produce radical changes in the physical and chemical characteristics of a polypeptide, in which case polypeptides containing the substitution,

deletion, or insertion would be considered to be substantially equivalent to polypeptides lacking the substitution, deletion, or insertion. Substantially equivalent substitutes for an amino acid within the sequence may be selected from other members of the class to which the amino acid belongs.

The non-polar (hydrophobic) amino acids include alanine, leucine, isoleucine, valine, proline, phenylalanine, tryptophan and methionine. The polar neutral amino acids include glycine, serine, threonine, cysteine, tyrosine, asparagine, and glutamine. The positively charged (basic) amino acids include arginine, lysine and histidine. The negatively charged (acidic) amino acids include aspartic acid and glutamic acid.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 depicts the community (homology) of the sequence of 5' side synthetic DNA primers (HS-1) having a nucleotide sequence represented by SEQ ID NO: 1 with the nucleotide sequence each of other G protein coupled receptor protein-encoding cDNAs and genes.

Figure 2 depicts the community (homology) of the sequence which is complementary to 3' side synthetic DNA primers (HS-2) having a nucleotide sequence represented by SEQ ID NO: 2 with the nucleotide sequence each of other G protein coupled receptor protein-encoding cDNAs and genes.

Figure 3 depicts the community (homology) of the sequence of 5' side synthetic DNA primers (3A) having a nucleotide sequence represented by SEQ ID NO: 5 or 5' side synthetic DNA primers (3B) having a nucleotide sequence represented by SEQ ID NO: 6 relative to the nucleotide sequence each of other G protein coupled receptor protein-encoding cDNAs and genes.

Figure 4 depicts the community (homology) of the sequence of 5' side synthetic DNA primers (3C) having a nucleotide sequence represented by SEQ ID NO: 7 or 5' side synthetic DNA primers (3D) having a nucleotide sequence represented by SEQ ID NO: 3 relative to the nucleotide sequence each of other G protein coupled receptor protein-encoding cDNAs

and genes.

Figure 5 depicts the community (homology) of the sequence (6A) which is complementary to 3' side synthetic DNA primers having a nucleotide sequence represented by SEQ ID NO: 8 or the nucleotide sequence (6B) which is complementary to 3' side synthetic DNA primers having a nucleotide sequence represented by SEQ ID NO: 9 relative to the nucleotide sequence each of other G protein coupled receptor protein-encoding cDNAs and genes.

Figure 6 depicts the community (homology) of the sequence (6C) which is complementary to 3' side synthetic DNA primers having a nucleotide sequence represented by SEQ ID NO: 4 relative to the nucleotide sequence each of other G protein coupled receptor protein-encoding cDNAs and genes.

Figure 7 depicts the community (homology) of the sequence of 5' side synthetic DNA primers (T2A) having a nucleotide sequence represented by SEQ ID NO: 10 with the nucleotide sequence each of other G protein coupled receptor protein-encoding cDNAs and genes.

Figure 8 depicts the community (homology) of the sequence which is complementary to 3' side synthetic DNA primers (T7A) having a nucleotide sequence represented by SEQ ID NO: 11 relative to the nucleotide sequence each of other G protein coupled receptor protein-encoding cDNAs and genes.

Figure 9 depicts the community (homology) of the sequence of 5' side synthetic DNA primers (TM1-A2) having a nucleotide sequence represented by SEQ ID NO: 12 relative to the nucleotide sequence each of other G protein coupled receptor protein-encoding cDNAs and genes.

Figure 10 depicts the community (homology) of the sequence which is complementary to 3' side synthetic DNA primers (TM3-B2) having a nucleotide sequence represented by SEQ ID NO: 13 relative to the nucleotide sequence each of other G protein coupled receptor protein-encoding cDNAs and genes.

Figure 11 depicts the community (homology) of the sequence of 5' side synthetic DNA primers (TM3-C2) having a nucleotide sequence represented by SEQ ID NO: 14 relative to

- 63 -

the nucleotide sequence each of other G protein coupled receptor protein-encoding cDNAs and genes.

5 Figure 12 depicts the community (homology) of the sequence which is complementary to 3' side synthetic DNA primers (TM6-E2) having a nucleotide sequence each of other SEQ ID NO: 15 relative to the nucleotide sequence each of other G protein coupled receptor protein-encoding cDNAs and genes.

10 Figure 13 depicts the community (homology) of the sequence of 5' side synthetic DNA primers (TM2F18) having a nucleotide sequence each of other G protein coupled receptor protein-encoding cDNAs and genes.

15 Figure 14 depicts the community (homology) of the sequence which is complementary to 3' side synthetic DNA primers (TM6R21) having a nucleotide sequence each of other G protein coupled receptor protein-encoding cDNAs and genes.

20 Figure 15 depicts the community (homology) of the sequence of 5' side synthetic DNA primers (S3A) having a nucleotide sequence each of other G protein coupled receptor protein-encoding cDNAs and genes.

25 Figure 16 depicts the community (homology) of the sequence which is complementary to 3' side synthetic DNA primers (S6A) having a nucleotide sequence each of other G protein coupled receptor protein-encoding cDNAs and genes.

30 Figure 17 is the 1.2% agarose gel electrophoresis profile of cDNA products each obtained from human brain amygdala (1, 2, 7), human pituitary body (3, 4, 8) and rat brain (5, 6, 9) by PCR amplification using the synthetic DNA primers having a nucleotide sequence represented by SEQ ID NO: 1 and the synthetic DNA primers having a nucleotide sequence represented by SEQ ID NO: 2, wherein lanes 1 to 6 show the results of when PCR is carried out under severe conditions as disclosed in Examples, lanes 7 to 9 show the results of when PCR is carried out under mild conditions, and M denotes a size

35

marker which is obtained by cutting λ -phage DNA with restriction enzyme, EcoT14I.

Figure 18 shows the nucleotide sequence determined by sequencing of clone A58 with a T7 primer wherein the clone A58 is obtained by amplifying human brain amygdala-derived cDNA by PCR under mild conditions and subcloning it to pCRTM II.

Figure 19 shows the nucleotide sequence determined by sequencing of clone A58 with an SP6 primer.

Figure 20 shows the nucleotide sequence determined by sequencing of clone 57-A-2 by using a -21M13 primer wherein the clone 57-A-2 is obtained by amplifying human brain amygdala-derived cDNA by PCR under severe conditions and subcloning it to pCRTM II.

Figure 21 shows the nucleotide sequence determined by sequencing of clone B54 with a T7 primer wherein the clone B54 is obtained by amplifying rat whole brain-derived cDNA by PCR under mild conditions and subcloning it to pCRTM II.

Figure 22 illustrates the nucleotide sequence of the human pituitary gland-derived G protein coupled receptor protein cDNA fragment included in the cDNA clone p19P2 isolated by PCR using a human pituitary gland-derived cDNA and the amino acid sequence encoded thereby, wherein the primer used for sequencing is -21M13, and the underlined part corresponds to the synthetic primer.

Figure 23 illustrates the nucleotide sequence of the human pituitary gland-derived G protein coupled receptor protein cDNA fragment included in the cDNA clone p19P2 isolated by PCR using a human pituitary gland-derived cDNA and the amino acid sequence encoded thereby, wherein the primer used for sequencing is M13RV-N (Takara, Japan), and the underlined part corresponds to the synthetic primer.

Figure 24 is the partial hydrophobicity plotting profile of the protein encoded by the human pituitary gland-derived G protein coupled receptor protein cDNA fragment included in p19P2, prepared based upon the amino acid sequence

shown in Figure 22.

Figure 25 is the partial hydrophobicity plotting profile of the protein encoded by the human pituitary gland-derived G protein coupled receptor protein cDNA fragment included in p19P2, prepared based upon the amino acid sequence shown in Figure 23.

Figure 26 shows the partial amino acid sequence (p19P2) of the protein encoded by the human pituitary gland-derived G protein coupled receptor protein cDNA fragment included in p19P2, as shown in Figures 22 and 23, relative to the known G protein coupled receptor protein, S12863, wherein reverse amino acid residues are in agreement, the 1st to 99th amino acid residues of the p19P2 sequence correspond to the 1st to 99th amino acid residues in Figure 22, and the 156th to 230th amino acid residues thereof correspond to the 1st to 68th amino acid residues in Figure 23.

Figure 27 is the nucleotide sequence of the MIN6-derived G protein coupled receptor protein cDNA fragment derived based upon the nucleotide sequences of the MIN6-derived G protein coupled receptor protein cDNA fragments each included in the cDNA clones, pG3-2 and pG1-10, isolated by PCR using a MIN6-derived cDNA and the amino acid sequence encoded thereby, wherein the underlined parts corresponds to the synthetic primers.

Figure 28 is the partial hydrophobicity plotting profile of the MIN6-derived G protein coupled receptor protein, prepared based upon the partial amino acid sequence shown in Figure 27.

Figure 29 is the partial nucleotide sequence of the novel receptor protein cDNA clone, p63A2, obtained from the human amygdaloid nucleus by PCR amplification and the amino acid sequence encoded thereby, wherein the underlined part corresponds to the synthetic primer.

Figure 30 is the partial nucleotide sequence of the novel receptor protein cDNA clone, p63A2, obtained from the human amygdaloid nucleus by PCR amplification and the amino acid sequence encoded thereby, wherein the underlined part

corresponds to the synthetic primer.

Figure 31 is the hydrophobicity plotting profile, prepared based upon the amino acid sequence shown in Figure 29, suggesting the presence of hydrophobic domains as designated
5 by 1 to 3.

Figure 32 is the hydrophobicity plotting profile, prepared based upon the amino acid sequence shown in Figure 30, suggesting the presence of hydrophobic domains as designated
by 4 to 6.

10 Figure 33 is the partial amino acid sequence (p63A2) of the protein encoded by the novel receptor protein cDNA fragment included in p63A2, relative to the partial amino acid sequence of the G protein coupled receptor protein (P30731) expressed and induced by a mouse T cell-derived glucocorticoid,
15 wherein reverse amino acid residues are in agreement.

Figure 34 is the whole nucleotide sequence of the the human pituitary gland-derived G protein coupled receptor protein cDNA, included in the cDNA clone, phGR3, isolated from the human-derived cDNA library by plaque hybridization using
20 an DNA insert in the p19P2 as a probe, and the amino acid sequence encoded thereby.

Figure 35 is the northern blotting profile of the human pituitary gland mRNA of the receptor gene encoded by the human pituitary gland-derived cDNA clone, phGR3.

25 Figure 36 is the hydrophobicity plotting profile of the protein encoded by the human pituitary gland-derived G protein coupled receptor protein cDNA included in phGR3, prepared based upon the amino acid sequence shown in Figure 34.

Figure 37 is the partial nucleotide sequence of the novel receptor protein cDNA clone, p3H2-17, obtained from mouse pancreatic β -cell strain, MIN6, by PCR amplification and the amino acid sequence encoded thereby, wherein the underlined part corresponds to the synthetic primer used for the PCR amplification.

35 Figure 38 is the hydrophobicity plotting profile, prepared based upon the amino acid sequence shown in Figure 37, suggesting the presence of hydrophobic domains as designated

by 3 to 6.

Figure 39 is the partial amino acid sequence encoded by the novel receptor protein cDNA included in p3H2-17, relative to the partial amino acid sequence each of chicken ATP receptor protein (P34996), human somatostatin receptor subtype 3 protein (A46226), human somatostatin receptor subtype 4 protein (JN0605) and bovine neuropeptide Y receptor protein (S28787), wherein reverse amino acid residues are in agreement.

Figure 40 is the partial nucleotide sequence of the novel receptor protein cDNA clone, p3H2-34, obtained from mouse pancreatic β -cell strain, MIN6, by PCR amplification and the amino acid sequence encoded thereby, wherein the underlined parts correspond to the synthetic primers used for the PCR amplification.

Figure 41 is the hydrophobicity plotting profile, prepared based upon the amino acid sequence shown in Figure 40, wherein the axis of ordinate represents an index of hydrophobicity, the axis of abscissa represents the number of amino acids and numerals 3 to 6 represent the presence of hydrophobic domains.

Figure 42 is the partial amino acid sequence encoded by the novel receptor protein cDNA included in p3H2-34, relative to the partial amino acid sequence each of human somatostatin receptor subtype 4 protein (JN0605), human somatostatin receptor subtype 2 protein (B41795) and rat-derived ligand unknown receptor protein (A39297), wherein reverse amino acid residues are in agreement.

Figure 43 is the nucleotide sequence of the rabbit gastropyloric part smooth muscle-derived G protein coupled receptor protein cDNA fragment included in the novel receptor protein cDNA clone, pMD4, obtained from rabbit gastropyloric part smooth muscles by PCR amplification, and the amino acid sequence encoded thereby, wherein the underlined parts correspond to the synthetic primers used for the PCR amplification.

Figure 44 is the hydrophobicity plotting profile of the protein encoded by the rabbit gastropyloric part smooth

muscle-derived G protein coupled receptor protein cDNA fragment included in pMD4, prepared based upon the amino acid sequence shown in Figure 35, wherein numerals 1 to 3 suggest the presence of hydrophobic domains.

5 Figure 45 is the partial amino acid sequence (pMD4) of the protein encoded by the rabbit gastropyrolic part smooth muscle-derived G protein coupled receptor protein cDNA fragment included in pMD4 as shown in Figure 43, relative to the known G protein coupled receptor protein, rat ligand unknown receptor
10 protein (A35639), wherein reverse amino acid residues are in agreement, the 1st to 88th amino acid residues of the pMD4 sequence correspond to the 1st to 88th amino acid residues in Figure 43.

15 Figure 46 shows the nucleotide sequence of the mouse-derived galanin receptor protein cDNA clone, pMGR20, which has been cloned with, as a probe, the cDNA insert in p3H2-34 and the amino acid sequence encoded thereby.

20 Figure 47 is the hydrophobicity plotting profile, prepared based upon the amino acid sequence shown in Figure 46, wherein the axis of ordinate represents an index of hydrophobic property, the axis of abscissa represents the number of amino acids, and numerals 1 to 7 represent the presence of hydrophobic domains.

25 Figure 48 is the amino acid sequence (MOUSEGALRECE) of the mouse-derived galanin receptor protein encoded by pMGR20, relative to the amino acid sequence (HUMAGALAMI) of the human-derived galanin receptor protein, wherein reverse amino acid residues are in agreement.

30 Figure 49 is the nucleotide sequence of the rabbit gastropyrolic part smooth muscle-derived G protein coupled receptor protein cDNA fragment included in the novel receptor protein cDNA clone, pMJ10, obtained from rabbit gastropyrolic part smooth muscles by PCR amplification and the amino acid sequence encoded thereby, wherein the underlined parts
35 corresponds to the synthetic primers used for the PCR amplification.

Figure 50 is the hydrophobicity plotting profile of

the protein encoded by the rabbit gastropyrolic part smooth muscle-derived G protein coupled receptor protein cDNA fragment included in pMJ10, prepared based upon the amino acid sequence shown in Figure 49, wherein numerals 4 to 6 suggest the presence of hydrophobic domains.

Figure 51 is the partial amino acid sequence (pMJ10) of the protein encoded by the rabbit gastropyrolic part smooth muscle-derived G protein coupled receptor protein cDNA fragment included in pMJ10 shown in Figure 49, relative to human ligand unknown receptor protein (B42009), human N-formylpeptide receptor protein (JC2014), rabbit N-formylpeptide receptor protein (A46520), mouse C5a anaphylatoxin receptor protein (A46525) and bovine neuropeptide Y receptor protein (S28787) which are known G protein coupled receptor proteins, wherein reverse amino acid residues are in agreement, and the 1st to 125th amino acid residues of pMJ10 correspond to the 1st to 125th amino acid residues in Figure 49.

Figure 52 is the nucleotide sequence of the rabbit gastropyrolic part smooth muscle-derived G protein coupled receptor protein cDNA fragment included in the novel receptor protein cDNA clone, PMH28, obtained from rabbit gastropyrolic part smooth muscles by PCR amplification and the amino acid sequence encoded thereby, wherein the underlined parts correspond to the synthetic primers used for the PCR amplification.

Figure 53 is the hydrophobicity plotting profile of the protein encoded by the rabbit gastropyrolic part smooth muscle-derived G protein coupled receptor protein cDNA fragment included in PMH28, prepared based upon the amino acid sequence shown in Figure 52, wherein numerals 4 to 6 suggest the presence of hydrophobic domains.

Figure 54 is the partial amino acid sequence (PMH28) of the protein encoded by the rabbit gastropyrolic part smooth muscle-derived G protein coupled receptor protein cDNA fragment included in PMH28 shown in Figure 52, relative to mouse IL-8 receptor protein (P35343), human somatostatin receptor protein 1 (A41795) and human somatostatin receptor protein 4 (A47457)

which are known G protein coupled receptor proteins, wherein reverse amino acid residues are in agreement, and the 1st to 119th amino acid residues of pMH28 correspond to the 1st to 119th amino acid residues in Figure 52.

5 Figure 55 is the nucleotide sequence of the rabbit gastropyloric part smooth muscle-derived G protein coupled receptor protein cDNA fragment included in the novel receptor protein cDNA clone, pMN7, obtained from rabbit gastropyloric part smooth muscles by PCR amplification and the amino acid
10 sequence encoded thereby, wherein the underlined 5'-end nucleotide sequence part corresponds to the synthetic primer used for the PCR amplification.

 Figure 56 is the nucleotide sequence of the rabbit gastropyloric part smooth muscle-derived G protein coupled
15 receptor protein cDNA fragment included in the novel receptor protein cDNA clone, pMN7, obtained from rabbit gastropyloric part smooth muscles by PCR amplification and the amino acid sequence encoded thereby, wherein the underlined 3'-end
20 nucleotide sequence part corresponds to the synthetic primer used for the PCR amplification.

 Figure 57 is the hydrophobicity plotting profile of the protein encoded by the rabbit gastropyloric part smooth muscle-derived G protein coupled receptor protein cDNA fragment included in pMN7, prepared based upon the amino acid sequences
25 shown in Figures 55 and 56, wherein numerals TM2 to TM6 suggest the presence of hydrophobic domains.

 Figure 58 is the partial hydrophobicity plotting profile of the protein encoded by the human pituitary gland-derived G protein coupled receptor protein cDNA fragment
30 included in p19P2, prepared based upon the amino acid sequence shown in Figure 22.

 Figure 59 is the partial hydrophobicity plotting profile of the protein encoded by the human pituitary gland-derived G protein coupled receptor protein cDNA fragment
35 included in p19P2, prepared based upon the amino acid sequence shown in Figure 23.

 Figure 60 shows the partial amino acid sequence

(p19P2) of the protein encoded by the human pituitary gland-derived G protein coupled receptor protein cDNA fragment included in p19P2, as shown in Figures 22 and 23, relative to the known G protein coupled receptor protein, S12863, wherein reverse amino acid residues are in agreement, the 1st to 99th amino acid residues of the p19P2 sequence correspond to the 1st to 99th amino acid residues in Figure 22, and the 156th to 230th amino acid residues thereof correspond to the 1st to 68th amino acid residues in Figure 23.

Figure 61 is the partial amino acid sequence (pG3-2/pG1-10) of the MIN6-derived G protein coupled receptor protein, as shown in Figure 27, relative to the partial amino acid sequence (p19P2) of the protein encoded by p19P2, as shown in Figures 22 and 23, wherein reverse amino acid residues are in agreement, the 1st to 99th amino acid residues of the p19P2 sequence correspond to the 1st to 99th amino acid residues in Figure 22, the 156th to 223rd amino acid residues thereof correspond to the 1st to 68th amino acid residues in Figure 23, and the 1st to 223rd amino acid residues of the pG3-2/pG1-10 sequence correspond to the 1st to 223rd amino acid residues in Figure 27.

Figure 62 is the nucleotide sequence of the MIN6-derived G protein coupled receptor protein cDNA fragment included in the cDNA clone, p5S38, isolated by PCR using a MIN6-derived cDNA and the amino acid sequence encoded thereby, wherein the underlined parts corresponds to the synthetic primers.

Figure 63 is the partial amino acid sequence (p5S38) of the MIN6-derived G protein coupled receptor protein, as shown in Figure 62, relative to the partial amino acid sequence (p19P2) of the G protein coupled receptor protein encoded by p19P2, as shown in Figures 22 and 23, as well as the partial amino acid sequence of the G protein coupled receptor protein encoded by the nucleotide sequence derived from the nucleotide sequence of the cDNA fragment included in pG3-2 and pG1-10, as shown in Figure 27, wherein reverse amino acid residues are in agreement, the 1st to 144th amino acid residues of the p5S38

sequence correspond to the 1st to 144th amino acid residues in Figure 62, the 1st to 99th amino acid residues of the p19P2 sequence correspond to the 1st to 99th amino acid residues in Figure 22, the 156th to 223rd amino acid residues thereof correspond to the 1st to 68th amino acid residues in Figure 23, and the 1st to 223rd amino acid residues of the pG3-2/pG1-10 sequence correspond to the 1st to 223rd amino acid residues in Figure 27.

Figure 64 is the partial hydrophobicity plotting profile of the protein encoded by the MIN6-derived G protein coupled receptor protein cDNA fragment included in p5S38, prepared based upon the amino acid sequence shown in Figure 62.

Figure 65 shows the northern blot analysis profile of the receptor gene encoded by the cDNA included in the mouse pancreatic β -cell strain MIN6-derived novel receptor protein cDNA clone, p3H2-17, for mouse cell line, MIN6, Neuro-2a cell and mouse brain, thymus, spleen and pancreas poly(A)⁺ RNA, wherein each arrow and number indicates the size marker position (unit of number: kb).

Figure 66 shows the agarose gel electrophoresis analysis profile of the PCR products obtained by 5'RACE PCR of the receptor gene included in p3H2-17 using mouse thymus and spleen poly(A)⁺ RNA.

Lane 1 indicates the size marker 6 (Wako Pure Chemical, Japan).

Lane 2 indicates the internal control which is the thymus-derived PCR product obtained by PCR amplification using the primer having SEQ ID NO: 20 and the primer having SEQ ID NO: 22 with Taq polymerase.

Lane 3 indicates the negative control which is the PCR product obtained by Ex Taq polymerase PCR amplification of thymus cDNA prior to addition of anchors.

Lane 4 indicates the negative control which is the PCR product obtained by Taq polymerase PCR amplification of thymus cDNA prior to addition of anchors.

Lane 5 indicates the PCR product obtained by 5'RACE of thymus poly(A)⁺ RNA with Pfu polymerase.

Lane 6 indicates the PCR product obtained by 5'RACE of thymus poly(A)⁺ RNA with Vent polymerase.

Lane 7 indicates the PCR product obtained by 5'RACE of thymus poly(A)⁺ RNA with Ex Taq polymerase.

5 Lane 8 indicates the PCR product obtained by 5'RACE of thymus poly(A)⁺ RNA with Taq polymerase.

Lane 9 indicates the size marker 5 (Wako Pure Chemical, Japan).

10 Lane 10 indicates the internal control which is the spleen-derived PCR product obtained by PCR amplification using the primer having SEQ ID NO: 20 and the primer having SEQ ID NO: 22 with Taq polymerase.

15 Lane 11 indicates the negative control which is the PCR product obtained by Ex Taq polymerase PCR amplification of spleen cDNA prior to addition of anchors.

Lane 12 indicates the negative control which is the PCR product obtained by Taq polymerase PCR amplification of spleen cDNA prior to addition of anchors.

20 Lane 13 indicates the PCR product obtained by 5'RACE of poly(A) RNA⁺ with Pfu polymerase.

Lane 14 indicates the PCR product obtained by 5'RACE of spleen poly(A)⁺ RNA with Vent polymerase.

Lane 15 indicates the PCR product obtained by 5'RACE of spleen poly(A)⁺ RNA with Ex Taq polymerase.

25 Lane 16 indicates the PCR product obtained by 5'RACE of spleen poly(A)⁺ RNA with Taq polymerase.

Lane 17 indicates the size marker 5 (Wako Pure Chemical, Japan).

Each blacked triangle indicates the band recovered.

30 Figure 67 shows the agarose gel electrophoresis analysis profile of the PCR products obtained by 3'RACE PCR of the receptor gene included in p3H2-17 using mouse thymus and spleen poly(A)⁺ RNA.

35 Lane 1 indicates the size marker 5 (Wako Pure Chemical, Japan).

Lane 2 indicates the PCR product obtained by 3'RACE of spleen poly(A)⁺ RNA with Taq polymerase.

Lane 3 indicates the PCR product obtained by 3'RACE of spleen poly(A)⁺ RNA with Ex Taq polymerase.

Lane 4 indicates the PCR product obtained by 3'RACE of spleen poly(A)⁺ RNA with Vent polymerase.

5 Lane 5 indicates the PCR product obtained by 3'RACE of spleen poly(A)⁺ RNA with Pfu polymerase.

Lane 6 indicates the PCR product obtained by 3'RACE of thymus poly(A)⁺ RNA with Taq polymerase.

10 Lane 7 indicates the PCR product obtained by 3'RACE of thymus poly(A)⁺ RNA with Ex Taq polymerase.

Lane 8 indicates the PCR product obtained by 3'RACE of thymus poly(A)⁺ RNA with Vent polymerase.

Lane 9 indicates the PCR product obtained by 3'RACE of thymus poly(A)⁺ RNA with Pfu polymerase.

15 Lane 10 indicates the size marker 6 (Wako Pure Chemical, Japan).

Each blacked triangle indicates the band recovered.

Figure 68 depicts the model of the RACE products of the receptor protein cDNA fragment included in p3H2-17 obtained
20 by 5'RACE and 3'RACE. Open squares represent regions which have already been isolated and included in p3H2-17. Small arrows, ①, ②, ③ and ④, indicate the positions and directions of the primers designed in Working Example 19. The big arrow shows a predicted full-length open reading frame of the
25 receptor protein held by p3H2-17. Numbers at both ends, N26, N64, N75, C2, C13 and C15, indicate clone numbers of the RACE products obtained. Among these RACE products, N26, N64 and N75 are inserted into pCRTM II vector and C2, C13 and C15 are inserted into the SmaI site of pUC18. The solid triangle
30 indicates the PCR error position which has been clarified through sequencing.

Figure 69 is the nucleotide sequence of the open reading frame and neighboring regions thereof of mouse G protein coupled receptor protein cDNA included in the cDNA
35 clone pMAH2-17 obtained from mouse spleen and thymus poly(A) RNA by RACE techniques based on the nucleotide sequence of the cDNA fragment included in p3H2-17 and the amino acid

sequence encoded thereby.

Figure 70 is the hydrophobicity plotting profile of the protein encoded by the receptor protein cDNA included in pMAH2-17, prepared based upon the amino acid sequence shown in Figure 69.

Figure 71 is the amino acid sequence (75+13CODING) of the protein encoded by the mouse-derived G protein coupled receptor protein cDNA fragment included in pMAH2-17, as shown in Figure 69, relative to the known G protein coupled receptor proteins, mouse P_{2U} purinoceptor (P2UR MOUSE) and chicken P_{2Y} purinoceptor (P2YR CHICK), wherein reverse amino acid residues are in agreement.

Figure 72 is the nucleotide sequence (from 1st to 540th nucleotides) of the rabbit gastropyrolic part smooth muscle-derived G protein coupled receptor protein cDNA fragment included in the novel rabbit gastropyrolic part smooth muscles by PCR amplification, and the amino acid sequence encoded thereby, wherein the underlined 5' part corresponds to the synthetic primer used for the PCR amplification.

Figure 73 is the nucleotide sequence (from 541st to 843rd nucleotides) of the rabbit gastropyrolic part smooth muscle-derived G protein coupled receptor protein cDNA fragment included in the novel rabbit gastropyrolic part smooth muscles by PCR amplification, and the amino acid sequence encoded thereby, wherein the underlined 3' part corresponds to the synthetic primer used for the PCR amplification.

Figure 74 is the hydrophobicity plotting profile of the protein encoded by the rabbit gastropyrolic part smooth muscle-derived G protein coupled receptor protein cDNA fragment included in pMN128, prepared based upon the amino acid sequences shown in Figures 72 and 73, suggesting the presence of hydrophobic domains.

Figure 75 shows inward currents evoked by ATP in Xenopus oocytes injected with cDNA of pMAH2-17-encoded receptor.

Figure 76 is the nucleotide sequence of the human-derived G protein coupled receptor protein cDNA fragment included in ph3H2-17, relative to the nucleotide sequence of the mouse-derived G protein coupled receptor protein cDNA fragment included in p3H2-17, wherein reverse base residues are in agreement.

Figure 77 is the nucleotide sequence of the open reading frame and neighboring regions thereof of human-derived G protein coupled receptor protein cDNA included in phAH2-17 and the amino acid sequence encoded thereby.

Figure 78 is the hydrophobicity plotting profile of the protein encoded by the human-derived G protein coupled receptor protein cDNA included in phAH2-17.

Figure 79 is the amino acid sequence of human type purinoceptor encoded by phAH2-17, relative to the mouse purinoceptor encoded by p3H2-17, wherein reverse amino acid residues are in agreement.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENT

According to the present invention, DNA sequences comprising each a nucleotide sequence indicated by a SEQ ID NO selected from the group consisting of SEQ ID NO: 1 to SEQ ID NO: 19 have been synthesized and characterized. The DNA is a potent primer for polymerase chain reaction in order to amplify DNA sequences encoding part or all of the polypeptide sequence of G protein coupled receptor protein. PCR amplification methods of the DNA coding for part or all of the polypeptide sequence of G protein coupled receptor protein can be advantageously carried out with the said primer DNA. Screening of DNA libraries for the DNA encoding part or all of the polypeptide sequence of G protein coupled receptor protein can be successfully carried out through polymerase chain reaction techniques with the said primer DNA. As a result, template DNAs coding for part or all of the polypeptide sequence of G protein coupled receptor protein, contained in the DNA library, can be selectively amplified and various DNA sequences encoding part or all of the

polypeptide sequence of G protein coupled receptor protein may be isolated and characterized. Further, G protein coupled receptor proteins, peptide segments or fragments derived from the G protein coupled receptor protein, modified derivatives or analogues thereof, and salts thereof may be recognized, predicted, deduced, produced, expressed, isolated and characterized.

The primer DNA useful in PCR amplification of the DNA sequence encoding part or all of the polypeptide sequence of G protein coupled receptor protein is a degenerate deoxynucleotide which has an oligonucleotide sequence to which a SEQ ID NO selected from the group consisting of SEQ ID NO: 1 to SEQ ID NO: 19 is assigned.

The nucleotide sequence represented by SEQ ID NO: 1 is a base sequence having the following formula:

5'-CGTGGSCMTSSTGGGCAACN₁YCCTG-3'

wherein S is G or C, M is A or C, N₁ = A, G, C, or T, and Y is T or C (Figure 1: HS-1).

The nucleotide sequence represented by SEQ ID NO: 2 (HS-2) is a base sequence having the following formula:

5'-GTN₁GWRRGGCAN₁CCAGCAGAKGGCAAA-3'

wherein N₁ = A, G, C, or T, W is A or T, R is A or G, and K is G or T, which is complementary to a nucleotide sequence having the following formula:

5'-TTTGCCMTCTGCTGGNTGCCYYWCNAC-3'

wherein N = A, C, G, or T, M is A or C, Y is T or C, and W is A or T (Figure 2).

The nucleotide sequence represented by SEQ ID NO: 3 is a base sequence having the following formula:

5'-CTCGCSGCYMTN₂RGYATGGAYCGN₂TAT-3'

wherein S is G or C, Y is C or T, M is A or C, R is A or G, and N₂ = I (Figure 4: 3D).

The nucleotide sequence represented by SEQ ID NO: 4 is a base sequence having the following formula:

5'-CATGTRGWAGGGAAN₂CCAGSAMAN₂RARRAA-3'

wherein R is A or G, W is T or A, S is G or C, M is A or C, and N₂ = I, which is complementary to a nucleotide sequence

having the following formula:




wherein Y is C or T, $N_1 = A, G, C, \text{ or } T$, K is G or T, S is G or C, W is A or T (Figure 6: 6C).

5 The nucleotide sequence represented by SEQ ID NO: 5 is a base sequence having the following formula:



wherein Y is C or T, R is A or G, S is G or C, M is A or C, and V is A, C or G, and N_2 is I (Figure 3: 3A).

10 The nucleotide sequence represented by SEQ ID NO: 6 is a base sequence having the following formula:




wherein Y is C or T, R is A or G, S is G or C, M is A or C, and V is A, C or G, and N_2 is I (Figure 3: 3B).

15 The nucleotide sequence represented by SEQ ID NO: 7 is a base sequence having the following formula:




wherein S is G or C, Y is C or T, M is A or C, R is A or G, and N_2 is I (Figure 4: 3C).

20 The nucleotide sequence represented by SEQ ID NO: 8 is a base sequence having the following formula:




wherein R is A or G, S is G or C, Y is C or T, and N_2 is I, which is complementary to a nucleotide sequence having the following formula:




wherein R is A or G, Y is C or T, S is G or C, and N_1 is A, T, G, or C (Figure 5: 6A).

30 The nucleotide sequence represented by SEQ ID NO: 9 is a base sequence having the following formula:



wherein R is A or G, S is G or C, Y is C or T, and N_2 is I, which is complementary to a nucleotide sequence having the following formula:



wherein R is A or G, Y is C or T, S is G or C, and N_1 is A,

T, G, or C (Figure 5: 6B).

The nucleotide sequence represented by SEQ ID NO: 10 is a base sequence having the following formula:

5'-GYCACCAACN₂WSTTCATCCTSWN₂HCTG-3'

5 wherein S is G or C, Y is C or T, W is A or T, H is A, C or T, and N₂ is I (Figure 7: T2A).

The nucleotide sequence represented by SEQ ID NO: 11 (Figure 8: T7A) is a base sequence having the following formula:

10 5'-ASN₂SAN₂RAAGSARTAGAN₂GAN₂RGGRTT-3'

wherein R is A or G, S is G or C, and N₂ is I, which is complementary to a nucleotide sequence having the following formula:

15 5'-AAYCCYN₂TCN₂TCTAYTSCTTYN₂TSN₂ST-3'

wherein Y is C or T, N₂ is I, and S is G or C (Figure 8).

The nucleotide sequence represented by SEQ ID NO: 12 is a base sequence having the following formula:

5'-TGN₂TSSTKMTN₂GSN₂GTKGTN₂GGN₂AA-3'

20 wherein S is G or C, K is G or T, M is A or C, and N₂ is I (Figure 9: TM1-A2).

The nucleotide sequence represented by SEQ ID NO: 13 (Figure 10: TM3-B2) is a base sequence having the following formula:

25 5'-AYCKGTAYCKGTCCAN₂KGWN₂ATKGC-3'

wherein Y is C or T, K is G or T, W is A or T, and N₂ is I, which is complementary to a nucleotide sequence having the following formula:

5'-GCMATN₂WCMN₂TGGACMGRTACMGRT-3'

30 wherein M is A or C, W is A or T, R is A or G, and N₂ is I (Figure 10).

The nucleotide sequence represented by SEQ ID NO: 14 is a base sequence having the following formula:

5'-CATKKCCSTGGASAGN₂TAYN₂TRGC-3'

35 wherein K is G or T, S is G or C, Y is C or T, R is A or G, and N₂ is I (Figure 11: TM3-C2).


The nucleotide sequence represented by SEQ ID NO: 15 (Figure 12: TM6-E2) is a base sequence having the following

formula:




wherein W is A or T, S is G or C, K is G or T, R is A or G, and N_2 is I, which is complementary to a nucleotide sequence

5 having the following formula:



wherein Y is C or T, S is G or C, M is A or C, W is A or T, and N_2 is I (Figure 12).

10 The nucleotide sequence represented by SEQ ID NO: 16 is a base sequence having the following formula:




wherein R is A or G, Y is C or T, N_1 is A, T, G, or C, and N_2 is I (Figure 13: TM2F18).

15 The nucleotide sequence represented by SEQ ID NO: 17 (Figure 14: TM6R21) is a base sequence having the following formula:



wherein R is A or G, N_1 is A, T, G, or C, and N_2 is I which is complementary to a nucleotide sequence having the
20 following formula:




wherein Y is C or T, N_1 is A, T, G, or C, and N_2 is I (Figure 14).

25 The nucleotide sequence represented by SEQ ID NO: 18 is a base sequence having the following formula:



wherein S is G or C, R is A or G, W is A or T, M is A or C, and N_2 is I (Figure 15: S3A).

30 The nucleotide sequence represented by SEQ ID NO: 19 (Figure 16: S6A) is a base sequence having the following formula:



wherein W is A or T, S is G or C, M is A or C, Y is C or T, R is A or G, and N_2 is I, which is complementary to a

35 nucleotide sequence having the following formula:



wherein W is A or T, Y is C or T, R is A or G, K is G or T, and S is G or C (Figure 16).

In a specific embodiment, symbols in the
aforementioned SEQ ID NOs (R, Y, M, K, S, W, H, V and N)
5 indicate the incorporation of plural bases, leading to multiple
oligonucleotides in the primer preparation. In other words,
SEQ ID NO: 1 to SEQ ID NO: 19 are degenerate nucleotide primers.

The nucleotide sequence represented by SEQ ID NO: 1
(Figure 1: HS-1) is a nucleotide sequence highly homologous to
10 the DNA sequence coding for the amino acid sequence
corresponding to or near the first membrane-spanning
(transmembrane) domain each of known G protein coupled receptor
proteins such as human-derived TRH receptor protein (HTRHR),
human-derived RANTES receptor protein (L10918, HUMRANTES),
15 human Burkitt's lymphoma-derived receptor protein with an
unknown ligand (X68149, HSBRL1A), human-derived
somatostatin receptor protein (L14856, HUMSOMAT0), rat-derived
 μ -opioid receptor protein (U02083, RNU02083), rat-derived
 κ -opioid receptor protein (U00442, U00442), human-derived
20 neuromedin B receptor protein (M73482, HUMNMBR),
human-derived muscarinic acetylcholine receptor
protein (X15266, HSHM4), rat-derived adrenaline
 α_1 B receptor protein (L08609, RATAADRE01), human-derived
somatostatin 3 receptor protein (M96738, HUMSSTR3X),
25 human-derived C_5a receptor protein (HUMC5AAR), human-derived
receptor protein with an unknown ligand (HUMRDC1A),
human-derived receptor protein with an unknown ligand
(M84605, HUMOPIODRE), rat-derived adrenaline
 α_2 B receptor protein (M91466, RATA2BAR) and the like
30 [Figure 1].

The nucleotide sequence represented by SEQ ID NO: 2
(HS-2) is a nucleotide sequence which is complementary to the
nucleotide sequence (Figure 2) highly homologous to the DNA
sequence coding for the amino acid sequence corresponding to or
35 near the sixth membrane-spanning domain of known G protein
coupled receptor proteins such as mouse-derived receptor

protein with an unknown ligand (M80481, MUSGIR), human-derived bombesin receptor protein (L08893, HUMBOMB3S), human-derived adenosine A2 receptor protein (S46950, S46950), mouse-derived receptor protein with an unknown ligand (D21061, MUSGPCR),
5 mouse-derived TRH receptor protein (S43387, S43387),
rat-derived neuromedin K receptor protein (J05189, RATNEURA),
rat-derived adenosine A1 receptor protein (M69045, RATA1ARA),
human-derived neurokinin A receptor protein (M57414, HUMNEKAR),
rat-derived adenosine A3 receptor protein (M94152, RATADENREC),
10 human-derived somatostatin 1 receptor protein (M81829, HUMSTRI1A), human-derived neurokinin 3 receptor protein (S86390, S86371S4), rat-derived receptor protein with an unknown ligand (X61496, RNCGPCR), human-derived somatostatin 4 receptor protein (L07061, HUMSSTR4Z), rat-derived GnRH
15 receptor protein (M31670, RATGNRHA) and the like [Figure 2].

The nucleotide sequence represented by SEQ ID NO: 5 (Figure 3: 3A) or the nucleotide sequence represented by SEQ ID NO: 6 (Figure 3: 3B) is a nucleotide sequence highly homologous to the DNA sequence coding for the amino acid
20 sequence corresponding to or near the third membrane-spanning domain each of known G protein coupled receptors such as mouse-derived κ -opioid receptor protein (L11064), mouse-derived δ -opioid receptor protein (L11065), rat-derived μ -opioid receptor protein (D16349), mouse-derived bradykinin
25 B2 receptor protein (X69676), rat-derived bradykinin B2 receptor protein (M59967), mouse-derived bombesin receptor protein (M35328), human-derived neuromedin B receptor protein (M73482), human-derived gastrin releasing peptide receptor protein (M73481), human-derived bombesin receptor protein
30 subtype 3 (L08893), mouse-derived substance K receptor protein (X62933), mouse-derived substance P receptor protein (X62934), rat-derived neurokinin 3 receptor protein (J05189), rat-derived endothelin receptor protein (M60786), rat-derived receptor protein with an unknown ligand (L04672), rat-derived
35 receptor protein with an unknown ligand (X61496), rat-derived receptor protein with an unknown ligand (X59249), rat-derived receptor protein with an unknown ligand (L09249),

- 83 -

mouse-derived receptor protein with an unknown ligand (P30731),
 human-derived receptor protein with an unknown ligand (M31210),
 and the like (Figure 3).

5 The nucleotide sequence represented by SEQ ID NO: 7
 (Figure 4: 3C) or the nucleotide sequence represented by SEQ
 ID NO: 3 (Figure 4: 3D) is a nucleotide sequence highly
 homologous to the DNA sequence coding for the amino acid
 sequence corresponding to or near the third membrane-spanning
 10 domain each of known G protein coupled receptors such
 as mouse-derived angiotensin II receptor protein (L32840),
 rat-derived angiotensin Ib receptor protein (X64052),
 human-derived angiotensin Ia receptor protein (M91464),
 15 rat-derived cholecystokinin a receptor protein (M88096),
 rat-derived cholecystokinin b receptor protein (M99418),
 human-derived low affinity interleukin 8 receptor protein
 mouse-derived high affinity interleukin 8 receptor
 20 (M73969), human-derived C5a anaphylatoxin receptor
 protein (X65858), mouse-derived N-formylpeptide receptor
 protein (S46665), human-derived C5a anaphylatoxin receptor
 protein (M60626) and the like (Figure 4).

The nucleotide sequence represented by SEQ ID NO: 10
 (Figure 7: T2A) is a nucleotide sequence highly homologous to
 the DNA sequence coding for the amino acid sequence
 25 corresponding to or near the second membrane-spanning domain
 each of known G protein coupled receptors such as
 human galanin receptor (HUMGALAREC), rat α -1B-adrenergic
 receptor (RATADR1B), human β -1-adrenergic receptor
 30 (HUMADRB1), rabbit IL-8 receptor (RABIL8RSB), human opioid
 receptor (HUMOPIODRE), bovine substance K receptor (BTSKR),
 human somatostatin receptor-2 (HUMSRI2A), human somatostatin
 receptor-3 (HUMSSTR3Y), human gastrin receptor (HUMGARE),
 35 human cholecystokinin A receptor (HUMCCKAR), human dopamine
 receptor-D5 (HUMD1B), human serotonin receptor 5HT1E
 (HUM5HT1E), human dopamine receptor D4 (HUMD4C), mouse
 serotonin receptor-2 (MMSERO), rat α -1A-adrenergic receptor

(RATADRALA), rat histamine H2 receptor (S57565) and the like [Figure 7].

The nucleotide sequence represented by SEQ ID NO: 8 (complementary to 6A of Figure 5) or the nucleotide sequence represented by SEQ ID NO: 9 (complementary to 6B of Figure 5) is a nucleotide sequence which is complementary to the nucleotide sequence (Figure 5) highly homologous to the DNA sequence coding for the amino acid sequence corresponding to or near the sixth membrane-spanning domain of known G protein coupled receptors such as mouse-derived κ -opioid receptor protein (L11064), mouse-derived δ -opioid receptor protein (L11065), rat-derived μ -opioid receptor protein (D16349), mouse-derived bradykinin B2 receptor protein (X69676), rat-derived bradykinin B2 receptor protein (M59967), mouse-derived bombesin receptor protein (M35328), human-derived neuromedin B receptor protein (M73482), human-derived gastrin releasing peptide receptor protein (M73481), human-derived bombesin receptor protein subtype 3 (L08893), mouse-derived substance K receptor protein (X62933), mouse-derived substance P receptor protein (X62934), rat-derived neurokinin 3 receptor protein (J05189), rat-derived endothelin receptor protein (M60786), rat-derived receptor protein with an unknown ligand (L04672), rat-derived receptor protein with an unknown ligand (X61496), rat-derived receptor protein with an unknown ligand (X59249), rat-derived receptor protein with an unknown ligand (L09249), mouse-derived receptor protein with an unknown ligand (P30731), human-derived receptor protein with an unknown ligand (M31210) human-derived receptor protein with an unknown ligand (U03642) and the like [Figure 5].

The nucleotide sequence represented by SEQ ID NO: 4 (complementary to 6C of Figure 6) is a nucleotide sequence which is complementary to the nucleotide sequence (Figure 6) highly homologous to the DNA sequence coding for the amino acid sequence corresponding to or near the sixth membrane-spanning domain of known G protein coupled receptors such as mouse-derived angiotensin II receptor protein (L32840),

rat-derived angiotensin Ib receptor protein (X64052),
rat-derived angiotensin receptor protein subtype (M90065),
human-derived angiotensin Ia receptor protein (M91464),
rat-derived cholecystokinin a receptor protein (M88096),
5 rat-derived cholecystokinin b receptor protein (M99418),
human-derived cholecystokinin 8 receptor protein (L04473),
mouse-derived low affinity interleukin 8 receptor protein
(M73969), human-derived high affinity interleukin 8 receptor
protein (X65858), mouse-derived C5a anaphylatoxin receptor
10 protein (S46665), human-derived N-formylpeptide receptor
protein (M60626) and the like [Figure 6].

The nucleotide sequence represented by SEQ ID NO: 11
(Figure 8: T7A) is a nucleotide sequence which is
complementary to the nucleotide sequence (Figure 8) highly
15 homologous to the DNA sequence coding for the amino acid
sequence corresponding to or near the seventh
membrane-spanning domain each of known G protein coupled
receptors such as human galanin receptor (HUMGALAREC),
rat A1 adenosine receptor (RAT1DREC), porcine angiotensin
20 receptor (PIGA2R), rat serotonin receptor (RAT5HTRTC),
human dopamine receptor (S58541), human gastrin releasing
peptide receptor (HUMGRPR), mouse GRP/bombesin receptor
(MUSGRPBOB), rat vascular type 1 angiotensin receptor
(RRVT1A1IR), human muscarinic acetylcholine receptor (HSHM4),
25 human β -1 adrenergic receptor (HUMDRB1), human gastrin
receptor (HUMGARE), rat cholecystokinin receptor (RATCCKAR),
rat receptor with an unknown ligand (S59748), human
somatostatin receptor (HUMSST28A), rat receptor with an unknown
ligand (RNGPROCR), mouse somatostatin receptor-1 (MUSSRI1A),
30 human α -A1-adrenergic receptor (HUMA1AADR), mouse
delta-opioid receptor (S66181), human somatostatin receptor-3
(HUMSSTR3Y) and the like [Figure 8].

The nucleotide sequence represented by SEQ ID NO: 12
(Figure 9: TM1-A2) is a nucleotide sequence highly homologous
35 to the DNA sequence coding for the amino acid sequence
within the first membrane-spanning (transmembrane) domain
each of known G protein coupled receptors such as

mouse-derived bradykinin B₂ receptor (MUSBB2R),
 bovine-derived substance K receptor (BTSKR), bovine-derived
 endothelin ET_B receptor (BOVEETBR), human-derived
 neuropeptide Y receptor (MMSUBKREC), human-derived
 5 prostaglandin E₂ receptor (HUMPGE2R), human-derived
 prostacyclin receptor (HUMPIR), human-derived κ -opioid
 receptor (HSU11053), rat-derived melanocortin 3 receptor
 (RRMC3RA), human-derived melanocortin receptor (HUMMR),
 mouse-derived bombesin/GRP receptor (MUSGRPBOM),
 10 rat-derived cholecystokinin B receptor (RATCHOLREC),
 rat-derived cholecystokinin A receptor (RATCCKAR) and the
 like [Figure 9].

The nucleotide sequence represented by SEQ ID NO: 13
 (Figure 10: TM3-B2) is a nucleotide sequence which
 15 is complementary to the nucleotide sequence (Figure 10) highly
 homologous to the DNA sequence coding for the amino acid
 sequence corresponding to or near the end of the third
 membrane-spanning domain of known G protein coupled receptors
 such as human-derived cholecystokinin receptor (HUMCCKR),
 20 human-derived cholecystokinin B receptor (HUMCCKBGR),
 mouse-derived melanocortin 5 receptor (MMGMC5R),
 human-derived vasopressin receptor (HUMV2R), rat-derived
 neuromedin K receptor (RATNEURA), dog-derived gastrin receptor
 (DOGGSTRN), rat-derived serotonin receptor (RAT5HT5A),
 25 mouse-derived α_2 -adrenaline receptor (MUSALP2ADA),
 human-derived adenosine A₁ receptor (HUMADORA1X),
 human-derived opioid (presumed) receptor (HUMOPIODRE),
 mouse-derived bombesin/GRP receptor (MUSGRPBOM),
 rat-derived cholecystokinin A receptor (RATCCKAR),
 30 human-derived TRH receptor (HSTRHREC) and the like [Figure 10].

The nucleotide sequence represented by SEQ ID NO: 14
 (Figure 11: TM3-C2) is a nucleotide sequence highly homologous
 to the DNA sequence coding for the amino acid sequence
 corresponding to or near the end of the third membrane-spanning
 35 domain of known G protein coupled receptors such as
 human-derived neurokinin 3 receptor (HUMNK3R), human-derived
 oxytocin receptor (HSMRNOXY), guinea pig-derived

cholecystokinin A receptor (S68242), dog-derived
cholecystokinin A receptor with an unknown ligand (CFGPCR4),
mouse-derived substance P receptor (MMSUBPREC), human-derived
receptor with an unknown ligand (HUMOPIODRE), human-derived
5 galanin receptor (HUMGALAREC), human-derived serotonin
receptor (HSS31G), human-derived β_3 -adrenaline receptor
(HUMARB3A), human-derived prostacyclin receptor (HUMHPR),
rat-derived cholecystokinin A receptor (RATCCKAR) and the
like [Figure 11].

10 The nucleotide sequence represented by SEQ ID NO: 15
(Figure 12: TM6-E2) is a nucleotide sequence which
is complementary to the nucleotide sequence (Figure 12) highly
homologous to the DNA sequence coding for the amino acid
sequence within the sixth membrane-spanning domain of known
15 G protein coupled receptors such as human-derived neurokinin A
receptor (HUMNEKAR), human-derived substance P receptor
(HUMSUBPRA), rat-derived substance K receptor (RATSKR),
mouse-derived bombesin/GRP receptor (MUSGRPBOM),
human-derived opioid (presumed) receptor (HUMOPIODRE),
20 human-derived adenosine A₂ receptor (HUMA2XXX),
human-derived β_2 -adrenaline receptor (HUMADRBR),
canine-derived receptor RDC5 with an unknown ligand (CFGPCR8),
human-derived endothelin receptor (HUMETSR), mouse-derived
neuropeptide Y1 receptor (MMNPY1CDS), human-derived oxytocin
25 receptor (HSMRNOXY), rat-derived cholecystokinin A receptor
(RATCCKAR) and the like [Figure 12].

 The nucleotide sequence represented by SEQ ID NO: 16
(Figure 13: TM2F18) is a nucleotide sequence highly homologous
to the DNA sequence coding for the amino acid sequence
30 corresponding to or near the second membrane-spanning domain
of known G protein coupled receptors such as human-derived TSH
receptor (HUMTSHX), human-derived neurokinin A receptor
(HUMNEKAR), human-derived FMLP receptor (HUMFMLP),
human-derived IL8 receptor B (HUMINTLEU8), human-derived
35 α -A1 adrenergic receptor (HUMA1AADR), human-derived IL8
receptor A (HUMIL8RA), human-derived dopamine D2 receptor
(HSDD2), human-derived angiotensin type I receptor (HUMANTIR),

human-derived somatostatin receptor (HUSOMAT), human-derived TRH receptor (HSTRHREC), human-derived delta-opioid receptor (HSUO7882) and the like [Figure 13].

The nucleotide sequence represented by SEQ ID NO: 17 (Figure 14: TM6R21) is a nucleotide sequence which is complementary to the nucleotide sequence (Figure 14) highly homologous to the DNA sequence coding for the amino acid sequence corresponding to or near the sixth membrane-spanning domain of known G protein coupled receptors such as human-derived β -adrenergic receptor (HSBAR), human-derived neurokinin A receptor (HUMNEKAR), human-derived endothelin-1 receptor (HUMETN1R), human-derived histamine H_2 receptor (HUMHISH2R), human-derived α -A1 adrenergic receptor (HUMA1AADR), human-derived IL8 receptor A (HUMIL8RA), human-derived neuromedin B receptor (HUMNMBR), human-derived neurokinin 1 receptor (HUMNKIRX), human-derived substance P receptor (HUMSUBPRA), human-derived 5-HT1D serotonin receptor (HUM5HT1DA), human-derived formylpeptide receptor (HUMPFPR2A), human-derived dopamine D2 receptor (HSDD2), human-derived neuropeptide Y receptor (HUMNEUYREC), human-derived adenosine A2 receptor (HUMA2XXX), human-derived bradykinin receptor BK-2 (HUMBK2A), human-derived FMLP-related receptor II (HUMFMLPX), human-derived somatostatin receptor subtype 3 (HUMSSTR3X), human-derived cholecystokinin receptor (HUMCCKR), human-derived neurotensin receptor (HSNEURA) and the like [Figure 14].

The nucleotide sequence represented by SEQ ID NO: 18 (Figure 15: S3A) is a nucleotide sequence highly homologous to the DNA sequence coding for the amino acid sequence corresponding to or near the third membrane-spanning domain of known G protein coupled receptors such as human-derived galanin receptor (HUMGALAREC), human-derived CCK-B receptor (S70057), human-derived ET_A receptor (S67127), human-derived ET_B receptor (S44866), human-derived C5A receptor (HUMC5AAR), human-derived angiotensin II receptor (HUMANTIR), human-derived bradykinin receptor (HUMBK2R), human-derived neurotensin receptor (HSNEURA), human-derived GRP receptor (HUMGRPR), human-derived somatostatin 5 receptor (HUMFSRS),

human-derived IL-8 receptor (HUMIL8RA), human-derived neurokinin 2 (neurokinin A) receptor (HUMNEKAR) and the like [Figure 15].

5 The nucleotide sequence represented by SEQ ID NO: 19 (Figure 16: S6A) is a nucleotide sequence which is complementary to the nucleotide sequence (Figure 16) highly homologous to the DNA sequence coding for the amino acid sequence corresponding to or near the sixth membrane-spanning domain of known G protein coupled receptors such as human-
10 derived galanin receptor (HUMGLAREC), human-derived CCK-B receptor (S70057), human-derived ET_A receptor (S67127), human-derived ET_B receptor (S44866), human-derived C5A receptor (HUMC5AAR), human-derived angiotensin II receptor (HUMANTIR), human-derived bradykinin receptor (HUMBK2R), human-derived
15 neurotensin receptor (HSNEURA), human-derived GRP receptor (HUMGRPR), human-derived somatostatin 5 receptor (HUMFSRS), human-derived IL-8 receptor (HUMIL8RA), human-derived neurokinin 2 (neurokinin A) receptor (HUMNEKAR) and the like [Figure 16].

20 The above-mentioned abbreviations in the parentheses are the identifiers (or reference numbers) which are shown when GenBank/EMBL Data Bank is searched using a DNASIS Gene/Protein Sequence Data Base (CD019; Hitachi Software Engineering, Japan) and are usually called "Accession Numbers" or "Entry Names".
25 HTRHR is, however, the sequence as described in Japanese Patent Application No. Hei 5-286986 (or No. 286986/1993) (EPA 638645).

The DNA (or nucleotides) of the present invention may be manufactured by DNA synthetic methods which are known
30 per se or by methods similar thereto. The DNA (or nucleotides) of the present invention may be an oligonucleotide sequence having 8 to 60 base residues, preferably 12 to 50 base residues, more preferably 15 to 40 residues and most preferably 18 to 30 residues.

35 Among the DNAs of the present invention, the DNA having the nucleotide sequence represented by SEQ ID NO: 1 or

SEQ ID NO: 12 is a nucleotide sequence which is commonly present in the nucleotide sequence of the DNA encoding the amino acid sequence corresponding to or near the first membrane-spanning domain of the above-mentioned known G protein coupled receptor protein. Therefore, it can be complementarily bonded (i.e. is hybridizable) with RNA or DNA (including genome DNA, cDNA) coding for the amino acid sequence corresponding to or near the first membrane-spanning domain of known or unknown G protein coupled receptor proteins and, furthermore, it can be complementarily bonded (i.e. is hybridizable) with nucleotide sequences encoding other membrane-spanning domains as well.

The DNA having a nucleotide sequence represented by SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 14 or SEQ ID NO: 18 is a nucleotide sequence which is commonly present in the nucleotide sequence of the DNA coding for the amino acid sequence corresponding to or near the third membrane-spanning domain of the above-mentioned known G protein coupled receptor protein. Therefore, it can be complementarily bonded with RNA or DNA (including genome DNA, cDNA) coding for the part corresponding to or near the third membrane-spanning domain of known or unknown G protein coupled receptor proteins and, furthermore, it can be complementarily bonded with nucleotide sequences encoding other membrane-spanning domains as well.

The DNA having a nucleotide sequence represented by SEQ ID NO: 10 or SEQ ID NO: 16 is a nucleotide sequence which is commonly present in the nucleotide sequence of the DNA coding for the amino acid sequence corresponding to or near the second membrane-spanning domain of the above-mentioned known G protein coupled receptor protein. Therefore, it can be complementarily bonded with RNA or DNA (including genome DNA, cDNA) coding for the part corresponding to or near the second membrane-spanning domain of known or unknown G protein coupled receptor proteins and, furthermore, it can be complementarily bonded with nucleotide sequences encoding other membrane-spanning domains as well.

The DNA having a nucleotide sequence represented by
SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID
NO: 15, SEQ ID NO: 17 or SEQ ID NO: 19 is a nucleotide sequence
which is commonly present in the nucleotide sequence of the DNA
coding for the amino acid sequence corresponding to or near
the sixth membrane-spanning domain of the above-mentioned known
G protein coupled receptor protein. Therefore, it can be
complementarily bonded with RNA or DNA (including genome DNA,
cDNA) coding for the part corresponding to or near the sixth
membrane-spanning domain of known or unknown G protein coupled
receptor proteins and, furthermore, it can be complementarily
bonded with nucleotide sequences encoding other
membrane-spanning domains as well.

The DNA having a nucleotide sequence represented by
SEQ ID NO: 11 is a nucleotide sequence which is commonly
present in the nucleotide sequence of the DNA coding for the
amino acid sequence corresponding to or near the seventh
membrane-spanning domain of the above-mentioned known G protein
coupled receptor protein. Therefore, it can be complementarily
bonded with RNA or DNA (including genome DNA, cDNA) coding for
the part corresponding to or near the seventh membrane-spanning
domain of known or unknown G protein coupled receptor proteins
and, further more, it can be complementarily bonded with
nucleotide sequences encoding other transmembrane domains
as well.

The DNA having a nucleotide sequence represented by
SEQ ID NO: 13 is a nucleotide sequence which is commonly
present in the nucleotide sequence of the DNA coding for the
amino acid sequence corresponding to or near the third
membrane-spanning domain of the above-mentioned known G protein
coupled receptor protein. Therefore, it can be complementarily
bonded with RNA or DNA (including genome DNA, cDNA) coding for
the part corresponding to or near the third membrane-spanning
domain of known or unknown G protein coupled receptor proteins
and, furthermore, it can be complementarily bonded with
nucleotide sequences encoding other membrane-spanning domains
as well.

Accordingly, the DNAs (or nucleotides) of the present invention can be used as DNA primers for a polymerase chain reaction (hereinafter, sometimes referred to as PCR).

For example:

- 5 (i) a polymerase chain reaction is carried out by mixing
 - (1) a small amount of DNA (or DNA fragment(s)) which codes for G protein coupled receptor protein, said DNA (or DNA fragment(s)) acting as a template,
 - 10 (2) at least one DNA primer selected from the group consisting of DNA primers having a nucleotide sequence represented by SEQ ID NO: 1, DNA primers having a nucleotide sequence represented by SEQ ID NO: 3, DNA primers having a nucleotide sequence represented by SEQ
15 ID NO: 5, DNA primers having a nucleotide sequence represented by SEQ ID NO: 6, DNA primers having a nucleotide sequence represented by SEQ ID NO: 7, DNA primers having a nucleotide sequence represented by SEQ
20 ID NO: 10, DNA primers having a nucleotide sequence represented by SEQ ID NO: 12, DNA primers having a nucleotide sequence represented by SEQ ID NO: 14, DNA primers having a nucleotide sequence represented by SEQ ID NO: 16 and DNA primers having a nucleotide sequence represented by SEQ ID NO: 18 and
 - 25 (3) at least one DNA primer selected from the group consisting of DNA primers having a nucleotide sequence represented by SEQ ID NO: 2, DNA primers having a nucleotide sequence represented by SEQ ID NO: 4, DNA primers having a nucleotide sequence represented by SEQ
30 ID NO: 8, DNA primers having a nucleotide sequence represented by SEQ ID NO: 9, DNA primers having a nucleotide sequence represented by SEQ ID NO: 11, DNA primers having a nucleotide sequence represented by SEQ ID NO: 15, DNA primers having a nucleotide sequence represented by SEQ ID NO: 17 and DNA primers having a nucleotide sequence represented by SEQ ID NO: 19; or
 - 35 (ii) a polymerase chain reaction is carried out by mixing

(1) a small amount of DNA (or DNA fragment(s)) coding for G protein coupled receptor protein, said DNA (or DNA fragment(s)) acting as a template,

(2) at least one DNA primer selected from the group consisting of DNA primers having a nucleotide sequence represented by SEQ ID NO: 1 and DNA primers having a nucleotide sequence represented by SEQ ID NO: 12 and

(3) at least one DNA primer selected from the group consisting of DNA primers having a nucleotide sequence represented by SEQ ID NO: 13

so that it is possible to amplify the target DNA (or DNA fragment(s)) coding for said receptor protein.

When the PCR is carried out using at least one DNA primer selected from the group consisting of DNA primers having a nucleotide sequence represented by SEQ ID NO: 2, DNA primers having a nucleotide sequence represented by SEQ ID NO: 4, DNA primers having a nucleotide sequence represented by SEQ ID NO: 8, DNA primers having a nucleotide sequence represented by SEQ ID NO: 9, DNA primers having a nucleotide sequence represented by SEQ ID NO: 11, DNA primers having a nucleotide sequence represented by SEQ ID NO: 15, DNA primers having a nucleotide sequence represented by SEQ ID NO: 17 and DNA primers having a nucleotide sequence represented by SEQ ID NO: 19, said DNA primer(s) is(are) bonded (hybridized) with the nucleotide sequence at the 3'-side of the + chain (plus chain) of template RNA or DNA (or fragment(s) thereof) coding for the sixth membrane-spanning domain or other membrane-spanning domains of G protein coupled receptor protein whereupon an elongation of the - chain (minus chain) proceeds in the 5' → 3' direction.

When the PCR is carried out using at least one DNA primer selected from the group consisting of DNA primers having a nucleotide sequence represented by SEQ ID NO: 11, said DNA primer is bonded with the nucleotide sequence at the 3'-side of the + chain (plus chain) of template RNA or DNA (or fragment(s) thereof) coding for the seventh membrane-spanning domain or other membrane-spanning domains of the G protein

coupled receptor protein whereupon an elongation of the - chain (minus chain) proceeds in the 5' - 3' direction.

When the PCR is carried out using at least one DNA primer selected from the group consisting of DNA primers having a nucleotide sequence represented by SEQ ID NO: 1 and DNA
5 primers having a nucleotide sequence represented by SEQ ID NO: 12, said DNA primer is bonded with the nucleotide sequence at the 3'-side of the - chain (minus chain) of template RNA or DNA (or fragment(s) thereof) coding for the first membrane-
10 spanning domain or other membrane-spanning domains of G protein coupled receptor protein whereupon an elongation of the + chain (plus chain) proceeds in the 5' - 3' direction.

When the PCR is carried out using at least one DNA primer selected from the group consisting of DNA primers having a nucleotide sequence represented by SEQ ID NO: 10 and DNA
15 primers having a nucleotide sequence represented by SEQ ID NO: 16, said DNA primer is bonded with the nucleotide sequence at the 3'-side of the - chain (minus chain) of template RNA or DNA (or fragment(s) thereof) coding for the second membrane-
20 spanning domain or other membrane-spanning domains of G protein coupled receptor protein whereupon an elongation of the + chain (plus chain) proceeds in the 5' - 3' direction.

When the PCR is carried out using at least one DNA primer selected from the group consisting of DNA primers
25 having a nucleotide sequence represented by SEQ ID NO: 3, DNA primers having a nucleotide sequence represented by SEQ ID NO: 5, DNA primers having a nucleotide sequence represented by SEQ ID NO: 6, DNA primers having a nucleotide sequence represented by SEQ ID NO: 7, DNA primers having a nucleotide
30 sequence represented by SEQ ID NO: 14 and DNA primers having a nucleotide sequence represented by SEQ ID NO: 18, said DNA primer is bonded with the nucleotide sequence at the 3'-side of the - chain (minus chain) of template RNA or DNA (or fragment(s) thereof) coding for the third membrane-spanning
35 domain or other membrane-spanning domains of G protein coupled receptor protein whereupon an elongation of the + chain (plus chain) proceeds in the 5' - 3' direction.

Accordingly, when the DNA primers having nucleotide sequences represented by any of SEQ ID NO: 1 to SEQ ID NO: 19 of the present invention are used in combination each other, DNA (or DNA fragment(s)) coding for G protein coupled receptor protein can be successfully amplified.

One embodiment of the present invention provides:

(A) a method of amplifying DNA coding for the G protein coupled receptor protein (e.g., from the first to sixth membrane-spanning (transmembrane) domains or other segments of the G protein coupled receptor protein), characterized in that a polymerase chain reaction is carried out by mixing

① a DNA coding for the G protein coupled receptor protein, said DNA acting as a template,

② at least one DNA primer selected from the group consisting of DNA primers having a nucleotide sequence represented by SEQ ID NO: 1 and DNA primers having a nucleotide sequence represented by SEQ ID NO: 12 and

③ at least one DNA primer selected from the group consisting of DNA primers having a nucleotide sequence represented by SEQ ID NO: 2, DNA primers having a nucleotide sequence represented by SEQ ID NO: 4, DNA primers having a nucleotide sequence represented by SEQ ID NO: 8, DNA primers having a nucleotide sequence represented by SEQ ID NO: 9, DNA primers having a nucleotide sequence represented by SEQ ID NO: 15, DNA primers having a nucleotide sequence represented by SEQ ID NO: 17 and DNA primers having a nucleotide sequence represented by SEQ ID NO: 19;

(B) a method of amplifying DNA coding for the G protein coupled receptor protein (e.g., from the first to seventh membrane-spanning (transmembrane) domains or other segments of the G protein coupled receptor protein), characterized in that a polymerase chain reaction is carried out by mixing

① a DNA coding for the G protein coupled receptor protein, said DNA acting as a template,

② at least one DNA primer selected from the group consisting of DNA primers having a nucleotide sequence represented by SEQ ID NO: 1 and DNA primers having a

nucleotide sequence represented by SEQ ID NO: 12 and

③ at least one DNA primer selected from the group consisting of DNA primers having a nucleotide sequence represented by SEQ ID NO:11;

5 (C) a method of amplifying a DNA coding for the G protein coupled receptor protein (e.g., from the second to sixth membrane-spanning (transmembrane) domains or other segments of the G protein coupled receptor protein), characterized in that a polymerase chain reaction is carried out by mixing

10 ① a DNA coding for the G protein coupled receptor protein, said DNA acting as a template,

② at least one DNA primer selected from the group consisting of DNA primers having a nucleotide sequence represented by SEQ ID NO: 10 and DNA primers having a
15 nucleotide sequence represented by SEQ ID NO: 16 and

③ at least one DNA primer selected from the group consisting of DNA primers having a nucleotide sequence represented by SEQ ID NO: 2, DNA primers having a nucleotide sequence represented by SEQ ID NO: 4, DNA primers having a
20 nucleotide sequence represented by SEQ ID NO: 8, DNA primers having a nucleotide sequence represented by SEQ ID NO: 9, DNA primers having a nucleotide sequence represented by SEQ ID NO: 15, DNA primers having a nucleotide sequence represented by SEQ ID NO: 17 and DNA primers having a
25 nucleotide sequence represented by SEQ ID NO: 19;

(D) a method of amplifying a DNA coding for the G protein coupled receptor protein (e.g., from the second to seventh membrane-spanning (transmembrane) domains or other segments of the G protein coupled receptor protein), characterized in
30 that a polymerase chain reaction is carried out by mixing

① a DNA coding for the G protein coupled receptor protein, said DNA acting as a template,

② at least one DNA primer selected from the group consisting of DNA primers having a nucleotide sequence
35 represented by SEQ ID NO: 10 and DNA primers having a nucleotide sequence represented by SEQ ID NO: 16 and

③ at least one DNA primer selected from the group

consisting of DNA primers having a nucleotide sequence represented by SEQ ID NO: 11;

(E) a method of amplifying a DNA coding for the G protein coupled receptor protein (e.g., from the third to sixth membrane-spanning (transmembrane) domains or other segments of the G protein coupled receptor protein), characterized in that a polymerase chain reaction is carried out by mixing

① a DNA coding for the G protein coupled receptor protein, said DNA acting as a template,

② at least one DNA primer selected from the group consisting of DNA primers having a nucleotide sequence represented by SEQ ID NO: 3, DNA primers having a nucleotide sequence represented by SEE ID NO: 5, DNA primers having a nucleotide sequence represented by SEQ ID NO: 6, DNA primers having a nucleotide sequence represented by SEQ ID NO: 7, DNA primers having a nucleotide sequence represented by SEQ ID NO: 14 and DNA primers having a nucleotide sequence represented by SEQ ID NO: 18 and

③ at least one DNA primer selected from the group consisting of DNA primers having a nucleotide sequence represented by SEQ ID NO: 2, DNA primers having a nucleotide sequence represented by SEQ ID NO: 4, DNA primers having a nucleotide sequence represented by SEQ ID NO: 8, DNA primers having a nucleotide sequence represented by SEQ ID NO: 9, DNA primers having a nucleotide sequence represented by SEQ ID NO: 15, DNA primers having a nucleotide sequence represented by SEQ ID NO: 17 and DNA primers having a nucleotide sequence represented by SEQ ID NO: 19;

(F) a method of amplifying a DNA coding for the G protein coupled receptor protein (e.g., from the third to seventh membrane-spanning (transmembrane) domains or other segments of the G protein coupled receptor protein), characterized in that a polymerase chain reaction is carried out by mixing

① a DNA coding for the G protein coupled receptor protein, said DNA acting as a template,

② at least one DNA primer selected from the group consisting of DNA primers having a nucleotide sequence

represented by SEQ ID NO: 3, DNA primers having a nucleotide sequence represented by SEQ ID NO: 5, DNA primers having a nucleotide sequence represented by SEQ ID NO: 6, DNA primers having a nucleotide sequence represented by SEQ ID NO: 7, DNA primers having a nucleotide sequence represented by SEQ ID NO: 14 and DNA primers having a nucleotide sequence represented by SEQ ID NO: 18 and

③ at least one DNA primer selected from the group consisting of DNA primers having a nucleotide sequence represented by SEQ ID NO:11; and

(G) a method of amplifying a DNA coding for the G protein coupled receptor protein (e.g., from the first to third membrane-spanning (transmembrane) domains or other segments of the G protein coupled receptor protein), characterized in that a polymerase chain reaction is carried out by mixing

① a DNA coding for the G protein coupled receptor protein, said DNA acting as a template,

② at least one DNA primer selected from the group consisting of DNA primers having a nucleotide sequence represented by SEQ ID NO: 1 and DNA primers having a nucleotide sequence represented by SEQ ID NO: 12 and

③ at least one DNA primer selected from the group consisting of DNA primers having a nucleotide sequence represented by SEQ ID NO: 13.

An example of more preferred combination of the DNA primers in the amplification according to the above-mentioned (A) includes a combination of a DNA primer having a nucleotide sequence represented by SEQ ID NO: 1 with a DNA primer having a nucleotide sequence represented by SEQ ID NO: 2 and the like.

An example of more preferred combination of the DNA primers in the amplification according to the above-mentioned (D) includes a combination of a DNA primer having a nucleotide sequence represented by SEQ ID NO: 10 with a DNA primer having a nucleotide sequence represented by SEQ ID NO: 11 and the like.

An example of more preferred combination of the DNA

primers in the amplification according to the above-mentioned (E) includes:

- (i) a combination of a DNA primer having a nucleotide sequence represented by SEQ ID NO: 5 or a DNA primer having a nucleotide sequence represented by SEQ ID NO: 6 with a DNA primer having a nucleotide sequence represented by SEQ ID NO: 8 or a DNA primer having a nucleotide sequence represented by SEQ ID NO: 9;
- (ii) a combination of a DNA primer having a nucleotide sequence represented by SEQ ID NO: 3 or a DNA primer having a nucleotide sequence represented by SEQ ID NO: 7 with a DNA primer having a nucleotide sequence represented by SEQ ID NO: 4 and the like.

An example of more preferred combination of the DNA primers in the amplification according to the above-mentioned (G) includes a combination of a DNA primer having a nucleotide sequence represented by SEQ ID NO: 12 with a DNA primer having a nucleotide sequence represented by SEQ ID NO: 13 and the like.

The amplification may be carried out in accordance with known PCR techniques. For example, it may be carried out by the method described in Saiki, R. K. et al., Science, 239:487-491 (1988). Temperature, time, buffer, number of reaction cycles, enzyme such as DNA polymerase, addition of 2'-deoxy-7-deazaguanosine triphosphate or inosine, etc. in the PCR amplification may be suitably selected depending upon the type of target DNA and other factors. When RNA is used as a template, PCR amplification may be carried out, for example, by the method described in Saiki, R. K. et al., Science, 239:487-491(1988).

Moreover, the DNA having a nucleotide sequence represented by SEQ ID NO: 1 or SEQ ID NO: 12 of the present invention can be selectively and complementarily bonded (hybridized) with the nucleotide sequence at the 3'-side of the - chain of the DNA coding for the amino acid sequence corresponding to or near the first membrane-spanning domain of the G protein coupled receptor protein; the DNA having a nucleotide sequence represented by SEQ ID NO: 10 or SEQ ID

NO: 16 of the present invention can be selectively and complementarily bonded (hybridized) with the nucleotide sequence at the 3'-side of the - chain of the DNA coding for the amino acid sequence corresponding to or near the second
5 membrane-spanning domain of the G protein coupled receptor protein; the DNA having a nucleotide sequence represented by SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 14 or SEQ ID NO: 18 of the present invention can be selectively and complementarily bonded (hybridized) with the
10 nucleotide sequence at the 3'-side of the - chain of the DNA coding for the amino acid sequence corresponding to or near the third membrane-spanning domain of the G protein coupled receptor protein; the DNA having a nucleotide sequence represented by SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 15, SEQ ID NO: 17 or SEQ ID NO: 19
15 of the present invention can be selectively and complementarily bonded (hybridized) with the nucleotide sequence at the 3'-side of the + chain of the DNA coding for the amino acid sequence corresponding to or near the sixth membrane-spanning domain of the G protein coupled receptor protein; the DNA having a nucleotide sequence represented by SEQ ID NO: 11 of the present invention can be selectively and complementarily bonded (hybridized) with the nucleotide sequence at the
20 3'-side of the + chain of the DNA coding for the amino acid sequence corresponding to or near the third membrane-spanning domain of the G protein coupled receptor protein; and the DNA having a nucleotide sequence represented by SEQ ID NO: 13 of the present invention can be selectively and complementarily bonded (hybridized) with the nucleotide sequence at the
25 3'-side of the + chain of the DNA coding for the amino acid sequence corresponding to or near the third membrane-spanning domain of the G protein coupled receptor protein and, accordingly, said DNA is also advantageously useful as a probe for screening DNA libraries for DNA (or DNA fragment(s))
30 encoding part or all of the polypeptide sequence of G protein coupled receptor proteins.

These screening methods for DNA (or DNA fragment(s))

encoding part or all of the polypeptide sequence of G protein coupled receptor proteins from the DNA library by using as a reagent, because it can be used as a probe the DNA of the present invention may be carried out according to DNA cloning methods known per se by those of skill in the art or methods similar thereto. Especially when the DNA of the present invention is used as a DNA primer for the PCR, both amplification and screening of the DNA (or DNA fragment) coding for the G protein coupled receptor protein can be conducted in a single step.

Thus, when the DNAs of the present invention are suitably combined and used as the DNA primer for the PCR, said DNA primer(s) is(are) bonded (hybridized) with RNA or DNA (or fragment(s) thereof) encoding the amino acid sequence of the first membrane-spanning (transmembrane) domain, the second membrane-spanning domain, the third membrane-spanning domain, the sixth membrane-spanning domain, the seventh membrane-spanning domain or other membrane-spanning domains of G protein coupled receptor proteins to amplify, for example,

① RNA or DNA (or fragment(s) thereof) coding for the amino acid sequence of from the first membrane-spanning to the sixth membrane-spanning domains of G protein coupled receptor proteins,

② RNA or DNA (or fragment(s) thereof) coding for the amino acid sequence of from the first membrane-spanning to the seventh membrane-spanning domains of G protein coupled receptor proteins,

③ RNA or DNA (or fragment(s) thereof) coding for the amino acid sequence of from the third membrane-spanning to the sixth membrane-spanning domains of G protein coupled receptor proteins,

④ RNA or DNA (or fragment(s) thereof) coding for the amino acid sequence of from the third membrane-spanning to the seventh membrane-spanning domains of G protein coupled receptor proteins,

⑤ RNA or DNA (or fragment(s) thereof) coding for the amino acid sequence of from the second membrane-spanning to the

sixth membrane-spanning domains of G protein coupled receptor proteins or RNA or DNA (or fragment(s) thereof) coding for the amino acid sequence of other domains thereof,

⑥ RNA or DNA (or fragment(s) thereof) coding for the amino acid sequence of from the second membrane-spanning to the seventh membrane-spanning domains of G protein coupled receptor proteins,

⑦ RNA or DNA (or fragment(s) thereof) coding for the amino acid sequence of from the first membrane-spanning to the third membrane-spanning domains of G protein coupled receptor proteins or

⑧ RNA or DNA (or fragment(s) thereof) coding for the amino acid sequence of other domains of G protein coupled receptor proteins.

15 Through using the DNA primer according to the present invention, therefore, selective amplifications of:

① RNA or DNA (or fragment(s) thereof) coding for the amino acid sequence covering from the first membrane-spanning domain to the sixth membrane-spanning domain of G protein coupled receptor proteins;

② RNA or DNA (or fragment(s) thereof) coding for the amino acid sequence covering from the first membrane-spanning domain to the seventh membrane-spanning domain of G protein coupled receptor proteins;

25 ③ RNA or DNA (or fragment(s) thereof) coding for the amino acid sequence covering from the third membrane-spanning domain to the sixth membrane-spanning domain of G protein coupled receptor proteins;

30 ④ RNA or DNA (or fragment(s) thereof) coding for the amino acid sequence covering from the third membrane-spanning domain to the seventh membrane-spanning domain of G protein coupled receptor proteins;

35 ⑤ RNA or DNA (or fragment(s) thereof) coding for the amino acid sequence covering from the second membrane-spanning domain to the sixth membrane-spanning domain of G protein coupled receptor proteins or RNA or DNA (or fragment(s) thereof) coding for the amino acid sequence covering other areas thereof,

⑥ RNA or DNA (or fragment(s) thereof) coding for the amino acid sequence covering from the second membrane-spanning domain to the seventh membrane-spanning domain of G protein coupled receptor proteins;

- 5 ⑦ RNA or DNA (or fragment(s) thereof) coding for the amino acid sequence covering from the first membrane-spanning domain to the third membrane-spanning domain of G protein coupled receptor proteins; and the like,
from DNA libraries can be successfully achieved.

10 Among the DNA primers of the present invention, the combination of

① a DNA primer having a nucleotide sequence represented by SEQ ID NO: 1 or SEQ ID NO: 2; with

- ② at least one DNA primer selected from the group consisting
15 of a DNA primer having a nucleotide sequence represented by SEQ ID NO: 2, a DNA primer having a nucleotide sequence represented by SEQ ID NO: 4, a DNA primer having a nucleotide sequence represented by SEQ ID NO: 8, a DNA primer having a
20 having a nucleotide sequence represented by SEQ ID NO: 15, a DNA primer having a nucleotide sequence represented by SEQ ID NO: 17 and a DNA primer having a nucleotide sequence represented by SEQ ID NO: 19;

is, unlike conventional primers, capable of selectively
25 amplifying a broad area covering from the first membrane-spanning domain to the sixth membrane-spanning domain or other domains of G protein coupled receptor proteins.

 Among the DNA primers of the present invention, the combination of

- 30 ① a DNA primer having a nucleotide sequence represented by SEQ ID NO: 1 or SEQ ID NO: 12; with

② a DNA primer having a nucleotide sequence represented by SEQ ID NO: 11;

is, unlike conventional primers, capable of selectively
35 amplifying a broad area covering from the first membrane-spanning domain to the seventh membrane-spanning domain or other domains of G protein coupled receptor proteins.

Among the DNA primers of the present invention, the combination of

- ① a DNA primer having a nucleotide sequence represented by SEQ ID NO: 10 or SEQ ID NO: 16; with
- 5 ② at least one DNA primer selected from the group consisting of a DNA primer having a nucleotide sequence represented by SEQ ID NO: 2, a DNA primer having a nucleotide sequence represented by SEQ ID NO: 4, a DNA primer having a nucleotide sequence represented by SEQ ID NO: 8, a DNA primer having a nucleotide
- 10 sequence represented by SEQ ID NO: 9, a DNA primer having a nucleotide sequence represented by SEQ ID NO: 15, a DNA primer having a nucleotide sequence represented by SEQ ID NO: 17 and a DNA primer having a nucleotide sequence represented by SEQ ID NO: 19;
- 15 is, unlike conventional primers, capable of selectively amplifying a broad area covering from the second membrane-spanning domain to the sixth membrane-spanning domain or other domains of G protein coupled receptor proteins.

Among the DNA primers of the present invention, the combination of

- 20 ① a DNA primer having a nucleotide sequence represented by SEQ ID NO:10 or SEQ ID NO:16; with
- ② a DNA primer having a nucleotide sequence represented by SEQ ID NO:11;
- 25 is, unlike conventional primers, capable of selectively amplifying a broad area covering from the second membrane-spanning domain to the seventh membrane-spanning domain or other domains of G protein coupled receptor proteins.

Among the DNA primers of the present invention, the combination of

- 30 ① at least one DNA primer selected from the group consisting of a DNA primer having a nucleotide sequence represented by SEQ ID NO: 3, a DNA primer having a nucleotide sequence represented by SEQ ID NO: 5, a DNA primer having a nucleotide sequence represented by SEQ ID NO: 6, a DNA primer having a nucleotide
- 35 sequence represented by SEQ ID NO: 7, a DNA primer having a nucleotide sequence represented by SEQ ID NO: 14 and a DNA

primer having a nucleotide sequence represented by SEQ ID NO: 18; with

② a DNA primer having a nucleotide sequence represented by SEQ ID NO: 11;

5 is, unlike conventional primers, capable of selectively amplifying a broad area covering from the third membrane-spanning domain to the seventh membrane-spanning domain or other domains of G protein coupled receptor proteins.

Therefore, the protein hydrophobicity plotting of G
10 protein coupled receptor proteins and the homology at the amino acid level or the nucleic acid level between G protein coupled receptor proteins and other similar receptor proteins [said hydrophobicity plotting and homology both serve as standards
15 thereof) obtained according to the present invention is(are) encoding part or all of the amino acid sequence of G protein coupled receptor protein] can now be more clearly calculated.

Among the DNA primers of the present invention, the combination of

20 ① at least one DNA primer selected from the group consisting of a DNA primer having a nucleotide sequence represented by SEQ ID NO: 3, a DNA primer having a nucleotide sequence represented by SEQ ID NO: 5, a DNA primer having a nucleotide sequence represented by SEQ ID NO: 6, a DNA primer having a nucleotide
25 sequence represented by SEQ ID NO: 7, a DNA primer having a nucleotide sequence represented by SEQ ID NO: 14 and a DNA primer having a nucleotide sequence represented by SEQ ID NO: 18; with

30 ② at least one DNA primer selected from the group consisting of a DNA primer having a nucleotide sequence represented by SEQ ID NO: 2, a DNA primer having a nucleotide sequence represented by SEQ ID NO: 4, a DNA primer having a nucleotide sequence represented by SEQ ID NO: 8, a DNA primer having a nucleotide sequence represented by SEQ ID NO: 9, a DNA primer having a
35 nucleotide sequence represented by SEQ ID NO: 15, a DNA primer having a nucleotide sequence represented by SEQ IS NO: 17 and a DNA primer having a nucleotide sequence represented by SEQ ID

NO: 19;

is capable of amplifying the areas covering from the third membrane-spanning domain to the sixth membrane-spanning domain thereof at once like the conventional DNA primers and,

5 moreover, it is capable of more selectively and efficiently amplifying DNA coding for G protein coupled receptor proteins though it has not been obtained through the conventional DNA primers.

Moreover, among the DNA primers of the present
10 invention, the combination of

① at least one DNA primer selected from DNA primers having a nucleotide sequence of SEQ ID NO: 1 and DNA primers having a nucleotide sequence of SEQ ID NO: 12; with

② a DNA primer having a nucleotide sequence represented by
15 SEQ ID NO: 13;

is capable of amplifying the areas covering from the first membrane-spanning domain to the third membrane-spanning domain thereof at once.

Then (a) the amplified DNA (or fragment(s) thereof)
20 coding for the amino acid sequence of from the first membrane-spanning domain to the sixth membrane-spanning domain of G protein coupled receptor protein, (b) the amplified DNA (or fragment(s) thereof) coding for the amino acid sequence of from the first membrane-spanning domain to the seventh
25 membrane-spanning domain of G protein coupled receptor protein, (c) the amplified DNA (or fragment(s) thereof) coding for the amino acid sequence of from the third membrane-spanning domain to the sixth membrane-spanning domain of G protein coupled receptor protein, (d) the amplified DNA (or fragment(s)
30 thereof) coding for the amino acid sequence of from the third membrane-spanning domain to the seventh membrane-spanning domain of G protein coupled receptor protein, (e) the amplified DNA (or fragment(s) thereof) coding for the amino acid sequence of from the second membrane-spanning domain to the sixth
35 membrane-spanning domain of G protein coupled receptor protein, (f) the amplified DNA (or fragment(s) thereof) coding for the amino acid sequence of from the second membrane-spanning domain

to the seventh membrane-spanning domain of G protein coupled receptor protein, (g) the amplified DNA (or fragment(s) thereof) coding for the amino acid sequence of from the first membrane-spanning domain to the third membrane-spanning domain of G protein coupled receptor protein or (h) the amplified DNA (or fragment(s) thereof) coding for the amino acid sequence of other domains of G protein coupled receptor protein may be used as a probe(s) to screen for full-length DNA which completely encodes G protein coupled receptor proteins from DNA libraries according to methods known per se by those of skill in the art or methods similar thereto.

The DNA libraries used in the present invention include any of genome DNA libraries, cDNA libraries and RNA libraries. The term "DNA library" or "DNA libraries" as used herein refers to a DNA library or DNA libraries including all of those libraries.

The present invention further provides screening methods for target DNA (or fragment(s) thereof) coding for G protein coupled receptor protein from the DNA library containing DNA (or fragment(s) thereof) coding for receptor proteins, which comprise employing the DNA of the present invention as a DNA primer for the PCR.

One preferred embodiment of the present invention is a method for cloning full-length DNA which completely encodes an amino acid sequence of G protein coupled receptor protein from DNA libraries which comprises the steps of

- (i) using the DNA of the present invention as a DNA primer for PCR;
- (ii) carrying out PCR in the presence of a mixture of said DNA primer with the DNA library to amplify and select (i.e. screen for) a DNA fragment coding for the amino acid sequence of from the first membrane-spanning domain to the sixth membrane-spanning domain of G protein coupled receptor protein, a DNA fragment coding for the amino acid sequence of from the first membrane-spanning domain to the seventh membrane-spanning domain of G protein coupled receptor protein, a DNA fragment coding for the amino acid sequence of from the third membrane-

spanning domain to the sixth membrane-spanning domain of G protein coupled receptor protein, a DNA fragment coding for the amino acid sequence of from the third membrane-spanning domain to the seventh membrane-spanning domain of G protein coupled receptor protein, a DNA fragment coding for the amino acid sequence of from the second membrane-spanning domain to the sixth membrane-spanning domain of G protein coupled receptor protein, a DNA fragment coding for the amino acid sequence of from the second membrane-spanning domain to the seventh membrane-spanning domain of G protein coupled receptor protein, a DNA fragment coding for the amino acid sequence of from the first membrane-spanning domain to the third membrane-spanning domain of G protein coupled receptor protein or a DNA fragment coding for other domains of G protein coupled receptor protein; and

(iii) cloning said full-length DNA from the DNA library according to cloning methods known per se by those of skill in the art or methods similar thereto by using, as a probe, the DNA fragment obtained in the above step (ii).

Preferably, an embodiment of the present invention is a screening method of DNA coding for G protein coupled receptor proteins from DNA libraries, which comprises carrying out a polymerase chain reaction in the presence of a mixture of

- ① the DNA library,
- ② at least one DNA primer selected from the group consisting of DNA primers having a nucleotide sequence represented by SEQ ID NO: 1, DNA primers having a nucleotide sequence represented by SEQ ID NO: 3, DNA primers having a nucleotide sequence represented by SEQ ID NO: 5, DNA primers having a nucleotide sequence represented by SEQ ID NO: 6, DNA primers having a nucleotide sequence represented by SEQ ID NO: 7, DNA primers having a nucleotide sequence represented by SEQ ID NO: 10, DNA primers having a nucleotide sequence represented by SEQ ID NO: 14, DNA primers having a nucleotide sequence represented by SEQ ID NO: 16 and DNA primers having a nucleotide sequence represented by SEQ ID NO: 18 and

③ at least one DNA primer selected from the group consisting of DNA primers having a nucleotide sequence represented by SEQ ID NO: 2, DNA primers having a nucleotide sequence represented by SEQ ID NO: 4, DNA primers having a nucleotide sequence represented by SEQ ID NO: 8, DNA primers having a nucleotide sequence represented by SEQ ID NO: 9, DNA primers having a nucleotide sequence represented by SEQ ID NO: 11, DNA primers having a nucleotide sequence represented by SEQ ID NO: 15, DNA primers having a nucleotide sequence represented by SEQ ID NO: 17 and DNA primers having a nucleotide sequence represented by SEQ ID NO: 19 to selectively amplify template DNA coding for G protein coupled receptor protein contained in the DNA library.

More preferably, embodiments of the present invention include:

(1) a screening method of DNA coding for the amino acid sequence of G protein coupled receptor protein and the like (e.g. the regions spanning from the first transmembrane domain to the sixth transmembrane domain of G protein coupled receptor protein or other domains thereof) from a DNA library, which comprises carrying out a polymerase chain reaction in the presence of a mixture of

① the DNA library,

② at least one DNA primer selected from the group consisting of DNA primers having a nucleotide sequence represented by SEQ ID NO: 1 and DNA primers having a nucleotide sequence represented by SEQ ID NO: 12 and

③ at least one DNA primer selected from the group consisting of DNA primers having a nucleotide sequence represented by SEQ ID NO: 2, DNA primers having a nucleotide sequence represented by SEQ ID NO: 4, DNA primers having a nucleotide sequence represented by SEQ ID NO: 8, DNA primers having a nucleotide sequence represented by SEQ ID NO: 9, DNA primers having a nucleotide sequence represented by SEQ ID NO: 15, DNA primers having a nucleotide sequence represented by SEQ ID NO: 17 and DNA primers having a nucleotide sequence represented by SEQ ID NO: 19

to selectively amplify the DNA coding for the amino acid sequence of G protein coupled receptor protein and the like (e.g. the regions spanning from the first transmembrane domain to the sixth transmembrane domain of G protein coupled receptor protein or other domains thereof) contained in the DNA library;

(2) a screening method of DNA coding for the amino acid sequence of G protein coupled receptor protein and the like (e.g. the regions spanning from the first transmembrane domain to the seventh transmembrane domain of G protein coupled receptor protein or other domains thereof) from a DNA library, which comprises carrying out a polymerase chain reaction in the presence of a mixture of

- ① the DNA library,
- ② at least one DNA primer selected from the group consisting of DNA primers having a nucleotide sequence represented by SEQ ID NO: 1 and DNA primers having a nucleotide sequence represented by SEQ ID NO: 12 and
- ③ at least one DNA primer selected from the group consisting of DNA primers having a nucleotide sequence represented by SEQ ID NO: 11

to selectively amplify the DNA coding for the amino acid sequence of G protein coupled receptor protein and the like (e.g. the regions spanning from the first transmembrane domain to the seventh transmembrane domain of G protein coupled receptor protein or other domains thereof) contained in the DNA library;

(3) a screening method of DNA coding for the amino acid sequence of G protein coupled receptor protein and the like (e.g. the regions spanning from the second transmembrane domain to the sixth transmembrane domain of G protein coupled receptor protein or other domains thereof) from a DNA library, which comprises carrying out a polymerase chain reaction in the presence of a mixture of

- ① the DNA library,
- ② at least one DNA primer selected from the group consisting of DNA primers having a nucleotide sequence represented by

SEQ ID NO: 10 and DNA primers having a nucleotide sequence represented by SEQ ID NO: 16 and

③ at least one DNA primer selected from the group consisting of DNA primers having a nucleotide sequence represented by

5 SEQ ID NO: 2, DNA primers having a nucleotide sequence represented by SEQ ID NO: 4, DNA primers having a nucleotide sequence represented by SEQ ID NO: 8, DNA primers having a nucleotide sequence represented by SEQ ID NO: 9, DNA primers having a nucleotide sequence represented by SEQ ID NO: 15,
10 DNA primers having a nucleotide sequence represented by SEQ ID NO: 17 and DNA primers having a nucleotide sequence represented by SEQ ID NO: 19

to selectively amplify the DNA coding for the amino acid sequence of G protein coupled receptor protein and the like
15 (e.g. the regions spanning from the second transmembrane domain to the sixth transmembrane domain of G protein coupled receptor protein or other domains thereof) contained in the DNA library;

(4) a screening method of DNA coding for the amino acid
20 sequence of G protein coupled receptor protein and the like (e.g. the regions spanning from the second transmembrane domain to the seventh transmembrane domain of G protein coupled receptor protein or other domains thereof) from a DNA library, which comprises carrying out a polymerase chain reaction in the
25 presence of a mixture of

① the DNA library,

② at least one DNA primer selected from the group consisting of DNA primers having a nucleotide sequence represented by
30 SEQ ID NO: 10 and DNA primers having a nucleotide sequence represented by SEQ ID NO: 16 and

③ at least one DNA primer selected from the group consisting of DNA primers having a nucleotide sequence represented by
SEQ ID NO: 11

to selectively amplify the DNA coding for the amino acid
35 sequence of G protein coupled receptor protein and the like (e.g. the regions spanning from the second transmembrane domain to the seventh transmembrane domain of G protein coupled

receptor protein or other domains thereof) contained in the DNA library;

- (5) a screening method of DNA coding for the amino acid sequence of G protein coupled receptor protein and the like (e.g. the regions spanning from the third transmembrane domain to the sixth transmembrane domain of G protein coupled receptor protein or other domains thereof) from a DNA library, which comprises carrying out a polymerase chain reaction in the presence of a mixture of
- 10 ① the DNA library,
- ② at least one DNA primer selected from the group consisting of DNA primers having a nucleotide sequence represented by SEQ ID NO: 3, DNA primers having a nucleotide sequence represented by SEQ ID NO: 5, DNA primers having a nucleotide sequence represented by SEQ ID NO: 6, DNA primers having a nucleotide sequence represented by SEQ ID NO: 7, DNA primers having a nucleotide sequence represented by SEQ ID NO: 14 and DNA primers having a nucleotide sequence represented by SEQ ID NO: 18 and
- 20 ③ at least one DNA primer selected from the group consisting of DNA primers having a nucleotide sequence represented by SEQ ID NO: 2, DNA primers having a nucleotide sequence represented by SEQ ID NO: 4, DNA primers having a nucleotide sequence represented by SEQ ID NO: 8, DNA primers having a nucleotide sequence represented by SEQ ID NO: 9, DNA primers having a nucleotide sequence represented by SEQ ID NO: 15, DNA primers having a nucleotide sequence represented by SEQ ID NO: 17 and DNA primers having a nucleotide sequence represented by SEQ ID NO: 19
- 25 to selectively amplify the DNA coding for the amino acid sequence of G protein coupled receptor protein and the like (e.g. the regions spanning from the third transmembrane domain to the sixth transmembrane domain of G protein coupled receptor protein or other domains thereof) contained in
- 30 the DNA library;
- 35 (6) a screening method of DNA coding for the amino acid sequence of G protein coupled receptor protein and the like

(e.g. the regions spanning from the third transmembrane domain to the seventh transmembrane domain of G protein coupled receptor protein or other domains thereof) from a DNA library, which comprises carrying out a polymerase chain reaction in the presence of a mixture of

① the DNA library,

② at least one DNA primer selected from the group consisting of DNA primers having a nucleotide sequence represented by SEQ ID NO: 3, DNA primers having a nucleotide sequence

represented by SEQ ID NO: 5, DNA primers having a nucleotide sequence represented by SEQ ID NO: 6, DNA primers having a nucleotide sequence represented by SEQ ID NO: 7, DNA primers having a nucleotide sequence represented by SEQ ID NO: 14 and DNA primers having a nucleotide sequence represented by

SEQ ID NO: 18 and

③ at least one DNA primer selected from the group consisting of DNA primers having a nucleotide sequence represented by SEQ ID NO: 11

to selectively amplify the DNA coding for the amino acid sequence of G protein coupled receptor protein and the like (e.g. the regions spanning from the third transmembrane domain to the seventh transmembrane domain of G protein coupled receptor protein or other domains thereof) contained in the DNA library; and

(7) a screening method of DNA coding for the amino acid sequence of G protein coupled receptor protein and the like (e.g. the regions spanning from the first transmembrane domain to the third transmembrane domain of G protein coupled receptor protein or other domains thereof) from a DNA library, which comprises carrying out a polymerase chain reaction in the presence of a mixture of

① the DNA library,

② at least one DNA primer selected from the group consisting of DNA primers having a nucleotide sequence represented by SEQ ID NO: 1 and DNA primers having a nucleotide sequence represented by SEQ ID NO: 12 and

③ at least one DNA primer selected from the group consisting

of DNA primers having a nucleotide sequence represented by
SEQ ID NO: 13

to selectively amplify the DNA coding for the amino acid
sequence of G protein coupled receptor protein and the like
5 (e.g. the regions spanning from the first transmembrane
domain to the third transmembrane domain of G protein coupled
receptor protein or other domains thereof) contained in
the DNA library.

Particularly preferably, embodiments of the present
10 invention include:

(8) a screening method of DNA coding for the amino acid
sequence of G protein coupled receptor protein from a DNA
library, which comprises carrying out a polymerase chain
reaction in the presence of a mixture of

- 15 ① the DNA library,
② a DNA primer having a nucleotide sequence represented by
SEQ ID NO: 1 and
③ a DNA primer having a nucleotide sequence represented by
SEQ ID NO: 2

20 to selectively amplify the DNA coding for the amino acid
sequence of G protein coupled receptor protein contained in
the DNA library;

(9) a screening method of DNA coding for the amino acid
sequence of G protein coupled receptor protein from a DNA
25 library, which comprises carrying out a polymerase chain
reaction in the presence of a mixture of

- ① the DNA library,
② a DNA primer having a nucleotide sequence represented by
SEQ ID NO: 3 and
30 ③ a DNA primer having a nucleotide sequence represented by
SEQ ID NO: 4

to selectively amplify the DNA coding for the amino acid
sequence of G protein coupled receptor protein contained in
the DNA library;

35 (10) a screening method of DNA coding for the amino acid
sequence of G protein coupled receptor protein from a DNA
library, which comprises carrying out a polymerase chain

- 117 -

segments) thereof.

Preferably, the RNA or DNA (or fragment(s) thereof) obtained via the instant screening method for G protein coupled receptor protein encoding DNA wherein said method uses the DNA according to the present invention is a G protein coupled receptor protein-encoding RNA or DNA (or fragment(s) thereof) contained in the used DNA library. More specifically, it is an RNA or DNA (or RNA fragment(s) or DNA fragment(s) for G protein coupled receptor proteins such as angiotensin receptor, bombesin receptor, glutamine receptor, serotonin receptor, bombesin receptor, canavanine receptor, opioid receptor, melatonin receptor, vasopressin receptor, serotonin receptor, purine receptor (vasoactive intestinal and related peptide receptor), somatostatin receptor, dopamine receptor, motilin receptor, amylin receptor, bradykinin receptor, pancreastatin receptor (calcitonin receptor, leukotriene receptor, α - and β -chemokine receptor, prostaglandin receptor, thromboxane receptor, adenosine receptor, adrenaline receptor, α - and β -chemokine receptor (receptors to IL-8, GRO α , GRO β , GRO γ , MCP-3, I-309, NAP-2, ENA-78, PF4, IP10, GCP-2, MCP-1, HC14, MCP-3, I-309, MIP1 α , MIP-1 β , RANTES, etc.), endothelin receptor, enterogastrin receptor, histamine receptor, neurotensin receptor, TRH receptor, pancreatic polypeptide receptor, galanin receptor, their family member receptors, etc.

When the DNA obtained by the screening method of the present invention is the DNA fragment which partially codes for a G protein coupled receptor protein, it is possible to isolate a G protein coupled receptor protein, it is possible to isolate a completely encoding said G protein coupled receptor protein from a suitable DNA library according to cloning techniques known per se by using said DNA fragment as a probe.

Means for cloning the DNA completely encoding G protein coupled receptor proteins may include a PCR amplification employing a synthetic DNA primer having the partial nucleotide sequence of the DNA fragment partially

coding for the G protein coupled receptor protein and a selection of the target DNA via a hybridization with DNA or synthetic DNA having part or all of the region of said DNA fragments. The hybridization may be conducted, for example,
5 by the methods described in Molecular Cloning, 2nd ed.; J. Sambrook et al., Cold Spring Harbor Lab. Press, 1989. When the commercially available library is used, it may be conducted according to the manners described in the protocols attached thereto.

10 The DNA completely encoding G protein coupled receptor protein (full-length G protein coupled receptor protein DNA) may be used, depending upon its object, either as it is or after digesting with a restriction enzyme or after ligating with a linker if desired. Said DNA may have
15 ATG at the 5'-terminal as the translation initiation codon and TAA, TGA or TAG at the 3' terminal as the translation termination codon. These translation initiation codons and translation termination codons may be added using a suitable synthetic DNA adaptor. In addition, it is possible to
20 determine said receptor protein-expressing tissues/cells by northern blottings using said DNA as a probe. It is also possible to express target receptor proteins by introducing DNA having the entire coding region of the receptor protein into animal cells after binding with a suitable promoter.

25 The G protein coupled receptor protein according to the present invention is a G protein coupled receptor protein encoded by the G protein coupled receptor protein-encoding DNA obtained by the screening method of the present invention. More specifically, the G protein coupled receptor
30 protein according to the present invention includes G protein coupled receptor proteins such as angiotensin receptor protein, bombesin receptor protein, canavaninoid receptor protein, cholecystokinin receptor protein, glutamine receptor protein, serotonin receptor protein, melatonin receptor protein,
35 neuropeptide Y receptor protein, opioid receptor protein, purine receptor protein, vasopressin receptor protein, oxytocin receptor protein, VIP receptor protein (vasoactive

intestinal and related peptide receptor protein), somatostatin receptor protein, dopamine receptor protein, motilin receptor protein, amylin receptor protein, bradykinin receptor protein, CGRP receptor protein (calcitonin gene related peptide receptor protein), adrenomedullin receptor protein, leukotriene receptor protein, pancreastatin receptor protein, prostaglandin receptor protein, thromboxane receptor protein, adenosine receptor protein, adrenaline receptor protein, α - and β -chemokine receptor protein (receptor protein responsive to IL-8, GRO α , GRO β , GRO γ , NAP-2, ENA-78, PF4, IP10, GCP-2, MCP-1, HC14, MCP-3, I-309, MIP1 α , MIP-1 β , RANTES, etc.), endothelin receptor protein, enterogastrin receptor protein, histamine receptor protein, neurotensin receptor protein, TRH receptor protein, pancreatic polypeptide receptor protein, galanin receptor protein, family members thereof, etc.

According to the present invention, novel G protein coupled receptors proteins, peptide segments or fragments derived from the G protein coupled receptor protein, modified derivatives or analogues thereof, and salts thereof may be recognized, cloned, produced, isolated or characterized.

These G protein coupled receptor proteins are those derived from all cells and tissues (e.g. pituitary gland, pancreas, brain, kidney, liver, gonad, thyroid gland, cholecyst, bone marrow, adrenal, skin, muscle, lung, digestive duct, blood vessel, heart, etc.) of warm-blooded animals (e.g. guinea pig, rat, mouse, swine, sheep, cattle, monkey, human beings, rabbit, cat, dog, horse, etc.), and any of proteins as long as they comprise an amino acid sequence selected from the group consisting of an amino acid sequence represented by SEQ ID NO: 24, an amino acid sequence represented by SEQ ID NO: 25, an amino acid sequence represented by SEQ ID NO: 26, an amino acid sequence represented by SEQ ID NO: 27, an amino acid sequence represented by SEQ ID NO: 28, an amino acid sequence represented by SEQ ID NO: 34, an amino acid sequence represented by SEQ ID NO: 35, an amino acid sequence represented by SEQ ID NO: 38, an amino acid sequence represented by SEQ ID NO: 39, an amino acid sequence

represented by SEQ ID NO: 56, and substantial equivalents to the amino acid sequence represented by SEQ ID NO: 24, SEQ ID NO: 25, SEQ ID NO: 26, SEQ ID NO: 27, SEQ ID NO: 28, SEQ ID NO: 34, SEQ ID NO: 35, SEQ ID NO: 38, SEQ ID NO: 39, and/or SEQ ID NO: 56.

In one embodiment of the present invention, G protein coupled receptor proteins are those derived from all cells and tissues (e.g. pituitary gland, pancreas, brain, kidney, liver, gonad, thyroid gland, cholecyst, bone marrow, adrenal, skin, muscle, lung, digestive duct, blood vessel, heart, etc.) of warm-blooded animals (e.g. guinea pig, rat, mouse, swine, sheep, cattle, monkey, human beings, cat, dog, horse, etc.), and any of proteins as long as they comprise an amino acid sequence selected from the group consisting of an amino acid sequence represented by SEQ ID NO: 24, an amino acid sequence represented by SEQ ID NO: 25, an amino acid sequence represented by SEQ ID NO: 26, an amino acid sequence represented by SEQ ID NO: 27, an amino acid sequence represented by SEQ ID NO: 28, and substantial equivalents to the amino acid sequence represented by SEQ ID NO: 24, SEQ ID NO: 25, SEQ ID NO: 26, SEQ ID NO: 27, or SEQ ID NO: 28. These G protein coupled receptor proteins may include proteins having an amino acid sequence selected from the group consisting of an amino acid sequence represented by SEQ ID NO: 24, an amino acid sequence represented by SEQ ID NO: 25, an amino acid sequence represented by SEQ ID NO: 26, an amino acid sequence represented by SEQ ID NO: 27 and an amino acid sequence represented by SEQ ID NO: 28, proteins wherein the amino acid sequence thereof is about 90% to 99.9% homologous to an amino acid sequence represented by SEQ ID NO: 24, an amino acid sequence represented by SEQ ID NO: 25, an amino acid sequence represented by SEQ ID NO: 26, an amino acid sequence represented by SEQ ID NO: 27 or an amino acid sequence represented by SEQ ID NO: 28 and the activity thereof is substantially equivalent to the protein having an amino acid sequence represented by SEQ ID NO: 24, an amino acid sequence represented by SEQ ID NO: 25, an amino acid sequence

reaction in the presence of a mixture of

① the DNA library,

② a DNA primer having a nucleotide sequence represented by
SEQ ID NO: 6 and

5 ③ a DNA primer having a nucleotide sequence represented by
SEQ ID NO: 8

to selectively amplify the DNA coding for the amino acid
sequence of G protein coupled receptor protein contained in
the DNA library; and

10 (11) a screening method of DNA coding for the amino acid
sequence of G protein coupled receptor protein from a DNA
library, which comprises carrying out a polymerase chain
reaction in the presence of a mixture of

① the DNA library,

15 ② a DNA primer having a nucleotide sequence represented by
SEQ ID NO: 10 and

③ a DNA primer having a nucleotide sequence represented by
SEQ ID NO: 11

to selectively amplify the DNA coding for the amino acid
20 sequence of G protein coupled receptor protein contained in
the DNA library.

The cloned DNAs can be analyzed, usually by
restriction enzyme analysis and/or sequencing.

25 Target RNA or DNA (or fragment(s) thereof) coding
for G protein coupled receptor protein in the amplification
and the screening by the PCR techniques wherein the DNA of
the present invention is employed may include RNA, DNA or
fragments thereof coding for known (or prior art) G protein
coupled receptor proteins and RNA, DNA or fragments thereof
30 coding for unknown (novel) G protein coupled receptor
proteins.

These target RNA or DNA (or fragment(s) thereof) may
include novel nucleotide sequences and even known nucleotide
sequences.

35 Examples of such nucleotide sequences are RNA or DNA
(or fragment(s)) coding for a G protein coupled receptor
protein, said RNA or DNA (or fragment(s)) being derived from

all cells and tissues (e.g. pituitary gland, brain, pancreas, lung, adrenal gland, etc.) of vertebrate animals (e.g. mice, rats, cats, dogs, swines, cattle, horses, monkeys, human beings, etc.), insects or other invertebrate animals (e.g. drosophilae, silkworms, Barathra brassicae, etc.), plants (e.g. rice plant, wheat, tomato, etc.) and cultured cell lines derived therefrom, etc.

Specific examples of the nucleotide sequences are RNA or DNA (or fragment(s)) coding for G protein coupled receptor proteins such as receptor proteins to angiotensin, bombesin, canavaninoid, cholecystokinin, glutamine, serotonin, melatonin, neuropeptide Y, opioid, purine, vasopressin, oxytocin, VIP (vasoactive intestinal and related peptide), somatostatin, dopamine, motilin, amylin, bradykinin, CGRP (calcitonin gene related peptide), adrenomedullin, leukotriene, pancreastatin, prostaglandin, thromboxane, adenosine, adrenaline, α - and β -chemokine (IL-8, GRO α , GRO β , GRO γ , NAP-2, ENA-78, PF4, IP10, GCP-2, MCP-1, HC14, MCP-3, I-309, MIP1 α , MIP-1 β , RANTES, etc.), endothelin, enterogastrin, histamine, neurotensin, TRH, pancreatic polypeptide, galanin, family members thereof, etc.

In the PCR amplification using the DNA of the present invention, the DNA (or DNA fragment) acting as a template may include any DNA so far as it is derived from the above-mentioned tissues and cells. More specifically, the template DNA (or DNA fragment) includes any of genome DNA, genome DNA libraries, cDNA derived from the tissues and cells and cDNA libraries derived from the tissues and cells. cDNA libraries derived from human tissues and cells are particularly suitable. Vectors to be used in the DNA library may include any of bacteriophages, plasmids, cosmids, phagimids, etc. It is also possible to directly amplify the template DNA (or DNA fragment) by reverse transcriptase polymerase chain reaction (RT-PCR) techniques using mRNA fractions prepared from the tissues and cells. The DNA which is to be a template may be either DNA completely coding for G protein coupled receptor proteins or DNA fragments (or

represented by SEQ ID NO: 26, an amino acid sequence
represented by SEQ ID NO: 27 or an amino acid sequence
represented by SEQ ID NO: 28 and the like. The substantially
equivalent activity may include ligand binding activity,
5 signal information transmitting, etc. The term "substantially
equivalent" or "substantial equivalent" means that the nature
of the ligand binding activity and the like is equivalent.
Therefore, it is allowable that even differences among grades
such as ligand binding affinity grades and ligand binding
10 activity grades and quantitative factors such as molecular
weights of receptor proteins are present.

In another embodiment of the present invention, G
protein coupled receptor proteins include human pituitary
gland-derived G protein coupled receptor proteins comprising
15 an amino acid sequence selected from the group consisting of
an amino acid sequence represented by SEQ ID NO: 24, and/or
an amino acid sequence represented by SEQ ID NO: 25, mouse
pancreas-derived G protein coupled receptor proteins comprising
an amino acid sequence represented by SEQ ID NO: 27, mouse
20 pancreas-derived G protein coupled receptor proteins comprising
an amino acid sequence represented by SEQ ID NO: 28, etc.
Examples of the human pituitary gland-derived G protein coupled
receptor protein comprising an amino acid sequence selected
from the group consisting of an amino acid sequence represented
25 by SEQ ID NO: 24, and an amino acid sequence represented by SEQ
ID NO: 25, are human pituitary gland-derived G protein coupled
receptor proteins comprising an amino acid sequence represented
by SEQ ID NO: 24, etc. These G protein coupled receptor
proteins may include proteins wherein one or more amino acid
30 residues (preferably from 2 to 30 amino acid residues, more
preferably from 2 to 10 amino acid residues) are deleted from
the amino acid sequence of SEQ ID NO: 24, SEQ ID NO: 25, SEQ
ID NO: 26, SEQ ID NO: 27 or SEQ ID NO: 28, proteins wherein one
or more amino acid residues (preferably from 2 to 30 amino acid
35 residues, more preferably from 2 to 10 amino acid residues) are
added to the amino acid sequence of SEQ ID NO: 24, SEQ ID NO:
25, SEQ ID NO: 26, SEQ ID NO: 27 or SEQ ID NO: 28, proteins

wherein one or more amino acid residues (preferably from 2 to 30 amino acid residues, more preferably from 2 to 10 amino acid residues) in the amino acid sequence of SEQ ID NO: 24, SEQ ID NO: 25, SEQ ID NO: 26, SEQ ID NO: 27 or SEQ ID NO: 28, are substituted with one or more amino acid residues, etc.

In yet another embodiment of the present invention, G protein coupled receptor proteins include those derived from all cells and tissues (e.g. amygdaloid nucleus, pituitary gland, pancreas, brain, kidney, liver, gonad, thyroid gland, cholecyst, bone marrow, lung, digestive duct, blood vessel, heart, thymus, spleen, leukocyte, etc.) of warm-blooded animals (e.g. guinea pig, rat, mouse, pig, sheep, cattle, monkey, human beings, etc.), and any of proteins as long as they comprise an amino acid sequence selected from the group consisting of an amino acid sequence represented by SEQ ID NO: 34 and/or an amino acid sequence represented by SEQ ID NO: 35. These G protein coupled receptor proteins may include proteins having an amino acid sequence selected from the group consisting of an amino acid sequence represented by SEQ ID NO: 34 or/and an amino acid sequence represented by SEQ ID NO: 35, proteins wherein the amino acid sequence thereof is about 90% to 99.9% homologous to an amino acid sequence represented by SEQ ID NO: 34 or/and an amino acid sequence represented by SEQ ID NO: 35 and the activity thereof is substantially equivalent to the protein having an amino acid sequence represented by SEQ ID NO: 34 and/or an amino acid sequence represented by SEQ ID NO: 35, and the like. The substantially equivalent activity may include ligand binding activity, signal information transmitting, etc. The term "substantially equivalent" or "substantial equivalent" means that the nature of the ligand binding activity and the like is equivalent. Therefore, it is allowable that even differences among grades such as ligand binding affinity grades and ligand binding activity grades and quantitative factors such as molecular weights of receptor proteins are present. Examples of the G protein coupled receptor protein are human amygdaloid nucleus-derived G protein coupled receptor proteins

having an amino acid sequence selected from the group consisting of an amino acid sequence represented by SEQ ID NO: 34 and/or an amino acid sequence represented by SEQ ID NO: 35, etc. These G protein coupled receptor proteins may include
5 proteins wherein one or more amino acid residues (preferably from 2 to 30 amino acid residues, more preferably from 2 to 10 amino acid residues) are deleted from the amino acid sequence of SEQ ID NO: 34 or SEQ ID NO: 35, proteins wherein one or more amino acid residues (preferably from 2 to 30 amino acid
10 residues, more preferably from 2 to 10 amino acid residues) are added to the amino acid sequence of SEQ ID NO: 34 or SEQ ID NO: 35, proteins wherein one or more amino acid residues (preferably from 2 to 30 amino acid residues, more preferably from 2 to 10 amino acid residues) in the amino acid sequence of
15 SEQ ID NO: 34 or SEQ ID NO: 35, are substituted with one or more amino acid residues, etc.

In still another embodiment of the present invention, these G protein coupled receptor proteins are those derived from all cells and tissues (e.g. amygdaloid nucleus, pituitary
20 body, pancreas, brain, kidney, liver, gonad, thyroid gland, cholecyst, bone marrow, lung, digestive duct, blood vessel, heart, thymus, leukocyte, etc.) of warm-blooded animals (e.g. guinea pig, rat, mouse, swine, sheep, cattle, monkey, human beings, etc.), and any of proteins as long as they
25 comprise an amino acid sequence represented by SEQ ID NO: 38, or substantial equivalents to the amino acid sequence represented by SEQ ID NO: 38, preferably an amino acid sequence represented by SEQ ID NO: 39, or substantial equivalents to the amino acid sequence represented by SEQ ID NO: 39.
30 These G protein coupled receptor proteins may include proteins having an amino acid sequence represented by SEQ ID NO: 38, proteins wherein the amino acid sequence thereof is about 90% to 99.9% homologous to an amino acid sequence represented by SEQ ID NO: 38 and the activity thereof is substantially
35 equivalent to the protein having an amino acid sequence represented by SEQ ID NO: 38 and the like.
These G protein coupled receptor proteins are preferably

proteins having an amino acid sequence represented by SEQ ID NO: 39, proteins wherein the amino acid sequence thereof is about 90% to 99.9% homologous to an amino acid sequence represented by SEQ ID NO: 39 and the activity thereof is substantially equivalent to the protein having an amino acid sequence represented by SEQ ID NO: 39, etc. The substantially equivalent activity may include ligand binding activity, signal information transmitting, etc. The term "substantially equivalent" or "substantial equivalent" means that the nature of the ligand binding activity and the like is equivalent. Therefore, it is allowable that even differences among grades such as ligand binding affinity grades and ligand binding activity grades and quantitative factors such as molecular sizes or weights of receptor proteins are present.

It is suggested by data that the mouse pancreatic β -cell strain, MIN6-derived receptor protein of the present invention (e.g., SEQ ID NO: 38 and SEQ ID NO: 39, or proteins encoded by pMAH2-17) is a novel purinoceptor subtype which is clearly distinct from prior art purinoceptors.

In another more specific embodiment of the present invention, G protein coupled receptor proteins include mouse pancreatic β -cell line, MIN6, derived G protein coupled receptor proteins comprising an amino acid sequence represented by SEQ ID NO: 38, mouse pancreatic β -cell line, MIN6, derived G protein coupled receptor proteins wherein one or more amino acid residues (preferably from 2 to 30 amino acid residues, more preferably from 2 to 10 amino acid residues) are deleted from the amino acid sequence of SEQ ID NO: 38, proteins wherein one or more amino acid residues (preferably from 2 to 30 amino acid residues, more preferably from 2 to 10 amino acid residues) are added to the amino acid sequence of SEQ ID NO: 38, proteins wherein one or more amino acid residues (preferably from 2 to 30 amino acid residues, more preferably from 2 to 10 amino acid residues) are substituted with other amino acid residues in the amino acid sequence of SEQ ID NO: 38, etc. Further preferably these G protein coupled receptor proteins include mouse pancreatic β -cell line, MIN6, derived G

protein coupled receptor proteins comprising an amino acid sequence represented by SEQ ID NO: 39, mouse pancreatic β -cell line, MIN6, derived G protein coupled receptor proteins wherein one or more amino acid residues (preferably from 2 to 30 amino acid residues, more preferably from 2 to 10 amino acid residues) are deleted from the amino acid sequence of SEQ ID NO: 39, proteins wherein one or more amino acid residues (preferably from 2 to 30 amino acid residues, more preferably from 2 to 10 amino acid residues) are added to the amino acid sequence of SEQ ID NO: 39, proteins wherein one or more amino acid residues (preferably from 2 to 30 amino acid residues, more preferably from 2 to 10 amino acid residues) in the amino acid sequence of SEQ ID NO: 39 are substituted with other amino acid residues, etc.

In still another embodiment of the present invention, these G protein coupled receptor proteins are those derived from all cells and tissues (e.g. placenta, gonad, amygdaloid nucleus, pituitary body, pancreas, brain, kidney, liver, thyroid gland, cholecyst, bone marrow, lung, digestive duct, blood vessel, heart, thymus, leukocyte, etc.) of human beings, and any of proteins as long as they comprise an amino acid sequence represented by SEQ ID NO: 56, or substantial equivalents to the amino acid sequence represented by SEQ ID NO: 56. These G protein coupled receptor proteins may include proteins having an amino acid sequence represented by SEQ ID NO: 56, proteins wherein the amino acid sequence thereof is about 90% to 99.9% homologous to an amino acid sequence represented by SEQ ID NO: 56 and the activity thereof is substantially equivalent to the protein having an amino acid sequence represented by SEQ ID NO: 56 and the like. The substantially equivalent activity may include ligand binding activity, signal information transmitting, etc. The term "substantially equivalent" or "substantial equivalent" means that the nature of the ligand binding activity and the like is equivalent. Therefore, it is allowable that even differences among grades such as ligand binding affinity grades and ligand binding activity grades and quantitative factors

such as molecular sizes or weights of receptor proteins are present.

In another more specific embodiment of the present invention, G protein coupled receptor proteins include G
5 protein coupled receptor proteins comprising an amino acid sequence represented by SEQ ID NO: 56, G protein coupled receptor proteins wherein one or more amino acid residues (preferably from 2 to 30 amino acid residues, more preferably from 2 to 10 amino acid residues) are deleted from the amino
10 acid sequence of SEQ ID NO: 56, proteins wherein one or more amino acid residues (preferably from 2 to 30 amino acid residues, more preferably from 2 to 10 amino acid residues) are added to the amino acid sequence of SEQ ID NO: 56, proteins wherein one or more amino acid residues
15 (preferably from 2 to 30 amino acid residues, more preferably from 2 to 10 amino acid residues) in the amino acid sequence of SEQ ID NO: 56, are substituted with other amino acid residues, etc.

A portion of the amino acid sequence may be modified
20 (e.g. addition, deletion, substitution with other amino acids, etc.) in the G protein coupled receptor proteins of the present invention.

Furthermore, the G protein coupled receptor proteins of the present invention includes those wherein N-terminal Met
25 is protected with a protecting group (e.g., C₁₋₆ acyl group such as formyl, acetyl, etc.), those wherein the N-terminal side of Glu is cleaved in vivo to make said Glu pyroglutaminated, those wherein the intramolecular side chain of amino acids is protected with a suitable protecting group
30 (e.g., C₁₋₆ acyl group such as formyl, acetyl, etc.), conjugated proteins such as so-called "glycoproteins" wherein saccharide chains are bonded, etc.

The salt of said G protein coupled receptor protein of the present invention includes preferably physiologically
35 acceptable acid addition salts. Examples of such salts are salts thereof with inorganic acids (e.g. hydrochloric acid, phosphoric acid, hydrobromic acid, sulfuric acid, etc.), salts

- 1 2 7 -

thereof with organic acids (e.g. acetic acid, formic acid, propionic acid, fumaric acid, maleic acid, succinic acid, tartaric acid, citric acid, benzenesulfonic acid, etc.), etc. acid, methanesulfonic acid, benzenesulfonic acid, etc.), etc.

5 The G protein coupled receptor protein or its salt or cells of warm-blooded animals by purifying methods which are known per se by those skilled in the art or methods similar thereto or may be manufactured by culturing the transformant protein coupled receptor protein encoding DNA. The protein or its salt of the present invention may be manufactured by the peptide synthesis as described herein below.

10 The G protein coupled receptor protein may be manufactured by partial peptide of said G protein coupled receptor protein fragment (the partial peptide of said G protein coupled receptor protein may include, for example, the site which is exposed outside cell membranes, among the G protein coupled receptor protein molecule. Examples of the fragment are peptides containing a region which is analyzed as an extracellular area (hydrophilic region or site) in a hydrophobic plotting analysis on the G protein coupled receptor protein represented by any of Figures 24, 25, 28, 31, 32, 36, 38, 41, 44, 47, 50, 53, 57, 58, 59, 64, 70, 74, and 78.

20 A peptide which partly contains a hydrophobic region or site may be used as well. Further, a peptide which separately contains each domain may be used too although the partial peptide (peptide fragment) which contains plural domains at the same time will be used as well.

25 The salt of said G protein coupled receptor protein fragment (partial peptide thereof) includes preferably physiologically acceptable acid addition salts. Examples of such salts are salts thereof with inorganic acids (e.g. hydrochloric acid, phosphoric acid, hydrobromic acid, sulfuric acid, etc.), salts thereof with organic acids (e.g. acetic acid, formic acid, propionic acid, fumaric acid, maleic acid, succinic acid, tartaric acid, citric acid, malic acid, oxalic acid, benzoic acid, methanesulfonic acid, benzenesulfonic

acid, etc.), etc.

The G protein coupled receptor protein fragment (the partial peptide of the G protein coupled receptor protein) may be manufactured by synthesizing methods for peptides which are known per se by those skilled in the art or methods similar thereto or by cleaving (digesting) G protein coupled receptor proteins by a suitable peptidase. Methods of synthesizing peptide may be any of a solid phase synthesis and a liquid phase synthesis. Thus, a partial peptide (peptide fragment) or amino acids which can construct the protein of the present invention is condensed with the residual part thereof and, when the product has a protective group, said protective group is detached whereupon a desired peptide can be manufactured. Examples of the known methods for condensation and for detachment of protective groups include the following ① to ⑤ :

- ① M. Bodanszky and M. A. Ondetti: Peptide Synthesis, Interscience Publishers, New York (1966).
- ② Schroeder and Luebke: The Peptide, Academic Press, New York, 1965.
- ③ Nobuo Izumiya et al.: Fundamentals and Experiments of the Peptide Synthesis, Maruzen KK, Japan (1975).
- ④ Haruaki Yajima and Shumpei Sakakibara: "Seikagaku Jikken Koza 1" (Experiments of Biochemistry, Part 1), "Tanpakusitu No Kagaku IV" (Chemistry of Protein, IV), p.205 (1977), Japan.
- ⑤ Haruaki Yajima (ed): Development of Pharmaceuticals (Second Series), Vol. 14, Peptide Synthesis, Hirokawa Shoten, Japan.

After the reaction, conventional purifying techniques such as salting-out, extraction with solvents, distillation, column chromatography, liquid chromatography, electrophoresis, recrystallization, etc. are optionally combined so that the protein of the present invention can be purified and isolated. When the protein obtained as such is a free compound, it may be

converted to a suitable salt by known methods while, when it is obtained as a salt, the salt may be converted to a free compound or other salt compounds by known methods.

Furthermore, the product may be manufactured by
5 culturing the transformant (transfectant) containing the DNA coding for said partial peptide.

The G protein coupled receptor protein-encoding DNA obtained by the above-mentioned screening method using the DNA of the present invention and the G protein coupled receptor
10 protein encoded by said DNA or the peptide fragment (partial peptide thereof) encoded by said DNA may, for example, be used for the determination of a ligand to said G protein coupled receptor protein or for the screening of a compound which inhibits the binding of said protein coupled receptor protein
15 with a ligand.

In that case, an expression system for the G protein coupled receptor protein-encoding DNA is at first constructed. Hosts for said DNA may be any of animal cells, insect cells, yeasts, Bacillus subtilis, Escherichia coli, etc.
20 Promoters used therefor may be anyone so far as it is suitable as a promoter for the host used for gene expression. Incidentally, the utilization of enhancers for expression is effective as well.

Then the expressing cells per se which constructed
25 to express the G protein coupled receptor protein or the cell membrane fractions prepared therefrom by methods known per se by those skilled in the art or methods similar thereto may be subjected to a variety of receptor binding experiments. Ligands used therefor may include any of compounds labeled by
30 a commercially available radioisotope, etc., culture supernatants and tissue extracts which are directly labeled by a chloramine T method or by a lactoperoxidase method. Separation of bonded or free ligands may be carried out by a direct washing when cells adhered to substrates are used,
35 while, in the case of floating cells or cell membrane fractions thereof, it may be carried out by means of centrifugal separation or filtration. Nonspecific binding with container,

etc. may be estimated by addition of unlabeled ligands which are about 100 times as much concentrated relatively to the poured labeled ligand.

The ligand which is obtained by such a receptor
5 binding experiment may be subjected to a discrimination of agonist versus antagonist.

To be more specific, a natural substance or compound which is presumed to be a ligand with the G protein coupled receptor protein-expressing cell is cultured and, after that,
10 the culture supernatant liquid is collected or the cell is extracted. A change in the components contained therein is measured by, for example, a commercially available measuring kit (e.g. kits for cAMP, diacylglycerol, cGMP, proteinkinase A, etc.). Alternatively, it is possible to measure
15 physiological responses such as liberation of Fura-2, [³H]arachidonic acid and [³H]inositol phosphate metabolites by methods known per se by those skilled in the art or methods similar thereto. The compound or natural substance which is obtained by such a screening is an agonist for said G
20 protein coupled receptor protein or an antagonist for said G protein coupled receptor protein and is presumed to act on the tissues and cells in which said receptor is distributed. Accordingly, it is possible to check the pharmaceutical response (pharmaceutical effect) more efficiently by referring
25 to the distribution disclosed (clarified) by a northern blotting or the like. Moreover, a development of compounds having a novel pharmaceutical response (pharmaceutical effect) in, for example, central nervous tissues, circulatory system, kidney, pancreas, etc. is expected. An efficient development
30 of pharmaceuticals can be proceeded by amplifying G protein coupled receptor protein-encoding DNA selectively from tissues.

The G protein coupled receptor protein-encoding DNA of the present invention may be any coding DNA as long as it contains a nucleotide sequence coding for a G protein
35 coupled receptor protein which contains an amino acid sequence substantially equivalent to the amino acid sequence having SEQ ID NO: 24 and/or which has an activity substantially

equivalent to the amino acid sequence having SEQ ID NO: 24,
a G protein coupled receptor protein which contains an amino
acid sequence substantially equivalent to the amino acid
sequence having SEQ ID NO: 25 and/or which has an activity
5 substantially equivalent to the amino acid sequence having SEQ
ID NO: 25, a G protein coupled receptor protein which contains
an amino acid sequence substantially equivalent to the amino
acid sequence having SEQ ID NO: 26 and/or which has an activity
substantially equivalent to the amino acid sequence having SEQ
10 ID NO: 26, a G protein coupled receptor protein which contains
an amino acid sequence substantially equivalent to the amino
acid sequence having SEQ ID NO: 27 and/or which has an activity
substantially equivalent to the amino acid sequence having SEQ
ID NO: 27, or a G protein coupled receptor protein which
15 contains an amino acid sequence substantially equivalent to the
amino acid sequence having SEQ ID NO: 28 and/or which has an
activity substantially equivalent to the amino acid sequence
having SEQ ID NO: 28.

Still the G protein coupled receptor protein-encoding
20 DNA of the present invention may be any coding DNA as long as
it contains a nucleotide sequence coding for a G protein
coupled receptor protein which contains an amino acid sequence
substantially equivalent to the amino acid sequence having
SEQ ID NO: 34 and/or which has an activity substantially
25 equivalent to the amino acid sequence having SEQ ID NO: 34,
or a G protein coupled receptor protein which contains an amino
acid sequence substantially equivalent to the amino acid
sequence having SEQ ID NO: 35 and/or which has an activity
substantially equivalent to the amino acid sequence having SEQ
30 ID NO: 35.

Yet the G protein coupled receptor protein-encoding
DNA of the present invention may be any coding DNA as long as
it contains a nucleotide sequence coding for a G protein
coupled receptor protein which contains an amino acid sequence
35 substantially equivalent to the amino acid sequence having
SEQ ID NO: 38 and/or which has an activity substantially
equivalent to the amino acid sequence having SEQ ID NO: 38, or

preferably a G protein coupled receptor protein which contains an amino acid sequence substantially equivalent to the amino acid sequence having SEQ ID NO: 39 and/or which has an activity substantially equivalent to the amino acid sequence having SEQ ID NO: 39.

Yet the G protein coupled receptor protein-encoding DNA of the present invention may be any coding DNA as long as it contains a nucleotide sequence coding for a G protein coupled receptor protein which contains an amino acid sequence substantially equivalent to the amino acid sequence having SEQ ID NO: 56 and/or which has an activity substantially equivalent to the amino acid sequence having SEQ ID NO: 56, or preferably a G protein coupled receptor protein which contains an amino acid sequence substantially equivalent to the amino acid sequence having SEQ ID NO: 56 and/or which has an activity substantially equivalent to the amino acid sequence having SEQ ID NO: 56.

The DNA of the present invention may be any one of a human genome DNA, a human genome DNA library, a human tissue and cell-derived cDNA, a human tissue and cell-derived cDNA library and a synthetic DNA. The vector used for the library may include bacteriophage, plasmid, cosmid, phagemid, etc. The DNA can be further amplified directly by the reverse transcriptase polymerase chain reaction (hereinafter briefly referred to as "RT-PCR") using mRNA fractions prepared from tissues and cells.

In an embodiment, the DNA coding for the human pituitary gland-derived G protein coupled receptor protein comprising the amino acid sequence of SEQ ID NO: 24 includes DNA having a nucleotide sequence represented by SEQ ID NO: 29, etc. The DNA coding for the human pituitary gland-derived G protein coupled receptor protein comprising the amino acid sequence of SEQ ID NO: 25 includes DNA having a nucleotide sequence represented by SEQ ID NO: 30, etc. The DNA coding for the human pituitary gland-derived G protein coupled receptor protein comprising the amino acid sequence of SEQ ID NO: 26 includes DNA having a nucleotide sequence represented by

SEQ ID NO: 31, etc. The DNA coding for the mouse pancreas-derived G protein coupled receptor protein comprising the amino acid sequence of SEQ ID NO: 27 includes DNA having a nucleotide sequence represented by SEQ ID NO: 32, etc. The DNA coding for the mouse pancreas-derived G protein coupled receptor protein comprising the amino acid sequence of SEQ ID NO: 28 includes DNA having a nucleotide sequence represented by SEQ ID NO: 33, etc.

In another embodiment, the DNA coding for the human amygdaloid nucleus-derived G protein coupled receptor protein comprising the amino acid sequence of SEQ ID NO: 34 includes DNA having a nucleotide sequence represented by SEQ ID NO: 36, etc. The DNA coding for the human amygdaloid nucleus-derived G protein coupled receptor protein comprising the amino acid sequence of SEQ ID NO: 35 includes DNA having a nucleotide sequence represented by SEQ ID NO: 37, etc. The DNA coding for the human amygdaloid nucleus-derived G protein coupled receptor protein comprising the amino acid sequence of SEQ ID NO: 34 or the amino acid sequence of SEQ ID NO: 35 includes DNA having a nucleotide sequence represented by SEQ ID NO: 36, DNA having a nucleotide sequence represented by SEQ ID NO: 37, etc. Still in another embodiment, the DNA coding for the mouse pancreatic β -cell line, MIN6-derived G protein coupled receptor protein comprising the amino acid sequence of SEQ ID NO: 38 includes DNA having a nucleotide sequence represented by SEQ ID NO: 40, etc. The DNA coding for the mouse pancreatic β -cell line, MIN6-derived G protein coupled receptor protein comprising the amino acid sequence of SEQ ID NO: 39 includes DNA having a nucleotide sequence represented by SEQ ID NO: 41, etc. Yet in another embodiment, the DNA coding for the human-derived G protein coupled receptor protein comprising the amino acid sequence of SEQ ID NO: 56 includes DNA having a nucleotide sequence represented by SEQ ID NO: 57, etc.

The DNA completely coding for the G protein coupled receptor protein of the present invention can be cloned by (1) carrying out the PCR amplification using a synthetic DNA primer having a partial nucleotide sequence (nucleotide

fragment) of the G protein coupled receptor protein; or
(2) effecting the selection of a DNA constructed in a
suitable vector, based on the hybridization with a labeled
DNA fragment having part or all of the region encoding a human
5 G protein coupled receptor protein or a labeled synthetic DNA
having part or all of the coding region thereof.

The hybridization is carried out according to methods as
disclosed in, for example, Molecular Cloning, 2nd Ed., J.
Sambrook et al., Cold Spring Harbor Lab. Press, 1989.

10 When a DNA library commercially available in the market is
used, the hybridization is carried out according to protocols
manuals attached thereto.

The cloned G protein coupled receptor protein-
encoding DNA of the present invention can be used as it is, or
15 can be used, as desired, after modifications including
digestion with a restriction enzyme or addition of a linker
or adapter, etc. depending upon objects. The DNA may have
an initiation codon, ATG, on the 5' terminal side and
a termination codon, TAA, TGA or TAG, on the 3' terminal side.
20 These initiation and termination codons can be ligated by
using a suitable synthetic DNA adapter.

An expression vector for G protein coupled receptor
proteins can be produced by, for example, (a) cutting out a
target DNA fragment from the G protein coupled receptor
25 protein-encoding DNA of the present invention and (b) ligating
the target DNA fragment with the downstream site of a promoter
in a suitable expression vector.

The vector may include plasmids derived from
Escherichia coli (e.g., pBR322, pBR325, pUC12, pUC13, etc.),
30 plasmids derived from Bacillus subtilis (e.g., PUB110, PTP5,
pC194, etc.), plasmids derived from yeasts (e.g., pSH19, pSH15,
etc.), bacteriophages such as λ -phage, and animal virus such
as retrovirus, vaccinia virus and baculovirus.

According to the present invention, any promoter can
35 be used as long as it is compatible with a host which is used
for expressing a gene. When the host for the transformation is
E. coli, the promoters are preferably trp promoters, lac

promoters, recA promoters, λ_{PL} promoters, lpp promoters, etc. When the host for the transformation is the Bacillus, the promoters are preferably SPO1 promoters, SPO2 promoters, penP promoters, etc. When the host is an yeast, the promoters are preferably PH05 promoters, PGK promoters, GAP promoters, ADH promoters, etc. When the host is an animal cell, the promoters include SV40-derived promoters, retrovirus promoters, metallothionein promoters, heat shock promoters, cytomegalovirus promoters, SR α promoters, etc. An enhancer can be effectively utilized for the expression.

As required, furthermore, a host-compatible signal sequence is added to the N-terminal side of the G protein coupled receptor protein. When the host is E. coli, the utilizable signal sequences may include alkaline phosphatase signal sequences, OmpA signal sequences, etc. When the host is the Bacillus, they may include α -amylase signal sequences, subtilisin signal sequences, etc. When the host is an yeast, they may include mating factor α signal sequences, invertase signal sequences, etc. When the host is an animal cell, they may include insulin signal sequences, α -interferon signal sequences, antibody molecule signal sequences, etc.

A transformant or transfectant is produced by using the vector thus constructed, which carries the G protein coupled receptor protein-encoding DNA of the present invention.

The host may be, for example, Escherichia microorganisms, Bacillus microorganisms, yeasts, insect cells, animal cells, etc. Examples of the Escherichia and Bacillus microorganisms include Escherichia coli K12-DH1 [Proc. Natl. Acad. Sci. USA, Vol. 60, 160 (1968)], JM103 [Nucleic Acids Research, Vol. 9, 309 (1981)], JA221 [Journal of Molecular Biology, Vol. 120, 517 (1978)], HB101 [Journal of Molecular Biology, Vol. 41, 459 (1969)], C600 [Genetics, Vol. 39, 440 (1954)], etc.

Examples of the Bacillus microorganism are, for example, Bacillus subtilis MI14 [Gene, Vol. 24, 255 (1983)], 207-21 [Journal of Biochemistry, Vol. 95, 87 (1984)], etc.

The yeast may be, for example, Saccharomyces cerevisiae AH22, AH22R, NA87-11A, DKD-5D, 20B-12, etc. The insect may include

a silkworm (Bombyx mori larva), [Maeda et al, Nature, Vol. 315, 592 (1985)] etc. The host animal cell may be, for example, monkey-derived cell line, COS-7, Vero, Chinese hamster ovary cell line (CHO cell), DHFR gene-deficient Chinese hamster cell line (dhfr⁻ CHO cell), mouse L cell, murine myeloma cell, human FL cell, etc.

Depending on the host cell used, transformation is done using standard techniques appropriate to such cells. Transformation of Escherichia microorganisms can be carried out in accordance with methods as disclosed in, for example, Proc. Natl. Acad. Sci. USA, Vol. 69, 2110 (1972), Gene, Vol. 17, 107 (1982), etc. Transformation of Bacillus microorganisms can be carried out in accordance with methods as disclosed in, for example, Molecular & General Genetics, Vol. 168, 111 (1979), etc. Transformation of the yeast can be carried out in accordance with methods as disclosed in, for example, Proc. Natl. Acad. Sci. USA, Vol. 75, 1929 (1978), etc. The insect cells can be transformed in accordance with methods as disclosed in, for example, Bio/Technology, 6, 47-55, 1988. The animal cells can be transformed by methods as disclosed in, for example, Virology, Vol. 52, 456, 1973, etc. The transformants or transfectants which are transformed with expression vectors containing a G protein coupled receptor protein-encoding DNA are produced according to the aforementioned techniques.

Cultivation of the transformant (transfectant) in which the host is Escherichia or Bacillus microorganism can be carried out suitably in a liquid culture medium. The culture medium may contains carbon sources, nitrogen sources, minerals, etc. necessary for growing the transformant. The carbon source may include glucose, dextrin, soluble starch, sucrose, etc. The nitrogen source may include organic or inorganic substances such as ammonium salts, nitrates, corn steep liquor, peptone, casein, meat extracts, bean-cakes, potato extracts, etc. Examples of the minerals may include calcium chloride, sodium dihydrogen phosphate, magnesium chloride, etc. It is further allowable to add yeasts, vitamins, growth-

promoting factors, etc. It is desired that the culture medium is pH from about 5 to about 8.

The Escherichia microorganism culture medium is preferably an M9 medium containing, for example, glucose and casamino acid (Miller, Journal of Experiments in Molecular Genetics), 431-433, Cold Spring Harbor Laboratory, New York, 1972. Depending on necessity, the medium may be supplemented with drugs such as 3 β -indolyl acrylic acid in order to improve efficiency of the promoter. In the case of the Escherichia host, the cultivation is carried out usually at about 15 to 43 °C for about 3 to 24 hours. As required, aeration and stirring may be applied. In the case of the Bacillus host, the cultivation is carried out usually at about 30 to 40 °C for about 6 to 24 hours. As required, aeration and stirring may be also applied. In the case of the transformant in which the host is an yeast, the culture medium used may include, for example, a Burkholder minimum medium [Bostian, K.L. et al., Proc. Natl. Acad. Sci. USA, Vol. 77, 4505 (1980)], an SD medium containing 0.5% casamino acid [Bitter, G.A. et al., Proc. Natl. Acad. Sci. USA, Vol. 81, 5330 (1984)], etc.

It is preferable that pH of the culture medium is adjusted to be from about 5 to about 8. The cultivation is carried out usually at about 20 to 35 °C for about 24 to 72 hours.

As required, aeration and stirring may be applied.

In the case of the transformant in which the host is an insect, the culture medium used may include those obtained by suitably adding additives such as passivated (or immobilized) 10% bovine serum and the like to the Grace's insect medium (Grace, T.C.C., Nature, 195, 788 (1962)). It is preferable that pH of the culture medium is adjusted to be about 6.2 to 6.4. The cultivation is usually carried out at about 27 °C for about 3 to 5 days. As desired, aeration and stirring may be applied.

In the case of the transformant in which the host is an animal cell, the culture medium used may include MEM medium [Science, Vol. 122, 501 (1952)], DMEM medium [Virology, Vol. 8, 396 (1959)], RPMI 1640 medium [Journal of the American Medical Association, Vol. 199, 519 (1967)], 199 medium [Proceedings of

the Society of the Biological Medicine, Vol. 73, 1 (1950)],
etc. which are containing, for example, about 5 to 20% of
fetal calf serum. It is preferable that the pH is from about
6 to about 8. The cultivation is usually carried out at about
5 30 to 40 °C for about 15 to 60 hours. As required, aeration
and stirring may be applied.

Separation and purification of the G protein coupled
receptor protein from the above-mentioned cultures can be
carried out according to methods described herein below.

10 To extract G protein coupled receptor proteins
from the cultured microorganisms or cells, the microorganisms
or cells are collected by known methods after the cultivation,
suspended in a suitable buffer solution, disrupted by
ultrasonic waves, lysozyme and/or freezing and thawing, etc.
15 and, then, a crude extract of the G protein coupled receptor
protein is obtained by centrifugation or filtration. Other
conventional extracting or isolating methods can be applied.
The buffer solution may contain a protein-denaturing agent
such as urea or guanidine hydrochloride or a surfactant such
20 as Triton X-100 (registered trademark, hereinafter often
referred to as "TM").

In case where G protein coupled receptor proteins
are secreted into culture media, supernatant liquids are
separated from the microorganisms or cells after the
25 cultivation is finished and the resulting supernatant liquid is
collected by widely known methods. The culture supernatant
liquid and extract containing G protein coupled receptor
proteins can be purified by suitable combinations of widely
known methods for separation, isolation and purification.
30 The widely known methods of separation, isolation and
purification may include methods which utilizes solubility,
such as salting out or sedimentation with solvents
methods which utilizes chiefly a difference in the molecular
size or weight, such as dialysis, ultrafiltration, gel
35 filtration and SDS-polyacrylamide gel electrophoresis, methods
utilizing a difference in the electric charge, such as
ion-exchange chromatography, methods utilizing specific

affinity such as affinity chromatography, methods utilizing a difference in the hydrophobic property, such as inverse-phase high-performance liquid chromatography, and methods utilizing a difference in the isoelectric point such as isoelectric electrophoresis, etc.

In case where the G protein coupled receptor protein thus obtained is in a free form, the free protein can be converted into a salt thereof by known methods or method analogous thereto. In case where the G protein coupled receptor protein thus obtained is in a salt form vice versa, the protein salt can be converted into a free form or into any other salt thereof by known methods or method analogous thereto.

The G protein coupled receptor protein produced by the transformant can be arbitrarily modified or a polypeptide can be partly removed therefrom, by the action of a suitable protein-modifying enzyme before or after the purification. The protein-modifying enzyme may include trypsin, chymotrypsin, arginyl endopeptidase, protein kinase, glycosidase, etc. The activity of the G protein coupled receptor protein thus formed can be measured by experimenting the coupling (or binding) with a ligand or by enzyme immunoassays (enzyme linked immunoassays) using specific antibodies.

The G protein coupled receptor protein-encoding DNA and the G protein coupled receptor protein of the present invention can be used for:

- ① methods of determining ligands for the G protein coupled receptor protein of the present invention,
- ② obtaining an antibody and an antiserum,
- ③ constructing a system for expressing a recombinant receptor protein,
- ④ developing a receptor-binding assay system using the above developing system and screening pharmaceutical candidate compounds,
- ⑤ designing drugs based upon the comparison with ligands and receptors which have a similar or analogous structure,

- ⑥ preparing a probe in the analysis of genes and preparing a PCR primer, and
- ⑦ gene manipulating therapy.

5 In particular, it is allowable to screen a G protein coupled receptor agonist or antagonist specific to a warm-blooded animal such as human being by a receptor-binding assay system which uses a system for expressing a recombinant G protein coupled receptor protein of the present invention. The agonist or antagonist thus screened or characterized
10 permits various applications including prevention and/or therapy of a variety of diseases.

Concretely described below are uses of G protein coupled receptor proteins, partial peptide thereof (peptide fragment thereof), G protein coupled receptor protein-encoding
15 DNAs and antibodies against the G protein coupled receptor protein according to the present invention.

As hereunder, more detailed description will be made on the usefulness of the G protein coupled receptor protein-encoding DNA obtained by the screening method for G protein
20 coupled receptor protein-encoding DNAs according to the present invention, the G protein coupled receptor proteins encoded by said DNA, peptide fragments or segments thereof (including partial peptides thereof) or salts thereof (hereinafter, those including their salts, will be referred to as the "G protein
25 coupled receptor protein or a peptide fragment thereof"), cells or cell membrane fractions thereof each containing the recombinant type G protein coupled receptor protein, etc. Their various applications are also disclosed herein below.

30 (1) Method for Determining Ligands to the G Protein Coupled Receptor Protein

The G protein coupled receptor protein (or the peptide segment thereof) is useful as a reagent for investigating or determining a ligand to said G protein coupled receptor protein.

35 According to the present invention, methods for determining a ligand to the G protein coupled receptor protein

which comprises contacting the G protein coupled receptor protein or the peptide segment or fragment thereof with the compound to be tested are provided.

The compound to be tested may include not only known
5 ligands such as angiotensins, bombesins, canavanoids, cholecystokinins, glutamine, serotonin, melatonins, neuropeptides Y, opioids, purine, vasopressins, oxytocins, VIP (vasoactive intestinal and related peptides),
10 somatostatins, dopamine, motilins, amylin, bradykinins, CGRP (calcitonin gene related peptides), adrenomedullins, leukotrienes, pancreastatins, prostaglandins, thromboxanes, adenosine, adrenaline, α - and β -chemokines (IL-8, GRO α , GRO β , GRO γ , NAP-2, ENA-78, PF4, IP10, GCP-2, MCP-1, HC14, MCP-3, I-309, MIP1 α , MIP-1 β , RANTES, etc.), endothelins,
15 enterogastrins, histamine, neurotensins, TRH, pancreatic polypeptides, galanin, modified derivatives thereof, analogues thereof, family members thereof and the like but also tissue extracts, cell culture supernatants, etc. of warm-blooded animals (such as mice, rats, swines, cattle, sheep, monkeys
20 and human being), etc. For example, said tissue extract, said cell culture supernatant, etc. is added to the G protein coupled receptor protein for measurement of the cell stimulating activity, etc. and fractionated by relying on the measurements whereupon a single ligand can be finally
25 obtained.

In one specific embodiment of the present invention, said method for determining the ligand includes a method for determining a compound or a salt thereof capable of
stimulating a target cell which comprises binding said compound
30 with the G protein coupled receptor protein either in the presence of the G protein coupled receptor protein or the peptide segment thereof or in a receptor binding assay system in which the expression system for the recombinant type receptor protein is constructed and used; and measuring the
35 receptor-mediated cell stimulating activity, etc. Examples of said cell stimulating activities include promoting activity or inhibiting activity on biological responses,

e.g. liberation of arachidonic acid, liberation of acetylcholine, liberation of endocellular Ca^{2+} , production of endocellular cAMP, production of endocellular cGMP, production of inositol phosphate, changes in the cell membrane potential, phosphorylation of endocellular protein, activation of c-fos, lowering in pH, activation of G protein, cell promulgation, etc. Examples of said compound or salt capable of stimulating the cell via binding with the G protein coupled receptor protein include peptides, proteins, nonpeptidic compounds, synthetic compounds, fermented products, etc.

In said method for determining the ligand, the characteristic feature is that when the G protein coupled receptor protein or the peptide segment thereof is contacted with the test compound, for example, the binding amount, the cell stimulating activity, etc. of the test compound to the G protein coupled receptor protein or the peptide segment thereof is measured.

In more specific embodiments of the present invention, said methods for determining the ligand includes:

- ① a method of determining a ligand to a G protein coupled receptor protein, which comprises contacting a labeled test compound with a G protein coupled receptor protein or a peptide segment thereof, and measuring the amount of the labeled test compound binding with said protein or salt thereof or with said peptide fragment or salt thereof;
- ② a method of determining a ligand to a G protein coupled receptor protein, which comprises contacting a labeled test compound with cells containing the G protein coupled receptor protein or the membrane fraction of said cell, and measuring the amount of the labeled test compound binding with said cells or said cell fraction;
- ③ a method of determining a ligand to a G protein coupled receptor protein, which comprises contacting a labeled test compound with the G protein coupled receptor protein expressed on cell membranes by culturing transformants containing the DNA coding for the G protein coupled receptor protein, and measuring the amount of the labeled test compound binding with

said G protein coupled receptor protein;

④ a method of determining a ligand to a G protein coupled receptor protein, which comprises contacting a test compound with cells containing the G protein coupled receptor protein, and measuring the cell stimulating activity (e.g. promoting or inhibiting activity on biological responses such as liberation of arachidonic acid, liberation of acetylcholine, liberation of endocellular Ca^{2+} , production of endocellular cAMP, production of endocellular cGMP, production of inositol phosphate, changes in the cell membrane potential, phosphorylation of endocellular protein, activation of c-fos, lowering in pH, activation of G protein, cell promulgation, etc.) via the G protein coupled receptor protein; and

⑤ a method of determining a ligand to the G protein coupled receptor protein, which comprises contacting a test compound with the G protein coupled receptor protein expressed on the cell membrane by culturing transformants containing the DNA coding for the G protein coupled receptor protein, and measuring the cell stimulating activity (activity for promoting or inhibiting physiological responses such as liberation of arachidonic acid, liberation of acetylcholine, liberation of endocellular Ca^{2+} , production of endocellular cAMP, production of endocellular cGMP, production of inositol phosphate, changes in the cell membrane potential, phosphorylation of endocellular protein, activation of c-fos, lowering in pH, activation of G protein, cell promulgation, etc.) via the G protein coupled receptor protein.

Described below are specific explanations on the determining method of ligands according to the present invention which are provided only for illustrative purposes.

First, the G protein coupled receptor protein used for the method for determining the ligand may include any material so far as it contains a G protein coupled receptor protein or a peptide fragment or segment thereof (including a partial peptide thereof) or a salt thereof although it is preferable to express a large amount of G protein coupled receptor proteins in animal cells.

In the manufacture of the G protein coupled receptor protein, the above-mentioned method can be used and it may be carried out by expressing said protein encoding DNA in mammalian cells or in insect cells. With respect to the DNA
5 fragment coding for the aimed region, complementary DNA may be used although it is not limited thereto. For example, gene fragments or synthetic DNA may be used as well.

In order to introduce the G protein coupled receptor protein-encoding DNA fragment into host animal cells and to
10 express it efficiently, it is preferred that said DNA fragment is incorporated into the downstream site of polyhedron promoters derived from nuclear polyhedrosis virus belonging to baculovirus, promoters derived from SV40, promoters derived from retrovirus, metallothionein promoters,
15 human heat shock promoters, cytomegalovirus promoters, SR α promoters, etc. Examinations of the quantity and the quality of the expressed receptor can be carried out by methods per se known to those of skill in the art or methods similar thereto. For example, they may be conducted by methods
20 described in publications such as Nambi, P. et al: The Journal of Biochemical Society, vol.267, pages 19555-19559 (1992).

Accordingly, with respect to the determination of the ligand, the material containing a G protein coupled receptor protein or peptide segment thereof may include
25 products containing G protein coupled receptor proteins which are purified by methods per se known to those of skill in the art or methods similar thereto, peptide fragments of said G protein coupled receptor protein, cells containing said G protein coupled receptor protein, membrane fractions of the
30 cell containing said protein, etc.

When the G protein coupled receptor protein-containing cell is used in the determining method of the ligand, said cell may be immobilized with binding agents including glutaraldehyde, formalin, etc. The immobilization
35 may be carried out by methods per se known to those of skill in the art or methods similar thereto.

The G protein coupled receptor protein-

containing cells are host cells expressing the G protein coupled receptor protein. Examples of said host cells are microorganisms such as Escherichia coli, Bacillus subtilis, yeasts, insect cells, animal cells, etc.

5 The cell membrane fraction is a cell membrane-rich fraction which is prepared by methods per se known to those of skill in the art or methods similar thereto after disruption of cells. Examples of cell disruption may include a method for squeezing cells using a Potter-Elvehjem homogenizer,
10 a disruption by a Waring blender or a Polytron (manufactured by Kinematica), a disruption by ultrasonic waves, a disruption via blowing out cells from small nozzles together with applying a pressure using a French press or the like, etc. In the fractionation of the cell membrane, a fractionation method by
15 means of centrifugal force such as a fractional centrifugal separation and a density gradient centrifugal separation is mainly used. For example, disrupted cellular liquid is centrifuged at a low speed (500 rpm to 3,000 rpm) for a short period (usually, from about one to ten minutes), the
20 supernatant liquid is further centrifuged at a high speed (1,500 rpm to 3,000 rpm) usually for 30 minutes to two hours and the resulting precipitate is used as a membrane fraction. Said membrane fraction contains a lot of the expressed G protein coupled receptor protein and a lot of membrane
25 components such as phospholipids and membrane proteins derived from the cells.

 The amount of the G protein coupled receptor protein in the membrane fraction cell containing said G protein coupled receptor protein is preferably 10^3 - 10^8 molecules
30 per cell or, suitably, 10^5 to 10^7 molecules per cell. Incidentally, the more the expressed amount, the higher the ligand binding activity (specific activity) per membrane fraction whereby the construction of a highly sensitive screening system becomes possible and, moreover, it may enable
35 us to measure the large amount of samples within the same lot.

 In conducting the above-mentioned methods ① to ② wherein ligands capable of binding with the G protein coupled

receptor protein are determined, a suitable G protein coupled receptor fraction and a labeled test compound are necessary. The G protein coupled receptor fraction is preferably a naturally occurring (natural type) G protein coupled receptor, a recombinant type G protein coupled receptor having the activity equivalent to that of the natural type. Here, the term "activity equivalent to" means the equivalent ligand binding activity, etc.

Suitable examples of the labeled test compound are angiotensin, bombesin, canavanine, cholecystokinin, glutamine, serotonin, melatonin, neuropeptide Y, opioid, purine, vasopressin, oxytocin, VIP (vasoactive intestinal and related peptides), somatostatin, dopamine, motilin, amylin, bradykinin, CGRP (calcitonin gene related peptides), adrenomedullin, leukotriene, pancreastatin, prostaglandin, thromboxane, adenosine, adrenaline, α - and β -chemokine (IL-8, GRO α , GRO β , GRO γ , NAP-2, ENA-78, PF4, IP10, GCP-2, MCP-1, HC14, MCP-3, I-309, MIP1 α , MIP-1 β , RANTES, etc.), endothelin, enterogastrin, histamine, neurotensin, TRH, pancreatic polypeptides, galanin, an analogue derivative thereof, etc. which are labeled with [3 H], [125 I], [14 C], [35 S], etc.

Specifically, the determination of ligands capable of binding with G protein coupled receptor proteins is carried out as follows:

First, cells or cell membrane fractions containing the G protein coupled receptor protein are suspended in a buffer suitable for the determining method to prepare the receptor sample in conducting the method of determining the ligand binding with the G protein coupled receptor protein. The buffer may include any buffer such as Tris-HCl buffer or phosphate buffer with pH 4-10 (preferably, pH 6-8), etc., as long as it does not inhibit the binding of the ligand with the receptor. In addition, surface-active agents such as CHAPS, Tween 80TM (Kao-Atlas, Japan), digitonin, deoxycholate, etc. and various proteins such as bovine serum albumin (BSA), gelatin, milk derivatives, etc. may be added to the buffer with an object of decreasing the non-specific binding.

Further, a protease inhibitor such as PMSF, leupeptin, E-64 (manufactured by Peptide Laboratory), pepstatin, etc. may be added with an object of inhibiting the decomposition of the receptor and the ligand by protease. A test compound labeled with a predetermined (or certain) amount (5,000 cpm to 500,000 cpm) of [^3H], [^{125}I], [^{14}C], [^{35}S], etc. is made copresent in 0.01 ml to 10 ml of said receptor solution.

In order to know the non-specific binding amount (NSB), a reaction tube to which a great excessive amount of the unlabeled test compound is added is prepared as well.

The reaction is carried out at 0-50°C (preferably at 4-37°C) for 20 minutes to 24 hours (preferably 30 minutes to three hours). After the reaction, it is filtered through a glass fiber filter or the like, washed with a suitable amount of the same buffer and the radioactivity remaining in the glass fiber filter is measured by means of a liquid scintillation counter or a gamma-counter. The test compound in which the count (B - NSB) obtained by subtracting the non-specific binding amount (NSB) from the total binding amount (B) is more than 0 cpm can be selected as a ligand to the G protein coupled receptor protein of the present invention.

In conducting the above-mentioned methods ④ to ⑤ wherein ligands capable of binding with the G protein coupled receptor protein are determined, the cell stimulating activity (e.g. the liberation of arachidonic acid, the liberation of acetylcholine, endocellular Ca^{2+} liberation, endocellular cAMP production, the production of inositol phosphate, changes in the cell membrane potential, the phosphorylation of endocellular protein, the activation of c-fos, lowering of pH, the activation of G protein, cell promulgation, etc.) mediated by the G protein coupled receptor protein may be measured by known methods or by the use of commercially available measuring kits. To be more specific, G protein coupled receptor protein-containing cells are at first cultured in a multi-well plate or the like.

In conducting the determination of ligand, it is substituted with a fresh medium or a suitable buffer which

does not show toxicity to the cells in advance of the experiment, and incubated for certain period after adding a test compound, etc. thereto. Then, the cells are extracted or the supernatant liquid is recovered and the resulting product is determined by each of the methods. When it is difficult to identify the production of the substance (e.g. arachdonic acid) which is to be an index for the cell stimulating activity due to the decomposing enzyme contained in the cell, an assay may be carried out by adding an inhibitor against said decomposing enzyme. With respect to the activity such as an inhibitory action against cAMP production, it may be detected as an inhibitory action against the production of the cells whose fundamental production is increased by forskolin or the like.

The kit used for the method of determining the ligand binding with the G protein coupled receptor protein includes a G protein coupled receptor protein or a peptide fragment thereof, cells containing the G protein coupled receptor protein, a membrane fraction from the cells containing the G protein coupled receptor protein, etc.

Examples of the kit for determining the ligand are as follows:

1. Reagent for Determining the Ligand.

① Buffer for Measurement and Buffer for Washing.

The buffering product wherein 0.05% of bovine serum albumin (manufactured by Sigma) is added to Hanks' Balanced Salt Solution (manufactured by Gibco).

This product may be sterilized by filtration through a membrane filter with a 0.45 μ m pore size, and stored at 4°C or may be formulated upon use.

② G Protein Coupled Receptor Protein Sample.

CHO cells in which G protein coupled receptor proteins are expressed are subcultured at the rate of 5×10^5 cells/well in a 12-well plate and cultured at 37°C in a humidified 5% CO₂/95% air atmosphere for two days to prepare the sample.

③ Labeled Test Compound.

The compound which is labeled with commercially

available [^3H], [^{125}I], [^{14}C], [^{35}S], etc. or labeled with a suitable method.

The product in a state of an aqueous solution is stored at 4°C or at -20°C and, upon use, diluted to 1 μM with a buffer for the measurement. In the case of the test compound which is hardly soluble in water, it is dissolved in dimethylformamide, DMSO, methanol, etc.

④ Unlabeled Test Compound.

The same compound for the labeled one is prepared in a concentration of 100 to 1,000-fold concentrated state.

2. Method of Measurement.

① G protein coupled receptor protein-expressing CHO cells cultured in a 12-well tissue culture plate are washed twice with 1 ml of buffer for the measurement and then 490 μl of buffer for the measurement is added to each well.

② Five μl of the labeled test compound is added and the mixture is made to react at room temperature for one hour. For measuring the nonspecific binding amount, 5 μl of the unlabeled test compound is added.

③ The reaction solution is removed from each well, which is washed with 1 ml of a buffer for the measurement three times. The labeled test compound which is binding with the cells is dissolved in 0.2N NaOH-1% SDS and mixed with 4 ml of a liquid scintillator A (manufactured by Wako Pure Chemical, Japan).

④ Radioactivity is measured using a liquid scintillation counter (manufactured by Beckmann).

The ligand which can bind with the G protein coupled receptor protein include substances occurring or existing, for example, in brain, pituitary gland, pancreas, etc. Examples of the ligand are angiotensin, bombesin, canavaninoid, cholecystokinin, glutamine, serotonin, melatonin, neuropeptide Y, opioid, purine, vasopressin, oxytocin, VIP (vasoactive intestinal and related peptide), somatostatin, dopamine, motilin, amylin, bradykinin, CGRP (calcitonin gene related

peptide), adrenomedullin, leukotriene, pancreastatin, prostaglandin, thromboxane, thromboxatin, adenosine, adrenaline, α - and β -chemokine (IL-8, GRO α , GRO β , GRO γ , NAP-2, ENA-78, PF4, IP10, GCP-2, MCP-1, HC14, MCP-3, I-309, 5 MIP1 α , MIP-1 β , RANTES, etc.), endothelin, enterogastrin, histamine, neurotensin, TRH, pancreatic polypeptide, galanin, modified derivatives thereof, analogues thereof, etc.

Since the receptor protein encoded by pMAH2-17 is highly homologous to prinoceptors, it is considered that there 10 are strong possibility of a subtype within prinoceptor families. All data including electrophysiological measurements are supporting that the mouse pancreatic β -cell strain, MIN6-derived receptor protein of the present invention (e.g., SEQ ID NO: 38 and SEQ ID NO: 39, or proteins encoded by 15 pMAH2-17) is a novel purinoceptor subtype. In other words, it is suggested that the ligand capable of binding with the mouse pancreatic β -cell strain, MIN6-derived receptor protein of the present invention (e.g., SEQ ID NO: 38 and SEQ ID NO: 39, or proteins encoded by pMAH2-17) is a purine 20 compound such as ATP. Further, the receptor protein (e.g., SEQ ID NO: 56, or proteins encoded by pMAH2-17) is considered to be a novel human type purinoceptor. It is presumed that it is advantageously useful in efficiently screening for agonists or antagonists to receptor proteins which control or regulate 25 functions in the central nervous system or immune system, related to purine compounds, and in developing pharmaceuticals.

(2) Preventive and Therapeutic Agent for of G Protein

Conjugated Receptor Protein Deficiency Diseases

If a ligand to the G protein coupled receptor protein 30 is disclosed via the aforementioned method (1), the G protein coupled receptor protein-encoding DNA can be used a preventive and/or therapeutic agent for treating said G protein coupled receptor protein deficiency diseases depending upon the action that said ligand exerts.

35 For example, when there is a patient for whom the physiological action of the ligand cannot be expected because

of a decrease in the G protein coupled receptor protein in vivo, the amount of the G protein coupled receptor protein in the brain cells of said patient can be increased whereby the action of the ligand can be fully achieved by:

- 5 (a) administering the G protein coupled receptor protein-encoding DNA to the patient to express it; or
- (b) inserting the G protein coupled receptor protein-encoding DNA into brain cells or the like to express it, followed by transplanting said brain cells or the like to said patient.

10 Accordingly, the G protein coupled receptor protein-encoding DNA can be used as a safe and less toxic preventive and therapeutic agent for the G protein coupled receptor protein deficiency diseases. In an embodiment, it is suggested that the ligands capable of binding with the mouse pancreatic

15 β -cell strain, MIN6-derived receptor protein of the present invention (e.g., SEQ ID NO: 38 and SEQ ID NO: 39, or proteins encoded by pMAH2-17) and further with the human-derived receptor protein of the present invention (e.g., SEQ ID NO: 56, or proteins encoded by pHAH2-17) are purine compounds such as

20 ATP. Therefore, the disease to be treated may include diseases or syndromes in connection with purine ligand compounds. Examples of such diseases may include cancer, immunodeficiency, autoimmune disease, rheumatoid arthritis, rejection on internal organ transplant, hypertension, diabetes, cystic fibrosis,

25 hypotension, incontinence of urine, pain, etc.

(3) Preventive and Therapeutic Pharmaceutical Composition for Human-Derived G Protein Conjugated Receptor Protein Deficiency Diseases

If the human-derived G protein coupled receptor

30 protein-encoding DNA is screened and a ligand for said human-derived G protein coupled receptor protein can be clarified using the above-mentioned method (1), the human-derived G protein coupled receptor protein-encoding DNA can be used as an agent for the prevention or therapy of the deficiency

35 diseases of said human-derived G protein coupled receptor protein depending upon the action that said ligand exhibits.

For example, when there is a patient for whom the physiological action of the ligand cannot be expected because of a decrease in the G protein coupled receptor protein in vivo, the amount of the G protein coupled receptor protein in the brain cells of said patient can be increased whereby the action of the ligand can be fully achieved by:

- (a) administering the G protein coupled receptor protein-encoding DNA to the patient to express it; or
- (b) inserting the G protein coupled receptor protein-encoding DNA into brain cells or the like to express it, followed by transplanting said brain cells or the like to said patient. Accordingly, the G protein coupled receptor protein-encoding DNA can be used as a safe and less toxic preventive and therapeutic agent for the G protein coupled receptor protein deficiency diseases.

When the G protein coupled receptor protein-encoding DNA is used as the above-mentioned agent, said DNA may be used alone or after inserting it into a suitable vector such as retrovirus vector, adenovirus vector, adenovirus-associated virus vector, etc. followed by subjecting the product vector to a conventional means. Thus, it may be administered orally parenterally, by inhalation spray, rectally, or topically as pharmaceutical compositions or formulations. Oral formulations include tablets (sugar-coated if necessary), capsules, elixirs, microcapsules, etc. Parenteral formulations include injections such as an aseptic solution or a suspension in water or in other pharmaceutically acceptable liquid. For example, the DNA of the present invention is admixed in a unit dose form which is required for preparing generally approved pharmaceutical preparations together with a physiologically acceptable carriers, flavoring agents, adjuvants, excipients, diluents, fillers, vehicles, antiseptics, stabilizers, binders, etc. whereupon the preparation can be manufactured. The amount of the effective component in those preparations is to be in such an extent that the suitable dose within an indicated range is achieved.

Examples of the additives which can be admixed in the

tablets, capsules, etc. are binders such as gelatin, corn starch, tragacanth and gum arabicum; fillers such as crystalline cellulose; swelling agents such as corn starch, gelatin and alginic acid; lubricating agents such as magnesium stearate; sweetening agents such as sucrose, lactose and saccharine; and flavoring agents such as pepper mint, akamono oil and cherry. When the unit dose form of the preparation is a capsule, a liquid carrier such as fat/oil may be further added in addition of the above-mentioned types of materials.

The aseptic composition for injection may be formulated by conventional practices for the preparations such as that the active substance in a vehicle such as water for injection is dissolved or suspended in naturally occurring plant oil such as sesame oil and palm oil.

Examples of an aqueous liquid for injection are a physiological saline solution and isotonic solutions containing glucose and other auxiliary agents (e.g. D-sorbitol, D-mannitol, sodium chloride, etc.) wherein a suitable auxiliary solubilizers such as alcohol (e.g. ethanol, etc.), polyalcohol (e.g. propylene glycol polyethylene glycol, etc.), nonionic surface-active agent (e.g. Polysorbate 80TM, HCO-50, etc.), etc. may be jointly used. Examples of an oily liquid include sesame oil, soybean oil, etc. wherein benzyl benzoate, benzyl alcohol, etc. may be jointly used as auxiliary solubilizers. In addition, buffers (e.g. phosphate buffer, sodium acetate buffer, etc.), analgesic agents (e.g. benzalkonium chloride, procaine hydrochloride, etc.), stabilizers (e.g. human serum albumin, polyethylene glycol, etc.), stabilizers (e.g. benzyl alcohol phenol, etc.), antioxidants, etc. may be admixed therewith too. The prepared injection solution is filled in suitable ampoules. The preparation prepared as such is safe and less toxic and, therefore, it can be administered to warm-blooded animals (e.g., rat, rabbit, sheep, swine, cattle, cat, dog, monkey, human beings, etc.).

Specific dose levels of said DNA may vary depending upon a variety of factors including the activity of drugs employed, the age, body weight, general health, sex, diet,

time of administration, route of administration, drug combination, and the severity of the symptom. In the case of oral administration, it is usually about 0.1-100 mg, preferably about 1.0-50 mg or, more preferably, about 1.0-20 mg per day
5 for adults (as 60 kg). When it is administered parenterally, its dose at a time may vary depending upon the object (patient) to be administered, organs to be administered, symptoms, administering methods, etc. but, in the case of injections, it is usually convenient to give by an intravenous route in an
10 amount of about 0.01-30 mg, preferably about 0.1-20 mg or, more preferably, about 0.1-10mg per day to adults (as 60 kg). In the case of other animals, the dose calculated for 60 kg may be administered as well.

15 (4) Quantitative Determination of Ligand to the G Protein Conjugated Receptor Protein of the Present Invention.

The G protein coupled receptor protein or a peptide fragment thereof has a binding property to ligand and, therefore, it is capable of determining quantitatively an amount of ligands in vivo with good sensitivity.

20 This quantitative determination may be carried out by, for example, combining with a competitive method. Thus, samples to be determined is contacted with G protein coupled receptor proteins or peptide fragments thereof so that the ligand concentration in said sample can be determined.
25 In one embodiment of the quantitative determination, the protocols described in the following ① and ② or the methods similar thereto may be used:

- ① Hiroshi Irie (ed): "Radioimmunoassay" (Kodansha, Japan, 1974); and
- 30 ② Hiroshi Irie (ed): "Radioimmunoassay, Second Series" (Kodansha, Japan, 1979).

(5) Screening of Compound Inhibiting the Binding of Ligand with the G Protein Conjugated Receptor Protein of the Present Invention.

35 G Protein coupled receptor proteins or peptide

fragments thereof are used. Alternatively, expression systems for recombinant type G Protein coupled receptor proteins or peptide fragments thereof are constructed and receptor binding assay systems using said expression system are used. In these
5 assay systems, it is possible to screen compounds (e.g. peptides, proteins, nonpeptidic compounds, synthetic compounds, fermented products, cell extracts, plant extracts, animal tissue extracts, etc.) or salts thereof which inhibits the binding of a ligand with the G protein coupled receptor
10 protein. Such a compound includes a compound exhibiting a G protein coupled receptor-mediated cell stimulating activity (e.g. activity of promoting or activity of inhibiting physiological reactions including liberation of arachidonic acid, liberation of acetylcholine, endocellular Ca^{2+} liberation,
15 endocellular cAMP production, endocellular cGMP production, production of inositol phosphate, changes in cell membrane potential, phosphorylation of endocellular proteins, activation of c-fos, lowering of pH, activation of G protein, cell promulgation, etc.) (so-called "G protein coupled
20 receptor-agonist"), a compound free of such a cell stimulating activity (so-called "G protein coupled receptor-antagonist"), etc.

Thus, the present invention provides a method of screening a compound which inhibits the binding of a ligand
25 with a G protein coupled receptor protein or a salt thereof, characterized in comparing the following two cases:
(i) the case wherein the ligand is contacted with the G protein coupled receptor protein or salt thereof, or a peptide fragment thereof or a salt thereof; and
30 (ii) the case wherein the ligand is contacted with a mixture of the G protein coupled receptor protein or salt thereof or the peptide fragment or salt thereof and said test compound.

In said screening method, one characteristic feature of the present invention resides in that the amount of the
35 ligand bonded with said G protein coupled receptor protein or the peptide fragment thereof, the cell stimulating activity of the ligand, etc. are measured in the case where (i) the

ligand is contacted with G protein coupled receptor proteins or peptide fragments thereof and in the case where (ii) the ligand and the test compound are contacted with the G protein coupled receptor protein or the peptide fragment thereof, respectively and then compared therebetween.

In one more specific embodiment of the present invention, the following is provided:

- ① a method of screening a compound or a salt thereof which inhibits the binding of a ligand with a G protein coupled receptor protein, characterized in that, when a labeled ligand is contacted with a G protein coupled receptor protein or a peptide fragment thereof and when a labeled ligand and a test compound are contacted with a G protein coupled receptor protein or a peptide fragment thereof, the amounts of the labeled ligand bonded with said protein or peptide fragment thereof or salt thereof are measured and compared;
- ② a method of screening a compound or a salt thereof which inhibits the binding of a ligand with a G protein coupled receptor protein, characterized in that, when a labeled ligand is contacted with cells containing G protein coupled receptor proteins or a membrane fraction of said cells and when a labeled ligand and a test compound are contacted with cells containing G protein coupled receptor proteins or a membrane fraction of said cells, the amounts of the labeled ligand binding with said protein or peptide fragment thereof or salt thereof are measured and compared;
- ③ a method of screening a compound or a salt thereof which inhibits the binding of a ligand with a G protein coupled receptor protein, characterized in that, when a labeled ligand is contacted with G protein coupled receptor proteins expressed on the cell membrane by culturing a transformant containing a G protein coupled receptor protein encoding DNA and when a labeled ligand and a test compound are contacted with G protein coupled receptor proteins expressed on the cell membrane by culturing a transformant containing a G protein coupled receptor protein encoding DNA, the amounts of the labeled ligand binding with said G protein coupled receptor

- 1 5 7 -

protein are measured and compared;
a method of screening a compound or a salt thereof which
④ inhibits the binding of a ligand with a G protein coupled
receptor protein, characterized in that, when a G protein
coupled receptor protein-activating compound (e.g. a ligand

5 to the G protein coupled receptor protein) is contacted with
cells containing G protein coupled receptor proteins and
when the G protein coupled receptor protein-activating compound
and a test compound are contacted with cells containing G
protein coupled receptor proteins, the resulting activities
10 coupled receptor protein-mediated cell stimulating activities
(e.g. activities of promoting or activities of inhibiting
physiological responses including liberation of arachdonic acid,
liberation of inositol phosphate, changes in cell membrane
production of c-fos, lowering of pH, activation of G protein,
15 potential, phosphorylation of endocellular proteins,
activation of c-fos, etc.) are measured and compared; and
cell promulgation, etc.) are measured and compared; and
a method of screening a compound or a salt thereof which

20 ⑤ inhibits the binding of a ligand with a G protein coupled
receptor protein, characterized in that, when a G protein coupled
receptor protein-activating compound (e.g. a ligand to
the G protein coupled receptor protein) is contacted with G
protein coupled receptor proteins expressed on cell membranes
25 by culturing transformants containing G protein coupled
receptor protein-encoding DNA and when a test compound are
contacted with the G protein coupled receptor protein
expressed on the cell membrane by culturing the transformant
30 containing the G protein coupled receptor protein-encoding DNA,
the resulting G protein coupled receptor protein-mediated cell
stimulating activities (activities of promoting or activities
of inhibiting physiological responses such as liberation of
35 arachdonic acid, liberation of acetylcholine, endocellular Ca^{2+}
liberation, endocellular cAMP production, endocellular cGMP
production, production of inositol phosphate, changes in cell

membrane potential, phosphorylation of endocellular proteins, activation of c-fos, lowering of pH, activation of G protein, and cell promulgation) are measured and compared.

Before the G protein coupled receptor protein of the present invention was obtained, the G protein coupled receptor agonist or antagonist had to be screened by, first, obtaining a candidate compound by using G protein coupled receptor protein-containing cells, tissues or cell membrane fractions derived from rat or the like (primary screening) and, then, making sure whether the candidate compound really inhibits the binding between human G protein coupled receptor proteins and ligands (secondary screening). Other receptor proteins inevitably exist when the cells, the tissues or the cell membrane fractions are used as they are, whereby they intrinsically make it difficult to screen agonists or antagonists to the desired receptor proteins. By using the human-derived G protein coupled receptor protein, however, there is no need of effecting the primary screening, whereby it is allowable to efficiently screen a compound that inhibits the binding between a ligand and a G protein coupled receptor. Besides, it is allowable to evaluate whether the compound that is screened is a G protein coupled receptor agonist or a G protein coupled receptor antagonist.

Specific explanations of the screening method will be given as hereunder.

First, with respect to the G protein coupled receptor protein used for the screening method of the present invention, any product may be used so far as it contains G protein coupled receptor proteins or peptide fragment thereof although the use of a membrane fraction of mammalian organs is suitable. However, human organs is extremely hardly available and, accordingly, G protein coupled receptor proteins which are expressed in a large amount using a recombinant are suitable for the screening.

In the manufacture of the G protein coupled receptor protein, the above-mentioned method can be used and it may be carried out by expressing the DNA coding for said protein in

mammalian cells or in insect cells. With respect to the DNA fragment coding for the target region, complementary DNA may be used although it is not limited thereto. Thus, for example, gene fragments or synthetic DNA may be used as well.

5 In order to introduce the G protein coupled receptor protein-encoding DNA fragment into host animal cells and to express it efficiently, it is preferred that said DNA fragment is incorporated into the downstream of polyhedron promoter of nuclear polyhedrosis virus belonging to baculovirus, promoter
10 derived from SV40, promoter of retrovirus, metallothionein promoter, human heat shock promoter, cytomegalovirus promoter, SRa promoter, etc. Examinations of the quantity and the quality of expressed receptors can be carried out by known methods per se or modified methods substantially analogous
15 thereto. For example, they may be conducted by the method described in publications such as Nambi, P. et al.: The Journal of Biochemical Society, vol.267, pages 19555-19559 (1992).

20 Accordingly, in the screening method, the substance containing a G protein coupled receptor protein or a peptide fragment thereof may be a G protein coupled receptor protein which is purified by known methods per se or a G protein coupled receptor protein fragment which is purified by known methods per se, or a cell containing said protein or a cell
25 membrane fraction of the cell containing said protein, etc.

When the G protein coupled receptor protein-containing cells are used in the screening method, said cells may be immobilized with glutaraldehyde, formalin, etc. The immobilization may be carried out by known methods per se
30 or modified methods substantially analogous thereto.

The G protein coupled receptor protein-containing cells are host cells expressing the G protein coupled receptor protein. Examples of said host cells may include Escherichia coli, Bacillus subtilis, yeasts, insect cells,
35 animal cells such as CHO cell and COS cell, etc.

Cell membrane fractions are fractions which contain a lot of cell membranes prepared by known methods per se or

modified methods substantially analogous thereto after disrupting or crushing the cells. Examples of disruptions of the cell may include methods by squeezing the cells with a Potter-Elvehjem homogenizer, disrupting or crushing by a Waring
5 blender or a Polytron (manufactured by Kinematica), disrupting or crushing by means of ultrasonic wave, disrupting by blowing out the cells from small nozzles together with applying a pressure with a French press or the like, etc.

Fractionation of the cell membrane is carried out mainly by
10 fractionation techniques by means of centrifugal force such as a fractional centrifugal separation and a density gradient centrifugal separation. For example, disrupted liquid of cells is centrifuged at a low speed (500 rpm to 3,000 rpm) for a short period (usually, from about one to ten minutes), the
15 supernatant liquid is further centrifuged at a high speed (1,500 rpm to 3,000 rpm) usually for 30 minutes to two hours and the resulting precipitate is used as a membrane fraction. Said membrane fraction contains a lot of expressed G protein coupled receptor proteins and membrane components such as
20 phospholipids and membrane proteins derived from the cells.

The amount of the G protein coupled receptor protein in the G protein coupled receptor protein-containing cell and in the cell membrane fraction obtained from the cell is preferably 10^3 - 10^8 molecules per cell or, suitably,
25 10^5 to 10^7 molecules per cell. Incidentally, the more the expressed amount, the higher the ligand binding activity (specific activity) per membrane fraction whereby the construction of a highly sensitive screening system is possible and, moreover, it is possible to measure the large
30 amount of samples in the same lot.

In conducting the above-mentioned methods ① to ③ for screening the compound capable of inhibiting the binding of the ligand with the G protein coupled receptor protein, a suitable G protein coupled receptor fraction and a labeled
35 ligand are necessary. With respect to the G protein coupled receptor fraction, it is preferred to use naturally occurring G protein coupled receptors (natural type G protein coupled

receptors) or recombinant type G protein coupled receptor fractions with the activity equivalent to that of the natural type G protein coupled. Here the term "activity equivalent to" means the same ligand binding activity, or the substantially equivalent ligand binding activity.

With respect to the labeled ligand, it is possible to use labeled ligands, labeled ligand analogized compounds, etc. For example, ligands labeled with [^3H], [^{125}I], [^{14}C], [^{35}S], etc. and other labeled substances may be utilized.

Specifically, G protein coupled receptor protein-containing cells or cell membrane fractions are first suspended in a buffer which is suitable for the determining method to prepare the receptor sample in conducting the screening for a compound which inhibits the binding of the ligand with the G protein coupled receptor protein.

With respect to the buffer, any buffer such as Tris-HCl buffer or phosphate buffer of pH 4-10 (preferably, pH 6-8) which does not inhibit the binding of the ligand with the receptor may be used.

In addition, a surface-active agent such as CHAPS, Tween 80TM (Kao-Atlas, Japan), digitonin, deoxycholate, etc. and/or various proteins such as bovine serum albumin (BSA), gelatine, etc. may be added to the buffer with an object of decreasing the nonspecific binding. Further, a protease inhibitor such as PMSF, leupeptin, E-64 (manufactured by Peptide Laboratory, Japan), pepstatin, etc. may be added with an object of inhibiting the decomposition of the receptor and the ligand by protease. A labeled ligand in a certain amount (5,000 cpm to 500,000 cpm) is added to 0.01 ml to 10 ml of said receptor solution and, at the same time, 10^{-4} M to 10^{-10} M of a test compound is made copresent. In order to determine the nonspecific binding amount (NSB), a reaction tube to which a great excessive amount of unlabeled test compounds is added is prepared as well.

The reaction is carried out at 0-50°C (preferably at 4-37°C) for 20 minutes to 24 hours (preferably 30 minutes to three hours). After the reaction, it is filtered through a

glass fiber filter, a filter paper, or the like, washed with a suitable amount of the same buffer and the radioactivity retained in the glass fiber filter, etc. is measured by means of a liquid scintillation counter or a gamma-counter.

5 Supposing that the count ($B_0 - \text{NSB}$) obtained by subtracting the nonspecific binding amount (NSB) from the total binding amount (B_0) wherein an antagonizing substance is not present is set at 100%, the test compound in which the specific binding amount ($B - \text{NSB}$) obtained by subtracting the nonspecific binding
10 amount (NSB) from the total binding amount (B) is, for example, less than 50% may be selected as a candidate ligand to the G protein coupled receptor protein of the present invention.

 In conducting the above-mentioned methods ④ to ⑤ for screening the compound which inhibits the binding of the
15 ligand with the G protein coupled receptor protein, the G protein coupled receptor protein-mediated cell stimulating activity (e.g. activities of promoting or activities of inhibiting physiological responses such as liberation of arachidonic acid, liberation of acetylcholine, endocellular
20 Ca^{2+} liberation, endocellular cAMP production, production of inositol phosphate, changes in the cell membrane potential, phosphorylation of endocellular proteins, activation of c-fos, lowering of pH, activation of G protein and cell promulgation, etc.) may be measured by known methods or by the use of
25 commercially available measuring kits. To be more specific, G protein coupled receptor protein-containing cells are at first cultured in a multiwell plate or the like.

 In conducting the screening, it is substituted with a suitable buffer which does not show toxicity to fresh media
30 or cells in advance, incubated for a certain period after adding a test compound, etc. thereto. The resultant cells are extracted or the supernatant liquid is recovered and the resulting product is determined, preferably quantitatively, by each of the methods. When it is difficult to identify the
35 production of the index substance (e.g. arachidonic acid, etc.) which is to be an index for the cell stimulating activity due to the presence of decomposing enzymes contained in the cell,

an assay may be carried out by adding an inhibitor against said decomposing enzyme. With respect to the activities such as an inhibitory action against cAMP production, it may be detected as an inhibitory action against the cAMP production in the cells whose fundamental production has been increased by forskolin or the like.

In conducting a screening by measuring the cell stimulating activity, cells in which a suitable G protein coupled receptor protein is expressed are necessary. Preferred G protein coupled receptor protein-expressing cells are naturally occurring G protein coupled receptor protein (natural type G protein coupled receptor protein)-containing cell lines or strains (e.g. mouse pancreatic β cell line, MIN6, etc.), the above-mentioned recombinant type G protein coupled receptor protein-expressing cell lines or strains, etc.

Examples of the test compound includes peptides, proteins, non-peptidic compounds, synthesized compounds, fermented products, cell extracts, plant extracts, animal tissue extracts, serum, blood, body fluid, etc. Those compounds may be novel or known.

A kit for screening the compound which inhibits the binding of the ligand with the G protein coupled receptor protein or a salt thereof of the present invention comprises a G protein coupled receptor protein or a peptide fragment thereof, or G protein coupled receptor protein-containing cells or cell membrane fraction thereof.

Examples of the screening kit include as follows:

1. Reagent for Determining Ligand.
- ① Buffer for Measurement and Buffer for Washing.

The product wherein 0.05% of bovine serum albumin (manufactured by Sigma) is added to Hanks' Balanced Salt Solution (manufactured by Gibco).

This may be sterilized by filtration through a membrane filter with a 0.45 μ m pore size, and stored at 4°C or may be prepared upon use.

② Sample of G Protein Conjugated Receptor Protein.

CHO cells in which a G protein coupled receptor protein is expressed are subcultured at the rate of 5×10^5 cells/well in a 12-well plate and cultured at 37°C with a 5% CO₂ and 95% air atmosphere for two days to prepare the sample.

③ Labeled Ligand.

The ligand which is labeled with commercially available [³H], [¹²⁵I], [¹⁴C], [³⁵S], etc.

The product in a state of an aqueous solution is stored at 4°C or at -20°C and, upon use, diluted to 1 μM with a buffer for the measurement.

④ Standard Ligand Solution.

Ligand is dissolved in PBS containing 0.1% of bovine serum albumin (manufactured by Sigma) to make 1 mM and stored at -20°C.

2. Method of the Measurement.

① CHO cells are cultured in a 12-well tissue culture plate to express G protein coupled receptor proteins. The G protein coupled receptor protein-expressing CHO cells are washed with 1 ml of buffer for the measurement twice. Then 490 μl of buffer for the measurement is added to each well.

② Five μl of a test compound solution of 10^{-3} to 10^{-10} M is added, then 5 μl of a labeled ligand is added and is made to react at room temperature for one hour. For knowing the non-specific binding amount, 5 μl of the ligand of 10^{-3} M is added instead of the test compound.

③ The reaction solution is removed from the well, which is washed with 1 ml of buffer for the measurement three times. The labeled ligand binding with the cells is dissolved in 0.2N NaOH-1% SDS and mixed with 4 ml of a liquid scintillator A (manufactured by Wako Pure Chemical, Japan).

④ Radioactivity is measured using a liquid scintillation counter (manufactured by Beckmann) and PMB (percent of maximum binding) is calculated by the following expression:

35

$$\text{PMB} = [(B - \text{NSB}) / (B_0 - \text{NSB})] \times 100$$

PMB: Percent of maximum binding
B: Value when a sample is added
NSB: Nonspecific binding
 B_0 : Maximum binding

5 The compound or a salt thereof obtained by the
screening method or by the screening kit is a compound which
inhibits the binding of a ligand with a G protein coupled
receptor protein and, more particularly, it is a compound
having a cell stimulating activity mediated via a G protein
10 coupled receptor or a salt thereof (so-called "G protein
coupled receptor agonist") or a compound having no said
stimulating activity (so-called "G protein coupled receptor
antagonist"). Examples of said compound are peptides,
proteins, non-peptidic compounds, synthesized compounds,
15 fermented products, etc. and the compound may be novel or
known.

 Said G protein coupled receptor agonist has the same
physiological action as the ligand to the G protein coupled
receptor protein has and, therefore, it is useful as a safe
20 and less toxic pharmaceutical composition depending upon said
ligand activity.

 On the other hand, said G protein coupled receptor
antagonist is capable of inhibiting the physiological activity
of the ligand to the G protein coupled receptor protein and,
25 there fore, it is useful as a safe and less toxic
pharmaceutical composition for inhibiting said ligand
activity.

 It is also strongly suggested that agonists and/or
antagonists related to the receptor encoded by pMAH2-17
30 obtained in Example 19 and/or the receptor encoded by phAH2-17
obtained in Example 21 would be useful in therapeutic or
prophylactic treatment of diseases or syndromes in connection
with purine ligand compounds or related analogues. It is
expected that the agonists of the receptor encoded by pMAH2-17
35 and/or of the receptor encoded by phAH2-17 are useful as an
immunomodulator or an antitumor agent, in addition they are

useful in therapeutically or prophylactically treating hypertension, diabetes, cystic fibrosis, etc. It is still expected that the antagonists of the receptor encoded by pMAH2-17 and/or of the receptor encoded by pAH2-17 are useful
5 as hypotensive agents, analgesics, agents for therapeutically or prophylactically treating incontinence of urine, etc. With regard to purinoceptors, the mutation of conserved basic amino acid residues in the 6th or 7th putative transmembrane domain of purinoceptors introduces alteration into the receptor's
10 responses to ATP (J. Biol. Chem., Vol. 270(9), pp. 4185-4188 (1995)). It is suggested that ATP is related to blood pressure control and circular systems via receptors (Circulation Research, Vol. 58(3), pp. 319-330 (1986)) and that ATP and purinoceptors are closely related (Am. Phys. Soc., pp.
15 C577-C606 (1993)).

When the compound or the salt thereof obtained by the screening method or by the screening kit is used as the above-mentioned pharmaceutical composition, a conventional means may be applied therefor. The compound or the salt
20 thereof may be orally, parenterally, by inhalation spray, rectally, or topically administered as pharmaceutical compositions or formulations (e.g. powders, granules, tablets, pills, capsules, injections, syrups, emulsions, elixirs, suspensions, solutions, etc.). For example, it may be used
25 by an oral route as tablets (sugar-coated if necessary), capsules, elixiers, microcapsules, etc. or by a parenteral route as injections such as an aseptic solution or a suspension in water or in other pharmaceutically acceptable liquid. The pharmaceutical compositions or formulations may
30 comprise at least one such compound alone or in admixture with pharmaceutically acceptable carriers, adjuvants, vehicles, excipients and/or diluents. The pharmaceutical compositions can be formulated in accordance with conventional methods. For example, said compound or the salt thereof is mixed in a
35 unit dose form which is required for preparing a generally approved pharmaceutical preparations together with a

physiologically acceptable carriers, flavoring and/or perfuming agents (fragrances), fillers, vehicles, antiseptics, stabilizers, binders, etc. whereupon the preparation can be manufactured. An amount of the effective component in those
5 preparations is to be in such an extent that the suitable dose within an indicated range is achieved.

Examples of the additives which can be admixed in the tablets, capsules, etc. are binders such as gelatin, corn starch, tragacanth and gum arabicum; fillers such as
10 crystalline cellulose; swelling agents such as corn starch, gelatin and alginic acid; lubricants such as magnesium stearate; sweetening agents such as sucrose, lactose and saccharine; preservatives such as parabens and sorbic acid; antioxidants such as ascorbic acid, α -tocopherol and cysteine;
15 fragrances such as peppermint, akamono oil and cherry; disintegrants; buffering agents; etc. Other additives may include mannitol, maltitol, dextran, agar, chitin, chitosan, pectin, collagen, casein, albumin, synthetic or semi-synthetic polymers, glyceride, lactide, etc. When the unit form of the
20 preparation is a capsule, a liquid carrier such as fat/oil may be further added besides the above-mentioned types of materials. The aseptic composition for injection may be formulated by a conventional technique or practice for the preparations such as that the active substance in a vehicle
25 such as water for injection is dissolved or suspended in a naturally occurring plant oil such as sesame oil and palm oil.

Examples of an aqueous liquid for the injection are a physiological saline solution and isotonic solutions containing glucose and other auxiliary agents (e.g. D-sorbitol,
30 D-mannitol, sodium chloride, etc.) wherein a suitable auxiliary solubilizers such as alcohol (e.g. ethanol, etc.), polyalcohol (e.g. propylene glycol, polyethylene glycol, etc.), nonionic surface-active agent (e.g. Polysorbate 80TM, HCO-50, etc.), etc. may be jointly used. In the case of the oily liquid,
35 sesame oil, soybean oil, etc. may be exemplified wherein benzyl benzoate, benzyl alcohol, etc. may be jointly used as auxiliary solubilizers.

In addition, buffers (e.g. phosphate buffer, sodium acetate buffer, etc.), analgesic agents (e.g. benzalkonium chloride, procaine hydrochloride, etc.), stabilizers (e.g. human serum albumin, polyethylene glycol, etc.), stabilizers
5 (e.g. benzyl alcohol, phenol, etc.), antioxidants, etc. may be compounded therewith too. The prepared injection solution is filled in suitable ampoules. The formulation prepared as such is safe and less toxic and, therefore, it can be administered to warm-blooded mammals such as rats, rabbits, sheep, swines,
10 cattle, cats, dogs, monkeys, human being, etc.

Dose levels of said compound or the salt thereof may vary depending upon the symptom. Specific dose levels for any particular patient will be employed depending upon a variety of factors including the activity of specific compounds employed,
15 the age, body weight, general health, sex, diet, time of administration, route of administration, rate of excretion, drug combination, and the severity of the particular disease undergoing therapy. In the case of oral administration, it is usually about 0.1-100 mg, preferably about 1.0-50 mg or, more
20 preferably, about 1.0-20 mg per day for adults (as 60 kg). When it is administered parenterally, its dose at a time may vary depending upon the object to be administered, organs to be administered, symptoms, administering methods, etc. The term "parenteral" as used herein includes subcutaneous injections,
25 intravenous, intramuscular, intraperitoneal injections, or infusion techniques. In the case of injections, it is usually convenient to give by an intravenous route in an amount of about 0.01-30 mg, preferably about 0.1-20 mg or, more preferably, about 0.1-10 mg per day to adults (as 60 kg). In the case of
30 other animals, the dose calculated for 60 kg may be administered as well.

(6) Manufacture of Antibody or Antiserum against the G Protein Coupled Receptor Protein of the Present Invention,
Its Peptide Fragment or Its Salt.

35 Antibodies (e.g. polyclonal antibody and monoclonal antibody) and antisera against the G protein coupled receptor

protein or salt thereof of the present invention or against the peptide fragment of the G protein coupled receptor protein or salt thereof of the present invention may be manufactured by antibody or antiserum-manufacturing methods per se known to those of skill in the art or methods similar thereto, using the G protein coupled receptor protein or its salt of the present invention or the peptide fragment of the G protein coupled receptor protein or its salt of the present invention. For example, monoclonal antibodies can be manufactured by the method as given below.

[Preparation of Monoclonal Antibody]

(a) Preparation of Monoclonal Antibody-Producing Cells.

The G protein coupled receptor protein of the present invention or its salt or the peptide fragment of the G protein coupled receptor protein of the present invention or its salt (hereinafter, may be abbreviated as the "G protein coupled receptor protein") is administered to warm-blooded animals either solely or together with carriers or diluents to the site where the production of antibody is possible by the administration. In order to potentiate the antibody productivity upon the administration, complete Freund's adjuvants or incomplete Freund's adjuvants may be administered. The administration is usually carried out once every two to six weeks and two to ten times in total. Examples of the applicable warm-blooded animals are monkeys, rabbits, dogs, guinea pigs, mice, rats, sheep, goats and chickens and the use of mice and rats is preferred.

In the preparation of the cells which produce monoclonal antibodies, an animal wherein the antibody titer is noted is selected from warm-blooded animals (e.g. mice) immunized with antigens, then spleen or lymph node is collected after two to five days from the final immunization and antibody-producing cells contained therein are fused with myeloma cells to give monoclonal antibody-producing hybridomas. Measurement of the antibody titer in antisera may, for example, be carried out by reacting a labeled

G protein coupled receptor protein (which will be mentioned later) with the antiserum followed by measuring the binding activity of the labeling agent with the antibody.

5 The operation for fusing may be carried out, for example, by a method of Koehler and Milstein (Nature, 256, 495, 1975). Examples of the fusion accelerator are polyethylene glycol (PEG), Sendai virus, etc. and the use of PEG is preferred.

10 Examples of the myeloma cells are NS-1, P3U1, SP2/0, AP-1, etc. and the use of P3U1 is preferred. The preferred fusion ratio of the numbers of antibody-producing cells used (spleen cells) to the numbers of myeloma cells is within a range of about 1:1 to 20:1. When PEG (preferably, PEG 1000 to PEG 6000) is added in a concentration of about 10-80% followed by incubating at 20-40°C (preferably, at 30-37°C) 15 for one to ten minutes, an efficient cell fusion can be carried out.

Various methods may be applied for screening a hybridoma which produces anti-G protein coupled receptor antibody. For example, a supernatant liquid of hybridoma 20 culture is added to a solid phase (e.g. microplate) to which the G protein coupled receptor protein antigen is adsorbed either directly or with a carrier, then anti-immunoglobulin antibody (anti-mouse immunoglobulin antibody is used when the cells used for the cell fusion are those of 25 mouse) which is labeled with a radioactive substance, an enzyme or the like, or protein A is added thereto and then anti-G protein coupled receptor monoclonal antibodies bound on the solid phase are detected; or a supernatant liquid of the hybridoma culture is added to the solid phase to which 30 anti-immunoglobulin or protein A is adsorbed, then the G protein coupled receptor labeled with a radioactive substance or an enzyme is added and anti-G protein coupled receptor monoclonal antibodies bonded with the solid phase is detected.

35 Selection and cloning of the anti-G protein coupled receptor monoclonal antibody-producing hybridoma may be carried out by methods per se known to those of skill in the art or methods similar thereto. Usually, it is carried out in a

medium for animal cells, containing HAT (hypoxanthine, aminopterin and thymidine). With respect to a medium for the selection, for the cloning and for the growth, any medium may be used so far as hybridoma is able to grow therein. Examples
5 of the medium are an RPMI 1640 medium (Dainippon Pharmaceutical Co., Ltd., Japan) containing 1-20% (preferably 10-20%) of fetal calf serum (FCS) , a GIT medium (Wako Pure Chemical, Japan) containing 1-20% of fetal calf serum and a serum-free medium for hybridoma culturing (SFM-101; Nissui Seiyaku, Japan).
10 The culturing temperature is usually 20-40°C and, preferably, about 37°C. The culturing time is usually from five days to three weeks and, preferably, one to two weeks. The culturing is usually carried out in 5% carbon dioxide gas. The antibody titer of the supernatant liquid of the hybridoma culture may be
15 measured by the same manner as in the above-mentioned measurement of the antibody titer of the anti-G protein coupled receptor in the antiserum.

The cloning can be usually carried out by methods known per se such as techniques in semi-solid agar and
20 limiting dilution. The cloned hybridoma is preferably cultured in modern serum-free culture media to obtain optimal amounts of antibody in supernatants. The target monoclonal antibody is also preferably obtained from ascitic fluid derived from a mouse, etc. injected intraperitoneally with
25 live hybridoma cells.

(b) Purification of the Monoclonal Antibody.

Like in the separation/purification of conventional polyclonal antibodies, the separation/purification of the anti-G protein coupled receptor monoclonal antibody may be
30 carried out by methods for separating/purifying immunoglobulin (such as salting-out, precipitation with an alcohol, isoelectric precipitation, electrophoresis, adsorption/deadsorption using ion exchangers such as DEAE, ultracentrifugation, gel filtration, specific purifying
35 methods in which only an antibody is collected by treatment with an active adsorbent (such as an antigen-binding solid

phase, protein A or protein G) and the bond is dissociated whereupon the antibody is obtained.

The G protein coupled receptor antibody of the present invention which is manufactured by the aforementioned method (a) or (b) is capable of specifically recognizing
5 G protein coupled receptors and, accordingly, it can be used for a quantitative determination of the G protein coupled receptor in test liquid samples and particularly for a quantitative determination by sandwich immunoassays.

10 Thus, the present invention provides, for example, the following methods:

- (i) a quantitative determination of a G protein coupled receptor in a test liquid sample, which comprises
 - (a) competitively reacting the test liquid sample and a
15 labeled G protein coupled receptor with an antibody which reacts with the G protein coupled receptor of the present invention, and
 - (b) measuring the ratio of the labeled G protein coupled receptor binding with said antibody; and
- 20 (ii) a quantitative determination of a G protein coupled receptor in a test liquid sample, which comprises
 - (a) reacting the test liquid sample with an antibody immobilized on an insoluble carrier and a labeled antibody simultaneously or continuously, and
 - 25 (b) measuring the activity of the labeling agent on the insoluble carrier

wherein one antibody is capable of recognizing the N-terminal region of the G protein coupled receptor while another antibody is capable of recognizing the C-terminal region of
30 the G protein coupled receptor.

When the monoclonal antibody of the present invention recognizing a G protein coupled receptor (hereinafter, may be referred to as "anti-G protein coupled receptor antibody") is used, G protein coupled receptors can be measured and,
35 moreover, can be detected by means of a tissue staining, etc. as well. For such an object, antibody molecules per se may be used or $F(ab')_2$, Fab' or Fab fractions of the antibody

molecule may be used too. There is no particular limitation for the measuring method using the antibody of the present invention and any measuring method may be used so far as it relates to a method in which the amount of antibody, antigen or antibody-antigen complex, depending on or corresponding to the amount of antigen (e.g. the amount of G protein coupled receptor, etc.) in the liquid sample to be measured, is detected by a chemical or a physical means and then calculated using a standard curve prepared by a standard solution containing the known amount of antigen. For example, nephrometry, competitive method, immunometric method and sandwich method are suitably used and, in terms of sensitivity and specificity, the sandwich method which will be described herein later is particularly preferred.

Examples of the labeling agent used in the measuring method using the labeling substance are radioisotopes, enzymes, fluorescent substances, luminescent substances, colloids, magnetic substances, etc. Examples of the radioisotope are [^{125}I], [^{131}I], [^3H] and [^{14}C]; preferred examples of the enzyme are those which are stable and with big specific activity, such as β -galactosidase, β -glucosidase, alkali phosphatase, peroxidase and malate dehydrogenase; examples of the fluorescent substance are fluorescamine, fluorescein isothiocyanate, etc.; and examples of the luminescent substance are luminol, luminol derivatives, luciferin, lucigenin, etc. Further, a biotin-avidin system may also be used for binding an antibody or antigen with a labeling agent.

In an insolubilization (immobilization) of antigens or antibodies, a physical adsorption may be used or a chemical binding which is usually used for insolubilization or immobilization of proteins or enzymes may be used as well. Examples of the carrier are insoluble polysaccharides such as agarose, dextran and cellulose; synthetic resins such as polystyrene, polyacrylamide and silicone; glass; etc.

In a sandwich (or two-site) method, the test liquid is made to react with an insolubilized anti-G protein coupled

receptor antibody (the first reaction), then it is made to react with a labeled anti-G protein coupled receptor antibody (the second reaction) and the activity of the labeling agent on the insoluble carrier is measured whereupon the amount of the G protein coupled receptor in the test liquid can be determined. The first reaction and the second reaction may be conducted reversely or simultaneously or they may be conducted with an interval. The type of the labeling agent and the method of insolubilization (immobilization) may be the same as those mentioned already herein. In the immunoassay by means of a sandwich method, it is not always necessary that the antibody used for the labeled antibody and the antibody for the solid phase is one type or one species but, with an object of improving the measuring sensitivity, etc., a mixture of two or more antibodies may be used too.

In the method of measuring G protein coupled receptors by the sandwich method of the present invention, the preferred anti-G protein coupled receptor antibodies used for the first and the second reactions are antibodies wherein their sites binding to the G protein coupled receptors are different each other. Thus, the antibodies used in the first and the second reactions are those wherein, when the antibody used in the second reaction recognizes the C-terminal region of the G protein coupled receptor, then the antibody recognizing the site other than C-terminal regions, e.g. recognizing the N-terminal region, is preferably used in the first reaction.

The anti-G protein coupled receptor antibody of the present invention may be used in a measuring system other than the sandwich method such as a competitive method, an immunometric method and a nephrometry. In a competitive method, an antigen in the test solution and a labeled antigen are made to react with an antibody in a competitive manner, then an unreacted labeled antigen (F) and a labeled antigen binding with an antibody (B) are separated (i.e. B/F separation) and the labeled amount of any of B and F is measured whereupon the amount of the antigen in the test

solution is determined. With respect to a method for such a reaction, there are a liquid phase method in which a soluble antibody is used as the antibody and the B/F separation is conducted by polyethylene glycol, a second antibody to the above-mentioned antibody, etc.; and a solid phase method in which an immobilized antibody is used as the first antibody or a soluble antibody is used as the first antibody while an immobilized antibody is used as the second antibody.

In an immunometric method, an antigen in the test solution and an immobilized antigen are subjected to a competitive reaction with a certain amount of a labeled antibody followed by separating into solid and liquid phases; or the antigen in the test solution and an excess amount of labeled antibody are made to react, then a immobilized antigen is added to bind an unreacted labeled antibody with the solid phase and separated into solid and liquid phases. After that, the labeled amount of any of the phases is measured to determine the antigen amount in the test solution.

In a nephrometry, the amount of insoluble sediment which is produced as a result of the antigen-antibody reaction in a gel or in a solution is measured. Even when the antigen amount in the test solution is small and only a small amount of the sediment is obtained, a laser nephrometry wherein scattering of laser is utilized can be suitably used.

In applying each of those immunological measuring methods (immunoassays) to the measuring method of the present invention, it is not necessary to set up any special condition, operation, etc. therefor. A measuring system (assay system) for G protein coupled receptor may be constructed taking the technical consideration of the persons skilled in the art into consideration in the conventional conditions and operations for each of the methods. With details of those conventional technical means, a variety of reviews, reference books, etc. may be referred to. They are, for example, Hiroshi Irie (ed): "Radioimmunoassay" (Kodansha, Japan, 1974); Hiroshi Irie (ed): "Radioimmunoassay; Second Series" (Kodansha, Japan, 1979); Eiji Ishikwa et al. (ed): "Enzyme Immunoassay" (Igaku Shoin,

Japan, 1978); Eiji Ishikawa et al. (ed): "Enzyme Immunoassay" (Second Edition) (Igaku Shoin, Japan, 1982); Eiji Ishikawa et al. (ed): "Enzyme Immunoassay" (Third Edition) (Igaku Shoin, Japan, 1987); "Methods in Enzymology" Vol. 70 (Immunochemical Techniques (Part A)); ibid. Vol. 73 (Immunochemical Techniques (Part B)); ibid. Vol. 74 (Immunochemical Techniques (Part C)); ibid. Vol. 84 (Immunochemical Techniques (Part D: Selected Immunoassays)); ibid. Vol. 92 (Immunochemical Techniques (Part E: Monoclonal Antibodies and General Immunoassay Methods)); ibid. Vol. 121 (Immunochemical Techniques (Part I: Hybridoma Technology and Monoclonal Antibodies)) (Academic Press); etc.

(7) Preparation of Animals Having the G Protein Coupled Receptor Protein-Encoding DNA of the Present Invention.

It is possible to prepare transgenic animals expressing G protein coupled receptors using G protein coupled receptor protein-encoding DNA. Examples of the animals are warm-blooded mammals such as rats, rabbit, sheep, swines, cattle, cats, dogs and monkeys.

In transferring the G protein coupled receptor protein-encoding DNA to the aimed animal, it is generally advantageous that said DNA is used by ligating with a site at the downstream of a promoter which is capable of expressing in animal cells. For example, when G protein coupled receptor protein DNA is to be transferred to a rabbit, a gene construct ligated with a site at the downstream of various promoters which are capable of expressing the G protein coupled receptor protein DNA derived from an animal compatible to the animal in animal host cells is subjected to a microinjection to the fertilized ovum (oosperm) of the aimed animal (e.g. fertilized ovum (embryo) of rabbit) whereupon the transgenic animal which produces the G protein coupled receptor protein in a high amount can be prepared.

Examples of the promoters used are promoters derived from virus and ubiquitous expression promoters such as metallothionein promoters may be used but, preferably,

enolase gene promoters and NGF gene promoters capable of specifically expressing in brain are used.

Transfer of the G protein coupled receptor protein DNA at a fertilized ovum cell stage is secured in order that the DNA can be present in all of embryonal cells and body somatic cells of an aimed animal. The fact that the G protein coupled receptor protein is present in the fertilized ovum cells of the produced transgenic animal after the DNA transfer means that all progeny of the produced transgenic animal have the G protein coupled receptor protein in all of their embryonal cells and somatic cells. Descendants (offsprings) of the animal of this type which inherited the gene have the G protein coupled receptor protein in all of their embryonal cells and somatic cells.

The transgenic animal to which the G protein coupled receptor protein DNA is transferred can be subjected to a mating and a breeding for generations under a common breeding circumstance as the animal holding said DNA after confirming that the gene can be stably retained. Moreover, male and female animals having the desired DNA are mated to give a homozygote having the transduced gene in both homologous chromosomes and then those male and female animals are mated whereby it is possible to breed for generations so that all descendants have said DNA.

The animal to which the G protein coupled receptor protein DNA is transferred highly expresses the G protein coupled receptor protein and, accordingly, it is useful as the animal for screening for an agonist or an antagonist to said G protein coupled receptor protein.

The DNA-transferred animal can be used as a cell source for a tissue culture. For example, DNA or RNA in the tissue of the DNA-transferred mouse is directly analyzed or protein tissues expressed by gene are analyzed whereupon the G protein coupled receptor protein can be analyzed. Cells of the G protein coupled receptor protein-containing tissue are cultured by standard tissue culture techniques whereupon it is possible to study the function of the cells

which are usually difficult to culture (e.g. those derived from brain and peripheral tissues) using the resulting culture. By using said cells, it is also possible to select the pharmaceuticals which can potentiate, for example, the functions of various tissues. Moreover, if a cell strain with a high expression is available, it is possible to separate and purify G protein coupled receptor proteins therefrom.

As such, the amount of G protein coupled receptor proteins can now be determined with a high precision using the anti-G protein coupled receptor antibody of the present invention.

(8) Antisense Oligonucleotides Capable of Inhibiting Replication of G Protein Coupled Receptor Protein Gene

In another aspect of the present invention, antisense oligonucleotides (nucleic acids) capable of inhibiting the replication or expression of G protein coupled receptor protein gene may be designed and synthesized based on information on the nucleotide sequences of cloned and determined G protein coupled receptor protein-encoding DNAs. Such an antisense oligonucleotide (nucleic acid) is capable of hybridizing with RNA of G protein coupled receptor protein genes to inhibit the synthesis or function of said RNA or of modulating the expression of a G protein coupled receptor protein gene via interaction with G protein coupled receptor protein-related RNA. Oligonucleotides complementary to, and specifically hybridizable with, selected sequences of G protein coupled receptor protein-related RNA are useful in controlling or modulating the expression of a G protein coupled receptor protein gene in vitro and in vivo, and in treating or diagnosing disease states of suspected animals. The term "corresponding" means homologous to or complementary to a particular sequence of the nucleotide sequence or nucleic acid including the gene. As between nucleotides (nucleic acids) and peptides (proteins), "corresponding" usually refers to amino acids of a peptide (protein) in an order derived from the sequence of a nucleotides (nucleic acids) or its complement.

The G protein coupled receptor protein gene 5' end hairpin loop, 5' end 6-base-pair repeats, 5' end untranslated region, polypeptide translation initiation codon, protein coding region, ORF translation initiation codon, 3' untranslated region, 3' end palindrome region, and 3' end hairpin loop may be selected as preferred targets though any region may be a target among G protein coupled receptor protein genes. The relationship between the target and oligonucleotides complementary to at least a portion of the target, specifically hybridizable with the target, is denoted as "antisense". The antisense oligonucleotides may be polydeoxynucleotides containing 2-deoxy-D-ribose, polyribonucleotides containing D-ribose, any other type of polynucleotide which is an N-glycoside of a purine or pyrimidine base, or other polymers containing nonnucleotide backbones (e.g., protein nucleic acids and synthetic sequence-specific nucleic acid polymers commercially available) or nonstandard linkages, providing that the polymers contain nucleotides in a configuration which allows for base pairing and base stacking such as is found in DNA and RNA. They may include double- and single-stranded DNA, as well as double- and single-stranded RNA and DNA:RNA hybrids, and also include, as well as unmodified forms of the polynucleotide or oligonucleotide, known types of modifications, for example, labels which are known to those skilled in the art, "caps", methylation, substitution of one or more of the naturally occurring nucleotides with analogue, internucleotide modifications such as, for example, those with uncharged linkages (e.g., methyl phosphonates, phosphotriesters, phosphoramidates, carbamates, etc.) and with charged linkages or sulfur-containing linkages (e.g., phosphorothioates, phosphorodithioates, etc.), those containing pendant moieties, such as, for example, proteins (including nucleases, nuclease inhibitors, toxins, antibodies, signal peptides, poly-L-lysine, etc.) and saccharides (e.g., monosaccharides, etc.), those with intercalators (e.g., acridine, psoralen, etc.), those containing chelators (e.g., metals, radioactive metals, boron, oxidative metals, etc.),

those containing alkylators, those with modified linkages (e.g., alpha anomeric nucleic acids, etc.). The terms "nucleoside", "nucleotide" and "nucleic acid" will include those moieties which contain not only the known purine and pyrimidine bases, but also other heterocyclic bases which have been modified. Such modifications include methylated purines and pyrimidines, acylated purines and pyrimidines, or other heterocycles. Modified nucleosides or nucleotides will also include modifications on the sugar moiety, e.g., wherein one or more of the hydroxyl groups are replaced with halogen, aliphatic groups, or are functionalized as ethers, amines, or the like.

The antisense nucleic acid of the present invention is RNA, DNA or a modified nucleic acid. Examples of modified nucleic acid are, but not limited to, degradation-resistant sulfurized and thiophosphate derivatives of nucleic acids, and poly- or oligonucleoside amides. Preferred design modifications of the antisense nucleic acids of the present invention are modifications that are designed to:

- (1) increase the intracellular stability of the nucleic acid;
- (2) increase the cellular permeability of the nucleic acid;
- (3) increase the affinity of the nucleic acid for the target sense strand; or
- (4) decrease the toxicity (if any) of the nucleic acid.

Many such modifications are known to those skilled in the art, as described in J. Kawakami et al., Pharm Tech Japan, Vol. 8, pp.247, 1992; Vol. 8, pp.395, 1992; S. T. Crooke et al. ed., Antisense Research and Applications, CRC Press, 1993; etc.

The nucleic acids may contain altered or modified sugars, bases or linkages, be delivered in specialized systems such as liposomes, microspheres or by gene therapy, or may have attached moieties. Such attached moieties include polycationic moieties such as polylysine that act as charge neutralizers of the phosphate backbone, or hydrophobic moieties such as lipids (e.g., phospholipids, cholesterol, etc.) that enhance interaction with cell membranes or increase uptake of the nucleic acid. Preferred lipids that may be attached are

cholesterols or derivatives thereof (e.g., cholesteryl
chloroformate, cholic acid, etc.). The moieties may be
attached at the 3' or 5' ends of the nucleic acids, and also
may be attached through a base, sugar, or internucleoside
5 linkage. Other moieties may be capping groups specifically
placed at the 3' or 5' ends of the nucleic acids to prevent
degradation by nuclease such as exonuclease, RNase, etc. Such
capping groups include, but are not limited to, hydroxyl
protecting groups known to those skilled in the art, including
10 glycols such as polyethylene glycols, tetraethylene glycol and
the like.

The inhibitory activity of antisense nucleic acids
can be examined using the transformant (or transfectant) of
the present invention, the in vitro and in vivo gene
15 expression system of the present invention, or the in vitro
and in vivo translation system of G protein coupled receptor
proteins. The nucleic acid can be placed in the cell through
any number of ways known per se.

In the specification and drawings of the present
20 application, the abbreviations used for bases (nucleotides),
amino acids and so forth are those recommended by the IUPAC-IUB
Commission on Biochemical Nomenclature or those conventionally
used in the art. Examples thereof are given below.
Amino acids for which optical isomerism is possible are, unless
25 otherwise specified, in the L form.

DNA : Deoxyribonucleic acid
cDNA: Complementary deoxyribonucleic acid
A : Adenine
T : Thymine
30 G : Guanine
C : Cytosine
RNA : Ribonucleic acid
mRNA : Messenger ribonucleic acid
dATP: Deoxyadenosine triphosphate
35 dTTP: Deoxythymidine triphosphate

	dGTP:	Deoxyguanosine triphosphate
	dCTP:	Deoxycytidine triphosphate
	ATP :	Adenosine triphosphate
	EDTA:	Ethylenediamine tetraacetic acid
5	SDS :	Sodium dodecyl sulfate
	EIA:	Enzyme Immunoassay
	G, Gly:	Glycine (or Glycyl)
	A, Ala:	Alanine (or Alanyl)
	V, Val:	Valine (or Valyl)
10	L, Leu:	Leucine (or Leucyl)
	I, Ile:	Isoleucine (or Isoleucyl)
	S, Ser:	Serine (or Seryl)
	T, Thr:	Threonine (or Threonyl)
	C, Cys:	Cysteine (or Cysteinyl)
15	M, Met:	Methionine (or Methionyl)
	E, Glu:	Glutamic acid (or Glutamyl)
	D, Asp:	Aspartic acid (or Aspartyl)
	K, Lys:	Lysine (or Lysyl)
	R, Arg:	Arginine (or Arginyl)
20	H, His:	Histidine (or Histidyl)
	F, Phe:	Pheylalanine (or Pheylalanyl)
	Y, Tyr:	Tyrossine (or Tyrosyl)
	W, Trp:	Tryptophan (or Tryptophanyl)
	P, Pro:	Proline (or Prolyl)
25	N, Asn:	Asparagine (or Asparaginyll)
	Q, Gln:	Glutamine (or Glutaminyl)
	NVal:	Norvaline (or Norvalyl)
	pGlu:	Pyroglutamic acid (or Pyroglutamyl)
	Blc:	γ -Butyrolacton- γ -carbonyl
30	Kpc:	2-Ketopiperidinyl-6-carbonyl
	Otc:	3-Oxoperhydro-1,4-thiazin-5-carbonyl
	Me:	Methyl
	Et:	Ethyl
	Bu:	Butyl
35	Ph:	Phenyl
	TC:	Thiazolidinyl-4(R)-carboxamide

The transformant *Escherichia coli*, designated INVα F'/p19P2, which is obtained in the Example 3 mentioned herein below, is on deposit under the terms of the Budapest Treaty from August 9, 1994, with the National Institute of Bioscience and Human-Technology (NIBH), Agency of Industrial Science and Technology, Ministry of International Trade and Industry, Japan and has been assigned the Accession Number FERM BP-4776. It is also on deposit from August 22, 1994 with the Institute for Fermentation, Osaka, Japan (IFO) and has been assigned the Accession Number IFO 15739.

The transformant *Escherichia coli*, designated INVα F'/pG3-2, which is obtained in the Example 4 mentioned herein below, is on deposit under the terms of the Budapest Treaty from August 9, 1994, with NIBH and has been assigned the Accession Number FERM BP-4775. It is also on deposit from August 22, 1994 with IFO and has been assigned the Accession Number IFO 15740.

The transformant *Escherichia coli*, designated INVα F'/p63A2, which is obtained in the Example 5 mentioned herein below, is on deposit under the terms of the Budapest Treaty from August 9, 1994, with NIBH and has been assigned the Accession Number FERM BP-4777. It is also on deposit from August 22, 1994 with IFO and has been assigned the Accession Number IFO 15738.

The transformant *Escherichia coli*, designated JM109/phGR3, which is obtained in the Example 6 mentioned herein below, is on deposit under the terms of the Budapest Treaty from September 27, 1994, with NIBH and has been assigned the Accession Number FERM BP-4807. It is also on deposit from September 22, 1994 with IFO and has been assigned the Accession Number IFO 15748.

The transformant *Escherichia coli*, designated JM109/p3H2-17, which is obtained in the Example 7 mentioned herein below, is on deposit under the terms of the Budapest Treaty from September 27, 1994, with NIBH and has been assigned the Accession Number FERM BP-4806. It is also on deposit from September 22, 1994 with IFO and has been assigned the Accession

Number IFO 15747.

The transformant *Escherichia coli*, designated JM109/p3H2-34, which is obtained in the Example 8 mentioned herein below, is on deposit under the terms of the Budapest Treaty from October 12, 1994, with NIBH and has been assigned the Accession Number FERM BP-4828. It is also on deposit from October 12, 1994 with IFO and has been assigned the Accession Number IFO 15749.

The transformant *Escherichia coli*, designated JM109/pMD4, which is obtained in the Example 9 mentioned herein below, is on deposit under the terms of the Budapest Treaty from November 11, 1994, with NIBH and has been assigned the Accession Number FERM BP-4888. It is also on deposit from November 17, 1994 with IFO and has been assigned the Accession Number IFO 15765.

The transformant *Escherichia coli*, designated JM109/pMGR20, which is obtained in the Example 10 mentioned herein below, is on deposit under the terms of the Budapest Treaty from December 15, 1994, with NIBH and has been assigned the Accession Number FERM BP-4937. It is also on deposit from December 14, 1994 with IFO and has been assigned the Accession Number IFO 15773.

The transformant *Escherichia coli*, designated JM109/pMJ10, which is obtained in the Example 12 mentioned herein below, is on deposit under the terms of the Budapest Treaty from December 15, 1994, with NIBH and has been assigned the Accession Number FERM BP-4936. It is also on deposit from December 16, 1994 with IFO and has been assigned the Accession Number IFO 15784.

The transformant *Escherichia coli*, designated JM109/pMH28, which is obtained in the Example 14 mentioned herein below, is on deposit under the terms of the Budapest Treaty from January 13, 1995, with NIBH and has been assigned the Accession Number FERM BP-4970. It is also on deposit from January 20, 1995 with IFO and has been assigned the Accession Number IFO 15791.

The transformant *Escherichia coli*, designated

JM109/pMN7, which is obtained in the Example 16 mentioned herein below, is on deposit under the terms of the Budapest Treaty from February 22, 1995, with NIBH and has been assigned the Accession Number FERM BP-5011. It is also on deposit from
5 February 27, 1995 with IFO and has been assigned the Accession Number IFO 15803.

The transformant *Escherichia coli*, designated JM109/p5S38, which is obtained in the Example 17 mentioned herein below, is on deposit under the terms of the Budapest
10 Treaty from October 27, 1994, with NIBH and has been assigned the Accession Number FERM BP-4856. It is also on deposit from October 25, 1994 with IFO and has been assigned the Accession Number IFO 15754.

The transformant *Escherichia coli*, designated
15 JM109/pMAH2-17, which is obtained in the Example 19 mentioned herein below, is on deposit under the terms of the Budapest Treaty from April 7, 1995, with NIBH and has been assigned the Accession Number FERM BP-5073. It is also on deposit from March 31, 1995 with IFO and has been assigned the Accession
20 Number IFO 15813.

The transformant *Escherichia coli*, designated JM109/pMN128, which is obtained in the Example 20 mentioned herein below, is on deposit under the terms of the Budapest Treaty from March 17, 1995, with NIBH and has been assigned the
25 Accession Number FERM BP-5039. It is also on deposit from March 22, 1995 with IFO and has been assigned the Accession Number IFO 15810.

The transformant *Escherichia coli*, designated JM109/pAH2-17, which is obtained in the Example 21 mentioned
30 herein below, is on deposit under the terms of the Budapest Treaty from July 20, 1995, with NIBH and has been assigned the Accession Number FERM BP-5168. It is also on deposit from July 14, 1995 with IFO and has been assigned the Accession Number IFO 15856.

35 Each SEQ ID NO set forth in the SEQUENCE LISTING of the specification refers to the following sequence:

[SEQ ID NO: 24] is a partial amino acid sequence of the human pituitary gland-derived G protein coupled receptor protein encoded by the human pituitary gland-derived G protein coupled receptor protein cDNA fragment included in p19P2,

5 [SEQ ID NO: 25] is a partial amino acid sequence of the human pituitary gland-derived G protein coupled receptor protein encoded by the human pituitary gland-derived G protein coupled receptor protein cDNA fragment included in p19P2,

10 [SEQ ID NO: 26] is an entire amino acid sequence of the human pituitary gland-derived G protein coupled receptor protein encoded by the human pituitary gland-derived G protein coupled receptor protein cDNA fragment included in phGR3,

[SEQ ID NO: 27] is a partial amino acid sequence of the mouse pancreatic β -cell line, MIN6-derived G protein coupled
15 receptor protein encoded by the mouse pancreatic β -cell line, MIN6-derived G protein coupled receptor protein cDNA fragment having a nucleotide sequence (SEQ ID NO: 32), derived based upon the nucleotide sequences of the mouse pancreatic β -cell line, MIN6-derived G protein coupled receptor protein cDNA
20 fragments each included in pG3-2 and pG1-10,

[SEQ ID NO: 28] is a partial amino acid sequence of the mouse pancreatic β -cell line, MIN6-derived G protein coupled receptor protein encoded by p5S38,

[SEQ ID NO: 29] is a nucleotide sequence of the human
25 pituitary gland-derived G protein coupled receptor protein cDNA fragment included in p19P2,

[SEQ ID NO: 30] is a nucleotide sequence of the human pituitary gland-derived G protein coupled receptor protein cDNA fragment included in p19P2,

30 [SEQ ID NO: 31] is an entire nucleotide sequence of the human pituitary gland-derived G protein coupled receptor protein cDNA included in phGR3,

[SEQ ID NO: 32] is a nucleotide sequence of the mouse pancreatic β -cell line, MIN6-derived G protein coupled
35 receptor protein cDNA, derived based upon the nucleotide sequences of the mouse pancreatic β -cell line, MIN6-derived G protein coupled receptor protein cDNA fragments each

included in pG3-2 and pG1-10,

[SEQ ID NO: 33] is a nucleotide sequence of the mouse pancreatic β -cell line, MIN6-derived G protein cDNA included in p5S38,

5 [SEQ ID NO: 34] is a partial amino acid sequence of the human amygdaloid nucleus-derived G protein coupled receptor protein encoded by the cDNA fragment included in p63A2,

[SEQ ID NO: 35] is a partial amino acid sequence of the human amygdaloid nucleus-derived G protein coupled receptor protein
10 encoded by the cDNA fragment included in p63A2,

[SEQ ID NO: 36] is a nucleotide sequence of the human amygdaloid nucleus-derived G protein coupled receptor protein cDNA fragment included in p63A2,

[SEQ ID NO: 37] is a nucleotide sequence of the human
15 amygdaloid nucleus-derived G protein coupled receptor protein cDNA fragment included in p63A2,

[SEQ ID NO: 38] is a partial amino acid sequence encoded by the mouse pancreatic β -cell line, MIN6-derived G protein coupled receptor protein cDNA included in p3H2-17,

20 [SEQ ID NO: 39] is a full-length amino acid sequence encoded by the open reading frame of the mouse pancreatic β -cell line, MIN6-derived G protein coupled receptor protein cDNA included in pMAH2-17,

[SEQ ID NO: 40] is a nucleotide sequence of the mouse
25 pancreatic β -cell line, MIN6-derived G protein coupled receptor protein cDNA included in p3H2-17,

[SEQ ID NO: 41] is a nucleotide sequence of the mouse pancreatic β -cell line, MIN6-derived G protein coupled receptor protein cDNA included in pMAH2-17,

30 [SEQ ID NO: 42] is a partial amino acid sequence encoded by the mouse pancreatic β -cell line, MIN6-derived G protein coupled receptor protein cDNA included in p3H2-34,

[SEQ ID NO: 43] is a nucleotide sequence of the mouse pancreatic β -cell line, MIN6-derived G protein coupled
35 receptor protein cDNA fragment included in p3H2-34,

[SEQ ID NO: 44] is a partial amino acid sequence encoded by the rabbit gastropyloric part smooth muscle-derived G

protein coupled receptor protein cDNA included in pMD4,
[SEQ ID NO: 45] is a nucleotide sequence of the rabbit
gastropyrolic part smooth muscle-derived G protein coupled
receptor protein cDNA fragment included in pMD4,
5 [SEQ ID NO: 46] is an entire amino acid sequence
encoded by the mouse pancreatic β -cell line, MIN6-derived G
protein coupled receptor protein cDNA included in pMGR20,
[SEQ ID NO: 47] is a nucleotide sequence of the mouse
pancreatic β -cell line, MIN6-derived G protein coupled
10 receptor protein cDNA included in pMGR20,
[SEQ ID NO: 48] is a partial amino acid sequence encoded
by the rabbit gastropyrolic part smooth muscle-derived G
protein coupled receptor protein cDNA included in pMJ10,
[SEQ ID NO: 49] is a nucleotide sequence of the rabbit
15 gastropyrolic part smooth muscle-derived G protein coupled
receptor protein cDNA fragment included in pMJ10,
[SEQ ID NO: 50] is a partial amino acid sequence encoded
by the rabbit gastropyrolic part smooth muscle-derived G
protein coupled receptor protein cDNA included in pMH28,
20 [SEQ ID NO: 51] is a nucleotide sequence of the rabbit
gastropyrolic part smooth muscle-derived G protein coupled
receptor protein cDNA fragment included in pMH28,
[SEQ ID NO: 52] is a partial amino acid sequence encoded
by the rabbit gastropyrolic part smooth muscle-derived G
25 protein coupled receptor protein cDNA included in pMN7,
[SEQ ID NO: 53] is a nucleotide sequence of the rabbit
gastropyrolic part smooth muscle-derived G protein coupled
receptor protein cDNA fragment included in pMN7,
[SEQ ID NO: 54] is a partial amino acid sequence encoded
30 by the rabbit gastropyrolic part smooth muscle-derived G
protein coupled receptor protein cDNA included in pMN128,
[SEQ ID NO: 55] is a nucleotide sequence of the rabbit
gastropyrolic part smooth muscle-derived G protein coupled
receptor protein cDNA fragment included in pMN128,
35 [SEQ ID NO: 56] is a full-length amino acid sequence of the
human-derived G protein coupled receptor protein encoded
by the human-derived G protein coupled receptor protein cDNA

included in phAH2-17, and
[SEQ ID NO: 57] is a nucleotide sequence of the human-derived
G protein coupled receptor protein cDNA included in phAH2-17.

EXAMPLES

5 Described below are working examples of the present
invention which are provided only for illustrative purposes,
and not to limit the scope of the present invention. In light
of the present disclosure, numerous embodiments within the
scope of the claims will be apparent to those of ordinary skill
10 in the art.

Example 1

Preparation of Synthetic DNA Primer for Amplifying DNA Coding for G Protein Coupled Receptor Protein

A comparison of deoxyribonucleotide sequences
15 coding for the known amino acid sequences corresponding to
or near the first membrane-spanning domain each of
human-derived TRH receptor protein (HTRHR), human-derived
RANTES receptor protein (L10918, HUMRANTES), human Burkitt's
lymphoma-derived unknown ligand receptor protein (X68149,
20 HSBRLR1A), human-derived somatostatin receptor protein
(L14856, HUMSOMAT), rat-derived μ -opioid receptor protein
(U02083, RNU02083), rat-derived κ -opioid receptor protein
(U00442, U00442), human-derived neuromedin B receptor protein
(M73482, HUMNMBR), human-derived muscarinic acetylcholine
25 receptor protein (X15266, HSHM4), rat-derived adrenaline
 α_1 B receptor protein (L08609, RATAADRE01), human-derived
somatostatin 3 receptor protein (M96738, HUMSSTR3X),
human-derived C_5a receptor protein (HUMC5AAR), human-derived
unknown ligand receptor protein (HUMRDC1A), human-derived
30 unknown ligand receptor protein (M84605, HUMOPIODRE) and
rat-derived adrenaline α_2 B receptor protein (M91466,
RATA2BAR) was made. As a result, highly homologous
regions or parts were found (Figure 1).

Further, a comparison of deoxynucleotide sequences
35 coding for the known amino acid sequences corresponding to or
near the sixth membrane-spanning domain each of mouse-derived

unknown ligand receptor protein (M80481, MUSGIR), human-derived bombesin receptor protein (L08893, HUMBOMB3S), human-derived adenosine A2 receptor protein (S46950, S46950), mouse-derived unknown ligand receptor protein (D21061, MUSGPCR), mouse-
5 derived TRH receptor protein (S43387, S43387), rat-derived neuromedin K receptor protein (J05189, RATNEURA), rat-derived adenosine A1 receptor protein (M69045, RATA1ARA), human-derived neurokinin A receptor protein (M57414, HUMNEKAR), rat-derived adenosine A3 receptor protein (M94152, RATADENREC), human-
10 derived somatostatin 1 receptor protein (M81829, HUMSRI1A), human-derived neurokinin 3 receptor protein (S86390, S86371S4), rat-derived unknown ligand receptor protein (X61496, RNCGPCR), human-derived somatostatin 4 receptor protein (L07061, HUMSSTR4Z) and rat-derived GnRH receptor protein
15 (M31670, RATGNRHA) was made. As a result, highly homologous regions or parts were found (Figure 2).

The aforementioned abbreviations in the parentheses are identifiers (reference numbers) which are indicated when GenBank/EMBL Data Bank is retrieved by using DNASIS
20 Gene/Protein Sequencing Data Base (CD019, Hitachi Software Engineering, Japan) and are usually called "Accession Numbers" or "Entry Names". HTRHR is, however, the sequence as disclosed in Japanese Unexamined Patent Publication No. 286986/1993 (EPA 638645).

25 Specifically, it was planned to incorporate mixed bases relying upon the base regions that were in agreement with cDNAs coding for a large number of receptor proteins in order to enhance base agreement of sequences with as many receptor cDNAs as possible even in other regions. Based upon
30 these sequences, the degenerate synthetic DNA having a nucleotide sequence represented by SEQ ID NO: 1 which is complementary to the homologous nucleotide sequence of Figure 1 and the degenerate synthetic DNA having a nucleotide sequence represented by SEQ ID NO: 2 which is complementary to the
35 homologous nucleotide sequence of Figure 2 were produced. Nucleotide synthesis was carried out by a DNA synthesizer.

- 1 9 1 -

[Synthetic DNAs]

5'-CGTGG (G or C) C (A or C) T (G or C) (G or C) TGGGCAAC
(A, G, C or T) (C or T) CCTG-3'

(SEQ ID NO: 1)

5 5'-GT (A, G, C or T) G (A or T) (A or G) (A or G) GGCA
(A, G, C or T) CCAGCAGA (G or T) GGCAAA-3'

(SEQ ID NO: 2)

10 The parentheses indicate the incorporation of a plurality of bases, leading to multiple oligonucleotides in the primer preparation. In other words, nucleotide residues in parentheses of the aforementioned DNAs were incorporated in the presence of a mixture of plural bases at the time of synthesis.

Example 2

15 Isolation of Human Somatostatin Receptor Protein-Encoding DNA,
Human D5 Dopamine Receptor Protein-Encoding DNA, and Rat
Somatostatin Receptor Protein-Encoding DNA

(1) Amplification of DNA by Polymerase Chain Reaction (PCR)
cDNAs (QuickClone, CLONTECH Laboratories, Inc.) prepared from human brain amygdaloid nucleus, human pituitary gland and rat brain each in an amount of 1 ng as templates, the synthetic DNA primers prepared in Example 1 each in an amount of 1 μ M, 2.5 mM dNTPs (deoxyribonucleoside triphosphates), and 2.5 units of Taq DNA polymerase (Takara Shuzo Co., Japan) were mixed together with a buffer attached to the enzyme kit
25 such that the total amount was 100 μ l. The polymerase chain reaction was carried out by using a Thermal Cycler manufactured by Perkin-Elmer Co. One cycle was set to include 96 °C for 30 sec., 45 °C for 1 min. and 60 °C for 3 min.. Totally this one cycle was repeated 30 times to amplify DNAs.
30 Amplification of DNAs was confirmed by 1.2% agarose electrophoresis [Figure 17].

(2) Isolation of Amplified DNA and Analysis of DNA Sequence

By using a TA Cloning Kit (Invitrogen Co.), the DNA amplified by the PCR was inserted into a plasmid vector, pCRTM II. The DNA was transfected into E. coli attached to the
5 kit to form an amplified DNA library. Colonies formed by the transformants were selected under guidance based on the activity of β -galactosidase on X-gal (5-bromo-4-chloro-3-indolyl- β -D-galactoside)-added LB (Luria-Bertani) plates in order to separate only white colonies in which DNA fragments
10 are inserted. They were cultured in an LB culture medium to which ampicillin was added and plasmid DNAs were prepared with an automatic plasmid extracting machine (Kurabo Co., Japan).

An aliquot of the DNA thus prepared was further digested with EcoRI to confirm DNA fragments that were
15 inserted, and a DNA yield each of clones was compared with a marker. An aliquot of the plasmid DNA thus prepared was treated with RNase, extracted with phenol/chloroform, precipitated in ethanol, and the resulting product was then reacted for sequencing by using a DyeDeoxy terminator cycle
20 sequencing kit (Applied Biosystems Co.).

Sequencing was carried out by using a 370A fluorescent automatic sequencer manufactured by Applied Biosystems Co. The nucleotide sequences obtained were analyzed by using DNASIS (Hitachi Software Engineering, Japan).
25 The nucleotide sequences obtained are shown in Figures 18, 19, 20 and 21. From these Figures and the results of homology retrieval, it was learned that the DNAs obtained were DNAs encoding human somatostatin receptor protein [Figures 18 and 19], human D5 dopamine receptor protein [Figure 20] and rat
30 somatostatin receptor protein [Figure 21] that can be classified each into a group of G protein coupled receptor proteins.

In Figure 18 as described herein, the nucleotide sequence of the DNA is in agreement with the nucleotide
35 sequence encoding somatostatin receptor (HUMSOMAT) and the clone, A58, is a human somatostatin receptor cDNA. The underlined part represents the 5' side synthetic DNA primer

used for the PCR. Thus, even when parts of the nucleotide sequence are mismatched, amplification is effected to a sufficient degree by the PCR.

It will be understood from Figure 19 that the clone, A58 is in good agreement with the nucleotide sequence coding for the human somatostatin receptor (HUMSOMAT) even when the sequencing is carried out from the opposite side. The underlined part represents the 3' side synthetic DNA primer used for the PCR. In this figure, the nucleotide sequences are mismatched to some extent even in the portions other than the primer portion presumably due to base substitution at the time of PCR or due to partial deviation in the sequencing reaction. It can be confirmed via sequencing of chains complementary thereto as required.

In Figure 20 as described herein, the nucleotide sequence of the DNA is in good agreement with a nucleotide sequence coding for the human D5 dopamine receptor (HUMDRD5A) except the primer portion (underlined). It was learned that the clone, 57-A-2, is a human D5 dopamine receptor cDNA.

In Figure 21 as described herein, the DNA is in good agreement with a nucleotide sequence coding for the rat somatostatin receptor (RNU04738) except the primer portion (underlined). It was learned that the clone, B54, is a rat somatostatin receptor cDNA.

25

Example 3

Isolation of Human Pituitary Gland-Derived G Protein Coupled Receptor Protein-Encoding DNA

(1) Amplification of Receptor cDNA by PCR Using Human Pituitary Gland-Derived cDNA

30

By using human pituitary gland-derived cDNA (QuickClone, CLONTECH Laboratories, Inc.) as a template, PCR amplification using the DNA primers synthesized in Example 1 was carried out. The composition of the reaction solution consisted of the synthetic DNA primers (SEQ: 5' primer sequence and 3' primer sequence) each in an amount of 1 μ M, 1 ng of the template cDNA, 0.25 mM dNTPs, 1 μ l of Taq DNA polymerase and

35

a buffer attached to the enzyme kit, and the total amount of the reaction solution was made to be 100 μ l. The cycle for amplification including 95 °C for 1 min., 55 °C for 1 min. and 72 °C for 1 min. was repeated 30 times by using a Thermal Cycler (Perkin-Elmer Co.). Prior to adding Taq DNA polymerase, the remaining reaction solution was mixed and was heated at 95 °C for 5 minutes and at 65 °C for 5 minutes. The amplified products were confirmed relying upon 1.2% agarose gel electrophoresis and ethidium bromide staining.

10 (2) Subcloning of PCR Product into Plasmid Vector and
Selection of Novel Receptor Candidate Clone via Decoding
Nucleotide Sequence of Inserted cDNA Region

The PCR products were separated by using a 0.8% low-melting temperature agarose gel, the band parts were excised from the gel with a razor blade, and were heat-melted, extracted with phenol and precipitated in ethanol to recover DNAs. According to the protocol attached to a TA Cloning Kit (Invitrogen Co.), the recovered DNAs were subcloned into the plasmid vector, pCRTM II (TM represents registered trademark). The recombinant vectors were introduced into E. coli INV α F' competent cells (Invitrogen Co.) to produce transformants. Then, transformant clones having a cDNA-inserted fragment were selected in an LB agar culture medium containing ampicillin and X-gal. Only transformant clones exhibiting white color were picked with a sterilized toothstick to obtain transformant Escherichia coli INV α F'/p19P2.

The individual clones were cultured overnight in an LB culture medium containing ampicillin and treated with an automatic plasmid extracting machine (Kurabo Co., Japan) to prepare plasmid DNAs. An aliquot of the DNA thus prepared was cut by EcoRI to confirm the size of the cDNA fragment that was inserted. An aliquot of the remaining DNA was further processed with RNase, extracted with phenol/chloroform, and precipitated in ethanol so as to be condensed. Sequencing was carried out by using a DyeDeoxy terminator cycle sequencing kit (ABI Co.), the DNAs were decoded by using a

fluorescent automatic sequencer, and the data of the nucleotide sequences obtained were read by using DNASIS (Hitachi System Engineering Co., Japan). The underlined portions represent regions corresponding to the synthetic primers [Figures 22 and 23].

Homology retrieval was carried out based upon the determined nucleotide sequences [Figures 22 and 23]. As a result, it was learned that a novel G protein coupled receptor protein was encoded by the cDNA fragment insert in the plasmid, p19P2, possessed by the transformant Escherichia coli INV α F'/p19P2. To further confirm this fact, by using DNASIS (Hitachi System Engineering Co., Japan) the nucleotide sequences were converted into amino acid sequences [Figures 22 and 23], and homology retrieval was carried out in view of hydrophobicity plotting [Figures 24 and 25] and at the amino acid sequence level to find homology relative to neuropeptide Y receptor proteins [Figure 26].

Example 4

Isolation of Mouse Pancreas-Derived G Protein Coupled Receptor Protein-Encoding DNA

(1) Preparation of Poly(A)⁺ RNA Fraction from Mouse Pancreatic β -Cell Strain, MIN6 and Synthesis of cDNA

A total RNA was prepared from the mouse pancreatic β -cell strain, MIN6 (Jun-ichi Miyazaki et al., Endocrinology, Vol. 127, No. 1, p.126-132) according to the guanidine thiocyanate method (Kaplan B.B. et al., Biochem. J., 183, 181-184 (1979)) and, then, poly(A)⁺ RNA fractions were prepared with a mRNA purifying kit (Pharmacia Co.). Next, to 5 μ g of the poly(A)⁺ RNA fraction was added a random DNA hexamer (BRL Co.) as a primer, and the resulting mixture was subjected to reaction with mouse Moloney Leukemia virus (MMLV) reverse transcriptase (BRL Co.) in the buffer attached to the MMLV reverse transcriptase kit to synthesize complementary DNAs. The reaction product was extracted with phenol/chloroform (1:1), precipitated in ethanol, and was then dissolved in 30 μ l of TE buffer (10 mM Tris-HCl at pH8.0, 1 mM EDTA at pH8.0).

- 1 9 6 -

(2) Amplification of Receptor cDNA by PCR Using MIN6-Derived cDNA and Sequencing

By using, as a template, 5 μ l of cDNA prepared from the mouse pancreatic β -cell strain, MIN6 in the above step (1), PCR amplification using the DNA primers synthesized in Example 1 was carried out under the same conditions as in Example 3(2). The resulting PCR product was subcloned into the plasmid vector, pCRTM II, in the same manner as in Example 2 to obtain a plasmid, pG3-2. The plasmid pG3-2 was transfected into E. coli INV α F' to obtain transformed Escherichia coli INV α F'/pG3-2.

By using, as a template, 5 μ l of the cDNA prepared from the mouse pancreatic β -cell strain, MIN6, PCR amplification using DNA primers as disclosed in Libert F. et al., "Science, 244:569-572, 1989", i.e., a degenerate synthetic primer represented by the following sequence:

5'-CTGTG (C or T) G (C or T) (G or C) AT (C or T) GCIIT
(G or T) GA (C or T) (A or C) G (G or C) TAC-3'

(SEQ ID NO: 60)

wherein I is inosine; and
a degenerate synthetic primer represented by the following sequence:

5'-A (G or T) G (A or T) AG (A or T) AGGGCAGCCAGCAGAI
(G or C) (A or G) (C or T) GAA-3'

(SEQ ID NO: 61)

wherein I is inosine,
was carried out under the same conditions as in Working Example 1. The resulting PCR product was subcloned into the plasmid vector, pCRTM II, in the same manner as described in Example 3(2) to obtain a plasmid, pG1-10.

The reaction for determining the nucleotide sequence (sequencing) was carried out with a DyeDeoxy terminator cycle sequencing kit (ABI Co.), the DNA was decoded with the fluorescent automatic sequencer (ABI Co.), and the data

of the nucleotide sequence obtained were analyzed with DNASIS (Hitachi System Engineering Co., Japan).

Figure 27 shows a mouse pancreatic β -cell strain MIN6-derived G protein coupled receptor protein-encoding DNA and an amino acid sequence encoded by the isolated DNA based upon the nucleotide sequences of plasmids pG3-2 and pG1-10 which are held by the transformant Escherichia coli INV α F'/pG3-2. The underlined portions represent regions corresponding to the synthetic primers.

Homology retrieval was carried out based upon the determined nucleotide sequence [Figure 27]. As a result, it was learned that a novel G protein coupled receptor protein was encoded by the cDNA fragment obtained. To further confirm this fact, by using DNASIS (Hitachi System Engineering Co., Japan) the nucleotide sequence was converted into an amino acid sequence [Figure 27], hydrophobicity plotting was carried out to confirm the presence of six hydrophobic regions [Figure 28]. Upon comparing the amino acid sequence with that of p19P2 obtained in Example 3, furthermore, a high degree of homology was found as shown in [Figure 61]. As a result, it is strongly suggested that the G protein coupled receptor proteins encoded by pG3-2 and pG1-10 recognize the same ligand as the G protein coupled receptor protein encoded by p19P2 does while the animal species from which the receptor proteins encoded by pG3-2 and pG1-10 are derived is different from that from which the receptor protein encoded by p19P2 is.

Example 5

Isolation of Human Amygdaloid Nucleus-Derived G Protein Coupled Receptor Protein-Encoding DNA

(1) Amplification of Receptor cDNA by PCR Using Human Amygdaloid Nucleus-Derived cDNA

By using an amplified human amygdala-derived cDNA (QuickClone, CLONTECH Laboratories, Inc.) as a template, PCR amplification using the DNA primers synthesized in Example 1 was carried out. The composition of the reaction solution

consisted of the synthetic DNA primers (SEQ: 5' primer sequence and 3' primer sequence) each in an amount of $1\mu\text{M}$, 1 ng of the template cDNA, 0.25 mM dNTPs, $1\mu\text{l}$ of Taq DNA polymerase and a buffer attached to the enzyme kit, and the total amount of the reaction solution was made to be $100\mu\text{l}$. The cycle for amplification including 95°C for 1 min., 55°C for 1 min. and 72°C for 1 min. was repeated 30 times by using a Thermal Cycler (Perkin-Elmer Co.). Prior to adding Taq DNA polymerase, the remaining reaction solution was mixed and was heated at 95°C for 5 minutes and at 65°C for 5 minutes. The amplified products were confirmed relying upon 1.2% agarose gel electrophoresis and ethidium bromide staining.

(2) Subcloning of PCR Product into Plasmid Vector and Selection of Novel Receptor Candidate Clone via Decoding Nucleotide Sequence of Inserted cDNA Region

The PCR products were separated by using a 0.8% low-melting temperature agarose gel, the band parts were excised from the gel with a razor blade, and were heat-melted, extracted with phenol and precipitated in ethanol to recover DNAs. According to the protocol attached to a TA Cloning Kit (Invitrogen Co.), the recovered DNAs were subcloned to the plasmid vector, PCRTM II. The recombinant vectors were introduced into E. coli INV α F' competent cells (Invitrogen Co.) to produce transformants. Then, transformant clones having a cDNA-inserted fragment were selected in an LB agar culture medium containing ampicillin and X-gal. Only transformant clones exhibiting white color were picked with a sterilized toothstick to obtain transformant Escherichia coli INV α F'/p63A2.

The individual clones were cultured overnight in an LB culture medium containing ampicillin and treated with an automatic plasmid extracting machine (Kurabo Co., Japan) to prepare plasmid DNAs. An aliquot of the DNA thus prepared was cut by EcoRI to confirm the size of the cDNA fragment that was inserted. An aliquot of the remaining DNA was further processed with RNase, extracted with phenol/chloroform, and

precipitated in ethanol so as to be condensed. Sequencing was carried out by using a DyeDeoxy terminator cycle sequencing kit (ABI Co.), the DNAs were decoded by using a fluorescent automatic sequencer, and the data of the nucleotide sequences obtained were read by using DNASIS (Hitachi System Engineering Co., Japan).

Homology retrieval was carried out based upon the determined nucleotide sequences [Figures 29 and 30]. As a result, it was learned that a novel G protein coupled receptor protein was encoded by the cDNA fragment insert in the plasmid, p63A2 possessed by the transformant Escherichia coli INV α F'/p63A2. To further confirm this fact, by using DNASIS (Hitachi System Engineering Co., Japan) the nucleotide sequences were converted into amino acid sequences [Figures 29 and 30], and homology retrieval was carried out in view of hydrophobicity plotting [Figures 31 and 32] and at the amino acid sequence level to find homology relative to mouse GIR [Figure 33].

Example 6

Cloning of Human Pituitary Gland-Derived G Protein Coupled Receptor Protein cDNA

(1) Cloning of cDNA Comprising Whole Coding Regions for Receptor Protein from Human Pituitary Gland-Derived cDNA Library

The DNA library constructed by Clontech Co. wherein λ gt11 phage vector is used (CLONTECH Laboratories, Inc.; CLH L1139b) was employed as a human pituitary gland-derived cDNA library. The human pituitary gland cDNA library (2×10^6 pfu (plaque forming units)) was mixed with E. coli Y1090⁻ treated with magnesium sulfate, and incubated at 37°C for 15 minutes followed by addition of 0.5% agarose (Pharmacia Co.) LB. The E. coli was plated onto a 1.5% agar (Wako-Junyaku Co.) LB plate (containing 50 μ g/ml of ampicillin). A nitrocellulose filter was placed on the plate on which plaques were formed and the plaque was transferred onto the filter. The filter was denatured with an alkali and

- 2 0 0 -

then heated at 80 °C for 3 hours to fix DNAs.

The filter was incubated overnight at 42 °C together with the probe mentioned herein below in a buffer containing 50% formamide, 5 x SSPE (20 x SSPE (pH 7.4) is 3 M NaCl, 0.2 M $\text{NaH}_2\text{PO}_4 \cdot \text{H}_2\text{O}$, 25 mM EDTA), 5 x Denhardt's solution (Nippon Gene, Japan), 0.1% SDS and 100 μ g/ml of salmon sperm DNA for hybridization.

The probe used was obtained by cutting the DNA fragment inserted in the plasmid, p19P2, obtained in Working Example 3, with EcoRI, followed by recovery and labelling by incorporation of [32 P]dCTP (Dupont Co.) with a random prime DNA labelling kit (Amasham Co.).

It was washed with 2 x SSC (20 x SSC is 3 M NaCl, 0.3 M sodium citrate), 0.1% SDS at 55 °C for 1 hour and, then, subjected to an autoradiography at -80 °C to detect hybridized plaques.

In this screening, hybridization signals were recognized in three independent plaques. Each DNA was prepared from the three clones. The DNAs digested with EcoRI were subjected to an agarose electrophoresis and were analyzed by the southern blotting using the same probe as the one used in the screening. Hybridizing bands were identified at about 0.7kb, 0.8 kb and 2.0kb, respectively. Among them, the DNA fragment corresponding to the band at about 2.0kb (λ hGR3) was selected. The λ hGR3-derived EcoRI fragment with a hybridizable size was subcloned to the EcoRI site of the plasmid, pUC18, and *E. coli* JM109 was transformed with the plasmid to obtain transformant *E. coli* JM109/phGR3. A restriction enzyme map of the plasmid, phGR3, was prepared relying upon a restriction enzyme map deduced from the nucleotide sequence as shown in Example 3. As a result, it was learned that it carried a full-length receptor protein-encoding DNA which was predicted from the receptor protein-encoding DNA as shown in Example 3.

(2) Sequencing of Human Pituitary Gland-Derived Receptor Protein cDNA.

Among the EcoRI fragments inserted in the plasmid, phGR3, obtained in the above step (1), the from EcoRI to NheI
5 nucleotide sequence with about 1330bp that is considered to be a receptor protein-coding region was sequenced. Concretely speaking, by utilizing restriction enzyme sites that exist in the EcoRI fragments, unnecessary parts were removed or
10 necessary fragments were subcloned in order to prepare template plasmids for analyzing the nucleotide sequence.

The reaction for determining the nucleotide sequence (sequencing) was carried out with a DyeDeoxy terminator cycle sequencing kit (ABI Co.), the DNA was decoded with the fluorescent automatic sequencer (ABI Co.), and the data of
15 the nucleotide sequence obtained were analyzed with DNASIS (Hitachi System Engineering Co., Japan).

Figure 34 shows a nucleotide sequence of from immediate after the EcoRI site up to the NheI site encoded by phGR3. The nucleotide sequence of the human pituitary gland-
20 derived receptor protein-encoding DNA corresponds to the nucleotide sequence of from 118th to 123rd nucleotides [Figure 34]. An amino acid sequence of the receptor protein that is encoded by the nucleotide sequence is shown in Figure 34. Figure 36 shows the results of hydrophobicity
25 plotting based upon the amino acid sequence.

(3) Northern Hybridization with Human Pituitary Gland-Derived Receptor Protein-Encoding phGR3

Northern blotting was carried out in order to detect the expression of phGR3-encoded human pituitary gland-derived
30 receptor proteins in the pituitary gland at a mRNA level. Human pituitary gland mRNA (2.5 μ g, Clontech Co.) was used as a template mRNA and the same as the probe used in Working Example 5 was used as a probe. Nylon membrane (Pall Biodyne, U.S.A.) was used as a filter for northern blotting and
35 migration of the mRNA and adsorption (sucking) thereof with the blotting filter was carried out according to the method as

disclosed in Molecular Cloning, Cold Spring Harbor Laboratory Press, 1989.

The hybridization was effected by incubating the above-mentioned filter and probe in a buffer containing 50% formamide, 5 x SSPE, 5 X Denhardt's solution, 0.1% SDS and 100 μ g/ml of salmon sperm DNA overnight at 42 °C. The filter was washed with 0.1 x SSC, 0.1% SDS at 50 °C and, after drying with an air, was exposed to an X-ray film (XAR5, Kodak) for three days at -80 °C. The results were as shown in Figure 35 from which it is considered that the receptor gene encoded by phGR3 is expressed in the human pituitary gland.

Example 7

Cloning of Mouse Pancreatic β -Cell Strain, MIN6-Derived G Protein Coupled Receptor Protein cDNA

(1) Preparation of Poly(A)⁺ RNA Fraction from Mouse Pancreatic β -Cell Strain, MIN6 and Synthesis of cDNA

A total RNA was prepared from the mouse pancreatic β -cell strain, MIN6 (Jun-ichi Miyazaki et al., Endocrinology, Vol. 127, No. 1, p.126-132) according to the guanidine thiocyanate method (Kaplan B.B. et al., Biochem. J., 183, 181-184 (1979)) and, then, poly(A)⁺ RNA fractions were prepared with a mRNA purifying kit (Pharmacia Co.). Next, to 5 μ g of the poly(A)⁺ RNA fraction was added a random DNA hexamer (BRL Co.) as a primer, and the resulting mixture was subjected to reaction with MMLV reverse transcriptase (BRL Co.) in the buffer attached to the MMLV reverse transcriptase kit to synthesize complementary DNAs. The reaction product was extracted with phenol/chloroform (1:1), precipitated in ethanol, and was then dissolved in 30 μ l of TE.

(2) Amplification of Receptor cDNA by PCR Using MIN6-Derived cDNA and Sequencing

By using, as a template, 5 μ l of cDNA prepared from the mouse pancreatic β -cell strain, MIN6 in the above step (1), PCR amplification using the DNA primers synthesized in Example 1 was carried out. A reaction solution was

composed of the synthetic DNA primers (SEQ: 5' primer sequence and 3' primer sequence) each in an amount of 100 pM, 0.25 mM dNTPs, 1 μ l of Taq DNA polymerase and 10 μ l of 10 \times buffer attached to the enzyme kit, and the total amount of the reaction solution was made to be 100 μ l. The cycle for amplification including 96 °C for 30 sec., 45 °C for 1 min. and 60 °C for 3 min. was repeated 30 times by using a Thermal Cycler (Perkin-Elmer Co.). Prior to adding Taq DNA polymerase, the remaining reaction solution was mixed and was heated at 95 °C for 5 minutes and at 65 °C for 5 minutes. The amplified products were confirmed relying upon 1.2% agarose gel electrophoresis and ethidium bromide staining.

(3) Subcloning of PCR Product into Plasmid Vector and Selection of Novel Receptor Candidate Clone via Decoding Nucleotide Sequence of Inserted cDNA Region

The PCR products obtained in the above step (2) were separated by using a 0.8% low-melting temperature agarose gel, the band parts were excised from the gel with a razor blade, and were heat-melted, extracted with phenol and precipitated in ethanol to recover DNAs. According to the protocol attached to a TA Cloning Kit (Invitrogen Co.), the recovered DNAs were subcloned to the plasmid vector, PCRTM II. The recombinant vectors were introduced into E. coli JM109 competent cells (Takara Shuzo Co., Japan) to produce transformants. Then, transformant clones having a cDNA-inserted fragment were selected in an LB agar culture medium containing ampicillin, IPTG (isopropylthio- β - D-galactoside) and X-gal. Only transformant clones exhibiting white color were picked with a sterilized toothstick to obtain transformant Escherichia coli JM109/p3H2-17.

The individual clones were cultured overnight in an LB culture medium containing ampicillin and treated with an automatic plasmid extracting machine (Kurabo Co., Japan) to prepare plasmid DNAs. An aliquot of the DNAs thus prepared was cut by EcoRI to confirm the size of the cDNA fragment that was inserted. An aliquot of the remaining DNAs was

further processed with RNase, extracted with phenol/chloroform, and precipitated in ethanol so as to be condensed. Sequencing was carried out by using a DyeDeoxy terminator cycle sequencing kit (ABI Co.), the DNAs were decoded by using a
5 fluorescent automatic sequencer, and the data of the nucleotide sequences obtained were read by using DNASIS (Hitachi System Engineering Co., Japan).

Homology retrieval was carried out based upon the determined nucleotide sequence [Figure 37]. As a result, it
10 was learned that a novel G protein coupled receptor protein was encoded by the cDNA fragment insert in the plasmid possessed by the transformant Escherichia coli JM109/p3H2-17. To further confirm this fact, by using DNASIS (Hitachi System Engineering Co., Japan) the nucleotide sequence were converted
15 into an amino acid sequence [Figure 37], and homology retrieval was carried out in view of hydrophobicity plotting [Figure 38] and at the amino acid sequence level to find homology relative to chicken ATP receptor (P34996), human somatostatin receptor subtype 3 (A46226), human somatostatin receptor subtype 4
20 (JN0605) and bovine neuropeptide Y receptor (S28787) [Figure 39]. Abbreviations in parentheses are reference numbers assigned when they are registered as data to NBRF-PIR/Swiss-PROT and are usually called "Accession Numbers".

Example 8

25 Cloning of Mouse Pancreatic β -Cell Strain, MIN6-Derived G Protein Coupled Receptor Protein cDNA

(1) Preparation of Poly(A)⁺ RNA Fraction from Mouse Pancreatic β -Cell Strain, MIN6 and Synthesis of cDNA

A total RNA was prepared from the mouse pancreatic
30 β -cell strain, MIN6 (Jun-ichi Miyazaki et al., Endocrinology, Vol. 127, No. 1, p.126-132) according to the guanidine thiocyanate method (Kaplan B.B. et al., Biochem. J., 183, 181-184 (1979)) and, then, poly(A)⁺ RNA fractions were prepared with a mRNA purifying kit (Pharmacia Co.). Next, to 5 μ g
35 of the poly(A)⁺ RNA fraction was added a random DNA hexamer (BRL Co.) as a primer, and the resulting mixture was subjected

to reaction with MMLV reverse transcriptase (BRL Co.)
in the buffer attached to the MMLV reverse transcriptase kit
to synthesize complementary DNAs. The reaction product
was extracted with phenol/chloroform (1:1), precipitated in
5 ethanol, and was then dissolved in 30 μ l of TE.

(2) Amplification of Receptor cDNA by PCR Using MIN6-Derived
cDNA and Sequencing

By using, as a template, 5 μ l of cDNA prepared
from the mouse pancreatic β -cell strain, MIN6, in the above
10 step (1), PCR amplification using the DNA primers synthesized
in Example 1 was carried out. A reaction solution was
composed of the synthetic DNA primers (SEQ: 5' primer sequence
and 3' primer sequence) each in an amount of 100 pM,
0.25 mM dNTPs, 1 μ l of Taq DNA polymerase and 10 μ l of 10 \times
15 buffer attached to the enzyme kit, and the total amount of the
reaction solution was made to be 100 μ l. The cycle for
amplification including 96 °C for 30 sec., 45 °C for 1 min.
and 60 °C for 3 min. was repeated 30 times by using a Thermal
Cycler (Perkin-Elmer Co.). Prior to adding Taq DNA
20 polymerase, the remaining reaction solution was mixed and was
heated at 95 °C for 5 minutes and at 65 °C for 5 minutes.
The amplified products were confirmed relying upon 1.2%
agarose gel electrophoresis and ethidium bromide staining.

(3) Subcloning of PCR Product into Plasmid Vector and
25 Selection of Novel Receptor Candidate Clone via Decoding
Nucleotide Sequence of Inserted cDNA Region

The PCR products obtained in the above step (2)
were separated with a 0.8% low-melting temperature agarose
gel, the band parts were excised from the gel with a razor
30 blade, and were heat-melted, extracted with phenol and
precipitated in ethanol to recover DNAs. According to the
protocol attached to a TA Cloning Kit (Invitrogen Co.),
the recovered DNAs were subcloned to the plasmid vector,
pCRTM II. The recombinant vectors were introduced into
35 E. coli JM109 competent cells (Takara Shuzo Co., Japan)

to produce transformants. Then, transformant clones having a cDNA-inserted fragment were selected in an LB agar culture medium containing ampicillin, IPTG and X-gal. Only transformant clones exhibiting white color were picked with a sterilized toothstick to obtain transformant Escherichia coli JM109/p3H2-34.

The individual clones were cultured overnight in an LB culture medium containing ampicillin and treated with an automatic plasmid extracting machine (Kurabo Co., Japan) to prepare plasmid DNAs. An aliquot of the DNAs thus prepared was cut by EcoRI to confirm the size of the cDNA fragment that was inserted. An aliquot of the remaining DNAs was further processed with RNase, extracted with phenol/chloroform, and precipitated in ethanol so as to be condensed. Sequencing was carried out by using a DyeDeoxy terminator cycle sequencing kit (ABI Co.), the DNAs were decoded by using a fluorescent automatic sequencer, and the data of the nucleotide sequences obtained were read by using DNASIS (Hitachi System Engineering Co., Japan).

Homology retrieval was carried out based upon the determined nucleotide sequence [Figure 40]. As a result, it was learned that a novel G protein coupled receptor protein was encoded by the cDNA fragment insert in the plasmid possessed by the transformant Escherichia coli JM109/p3H2-34. To further confirm this fact, by using DNASIS (Hitachi System Engineering Co., Japan) the nucleotide sequence were converted into an amino acid sequence [Figure 40], and homology retrieval was carried out in view of hydrophobicity plotting [Figure 41] and at the amino acid sequence level to find homology relative to human somatostatin receptor subtype 2 (B41795) and rat-derived ligand unknown receptor (A39297) [Figure 42]. Abbreviations in parentheses are reference numbers assigned when they are registered as data to NBRF-PIR/Swiss-PROT and are usually called "Accession Numbers" or "Entry Names".

Example 9Cloning of Rabbit Gastropyrolic Part Smooth Muscle-Derived G
Protein Coupled Receptor Protein cDNA(1) Preparation of Poly(A)⁺ RNA Fraction from Rabbit

5 Gastropyrolic Part Smooth Muscle and Synthesis of cDNA

A total RNA was prepared from rabbit gastropyrolic part smooth muscles by the guanidine thiocyanate method (Kaplan B.B. et al., Biochem. J. 183, 181-184 (1979)) and, then, poly(A)⁺ RNA fractions were prepared with a mRNA purifying kit (Pharmacia Co.). Next, to 5 μ g of the poly(A)⁺ RNA fraction was added a random DNA hexamer (BRL Co.) as a primer, and the resulting mixture was subjected to reaction with MMLV reverse transcriptase (BRL Co.) in the buffer attached to the MMLV reverse transcriptase kit to synthesize complementary DNAs. The reaction product was extracted with phenol/chloroform (1:1), precipitated in ethanol, and was then dissolved in 30 μ l of TE (Tris-EDTA solution).

20 (2) Amplification of Receptor cDNA by PCR Using Rabbit
Gastropyrolic Part Smooth Muscle-Derived cDNA and
Sequencing

By using, as a template, 1 μ l of cDNA prepared from the rabbit gastropyrolic part smooth muscle in the above step (1), PCR amplification using the DNA primers synthesized in Example 1 was carried out. A reaction solution was composed of the synthetic DNA primers (SEQ: 5' primer sequence and 3' primer sequence) each in an amount of 100 pM, 0.25 mM dNTPs, 1 μ l of Taq DNA polymerase and 10 μ l of buffer attached to the enzyme kit, and the total amount of the reaction solution was made to be 100 μ l. The cycle for amplification including 96 °C for 30 sec., 45 °C for 1 min. and 60 °C for 3 min. was repeated 25 times by using a Thermal Cycler (Perkin-Elmer Co.). The amplified products were confirmed relying upon 1.2% agarose gel electrophoresis and ethidium bromide staining.

(3) Subcloning of PCR Product into Plasmid Vector and
Selection of Novel Receptor Candidate Clone via Decoding
Nucleotide Sequence of Inserted cDNA Region

The PCR products obtained in the above step (2)
5 were separated with a 1.0% low-melting temperature agarose
gel, the band parts were excised from the gel with a razor
blade, and were heat-melted, extracted with phenol and
precipitated in ethanol to recover DNAs. According to the
protocol attached to a TA Cloning Kit (Invitrogen Co.),
10 the recovered DNAs were subcloned to the plasmid vector,
pCRTM II. The recombinant vectors were introduced into
E. coli JM109 competent cells (Takara Shuzo Co., Japan) to
produce transformants. Then, transformant clones having
a cDNA-inserted fragment were selected in an LB agar
15 culture medium containing ampicillin, IPTG and X-gal. Only
transformant clones exhibiting white color were picked with
a sterilized toothstick to obtain transformant Escherichia
coli JM109/pMD4.

The individual clones were cultured overnight in an
20 LB culture medium containing ampicillin and treated with an
automatic plasmid extracting machine (Kurabo Co., Japan) to
prepare plasmid DNAs. An aliquot of the DNAs thus prepared
was cut by EcoRI to confirm the size of the cDNA fragment
that was inserted. An aliquot of the remaining DNAs was
25 further processed with RNase, extracted with phenol/chloroform,
and precipitated in ethanol so as to be condensed. Sequencing
was carried out by using a DyeDeoxy terminator cycle
sequencing kit (ABI Co.), the DNAs were decoded by using a
fluorescent automatic sequencer, and the data of the
30 nucleotide sequences obtained were read by using DNASIS
(Hitachi System Engineering Co., Japan). The determined
nucleotide sequence was as shown in Figure 43. It was learned
from Figure 43 that the cloned cDNA fragment was amplified
from both sides with only the synthetic DNA primer having a
35 nucleotide sequence represented by SEQ ID NO: 1 as synthesized
in Example 1.

Homology retrieval was carried out based upon the

determined nucleotide sequence [Figure 43]. As a result, it was learned that a novel G protein coupled receptor protein was encoded by the cDNA fragment insert in the plasmid possessed by the transformant Escherichia coli JM109/pMD4.

5 To further confirm this fact, by using DNASIS (Hitachi System Engineering Co., Japan) the nucleotide sequence were converted into an amino acid sequence [Figure 43], and homology retrieval was carried out in view of hydrophobicity plotting [Figure 44] and at the amino acid sequence level to find homology relative
10 to rat ligand-unknown receptor protein (A35639) [Figure 45]. Abbreviations in parentheses are reference numbers assigned when they are registered as data to NBRF-PIR/Swiss-PROT and are usually called "Accession Numbers".

Example 10

15 Cloning of cDNA Comprising Whole Coding Regions for Receptor Protein from Mouse Pancreatic β -Cell Strain, MIN6-Derived cDNA Library

(1) Cloning of cDNA Comprising Whole Coding Regions for Receptor Protein from Mouse Pancreatic β -Cell Strain,
20 MIN6-Derived cDNA Library

SuperscriptTM Lambda System (BRL, Cat. 8256) distributed by BRL Co. and GigaPack II Gold (Stratagene, Cat. 200215) distributed by Stratagene Co. were used to construct MIN6-derived cDNA libraries. By using the above
25 kits, a MIN6 cDNA library with 2.2×10^6 pfu (plaque forming units) was constructed from $10 \mu\text{g}$ of MIN6 poly(A)⁺ RNA. The cDNA library was mixed with E. coli Y1090⁻ treated with magnesium sulfate, and incubated at 37 °C for 15 minutes followed by addition of 0.5% agarose (Pharmacia Co.) LB.
30 The E. coli was plated onto a 1.5% agar (Wako-Junyaku Co.) LB plate (containing 50 $\mu\text{g/ml}$ of ampicillin). A nitrocellulose filter was placed on the plate on which plaques were formed and the plaque was transferred onto the filter. The filter was denatured with an alkali and then heated at 80 °C for 3 hours
35 to fix DNAs.

The filter was incubated overnight at 42 °C together

with the probe mentioned herein below in a buffer containing 50% formamide, 5 x SSPE, 5 X Denhardt's solution, 0.1% SDS and 100 μ g/ml of salmon sperm DNA for hybridization.

5 The probe used was obtained by cutting the DNA fragment inserted in the plasmid, p3H2-34, obtained in Working Example 8, with EcoRI, followed by recovery and labeling by incorporation of [32 P]dCTP (Dupont Co.) with a random prime DNA labelling kit (Amasham Co.).

10 It was washed with 2 x SSC (150 mM NaCl and 15 mM sodium citrate), 0.1% SDS at 55 °C for 1 hour and, then, subjected to an autoradiography at -80 °C to detect hybridized plaques.

In this screening, hybridization signals were recognized in two independent plaques. Each DNA was
15 prepared from the two clones. The DNAs digested with SalI and NotI were subjected to an agarose electrophoresis and were analyzed. Inserted fragments were identified at about 2.0kb and 3.0kb, respectively. Between them, the DNA fragment corresponding to the band at about 3.0kb (λ No.20)
20 was selected. The λ No.20-derived NotI-SalI fragment with about 3.0kb was subcloned into the NotI-SalI site of the plasmid, pBluescriptTM II SK(+), and E. coli JM109 was transformed with the plasmid to obtain a transformant E. coli JM109/pMGR20. A restriction enzyme map of the
25 plasmid, pMGR20, was prepared relying upon a restriction enzyme map deduced from the nucleotide sequence as shown in Working Example 8. As a result, it was learned that it carried a full-length receptor protein-encoding DNA which was predicted from the receptor protein-encoding DNA as shown in Working
30 Example 8.

(2) Sequencing of MIN6-Derived Receptor Protein Full-Length cDNA

Among the NotI-SalI fragments inserted in the plasmid, pMGR20, obtained in the above step (1), the nucleotide
35 sequence with total 1607bp, including not only a region that is considered to be a receptor protein-coding region (ORF) but

- 2 1 1 -

also a neighboring region thereof was sequenced. Concretely speaking, by utilizing restriction enzyme sites that exist in the NotI-SalI fragments, unnecessary parts were removed or necessary fragments were subcloned in order to prepare template plasmids for analyzing the nucleotide sequence thereof. As for the nucleotide sequences of part of the regions, primers for sequencing were synthesized based upon the nucleotide sequences that were determined already and used to make confirmation.

5 The reaction for determining the nucleotide sequence (sequencing) was carried out with a DyeDeoxy terminator cycle sequencing kit (ABI Co.), the DNA was decoded with the fluorescent automatic sequencer (ABI Co.), and the data of the nucleotide sequence obtained were analyzed with DNASIS (Hitachi System Engineering Co., Japan).

10 Figure 46 shows a nucleotide sequence around an open reading frame (ORF) of a mouse galanin receptor protein encoded by the cDNA insert in pMGR20. The nucleotide sequence of mouse galanin receptor protein-encoding DNA corresponds to from the 481st to 1525th nucleotides of the nucleotide sequence in Figure 46. The nucleotide sequence converted into an amino acid sequence [Figure 47]. Since the amino acid sequence [Figure 46] has 92% homology to the hydrophobicity plotting was carried out [Figure 47]. Since the amino acid sequence [Figure 46] has 92% homology to the human-derived galanin receptor protein at the amino acid sequence level [Figure 48], it was learned that the cDNA insert in the pMGR20 is a mouse-derived galanin receptor protein-encoding cDNA.

Example 11

30 Preparation of Synthetic DNA Primer for Amplifying G Protein Coupled Receptor Protein-Encoding DNA

Highly homologous parts were found by comparing nucleotide sequences corresponding to or near the third membrane-spanning domain [3C and 3D in Figure 4] and the sixth membrane-spanning domain [6C of Figure 6] among known G protein coupled receptors, i.e., rat-derived angiotensin II

receptor protein (L32840), rat-derived angiotensin Ib receptor protein (X64052), rat-derived angiotensin receptor protein subtype (M90065), human-derived angiotensin Ia receptor protein (M91464), rat-derived cholecystokinin_A receptor protein (M88096), rat-derived cholecystokinin_B receptor protein (M99418), human-derived cholecystokinin_B receptor protein (L04473), mouse-derived low-affinity interleukin 8 receptor protein (M73969), human-derived high-affinity interleukin 8 receptor protein (X65858), mouse-derived C5a anaphylatoxin receptor protein (S46665), human-derived N-formyl peptide receptor protein (M60626), etc.

The aforementioned abbreviations in parentheses are reference numbers that are indicated when the GenBank/EMBL data base is retrieved, and are usually called "Accession Numbers".

It was planned to incorporate mixed bases relying upon the base regions that were in agreement with a large number of receptor protein cDNAs in order to enhance base agreement of sequences with as many receptor cDNAs as possible even in other regions. Based upon these sequences, the degenerate synthetic DNA (3D of Figure 4) having a nucleotide sequence represented by SEQ ID NO: 3 which is complementary to the homologous nucleotide sequence of Figure 4 and the degenerate synthetic DNA (nucleotide sequence complementary to 6C of Figure 6) having a nucleotide sequence represented by SEQ ID NO: 4 were produced. Nucleotide synthesis was carried out by a DNA synthesizer.

[Synthetic DNAs]

5'-CTCGC (G or C) GC (C or T) (A or C) TI (A or G) G
(C or T) ATGGA (C or T) CGITAT-3'
(SEQ ID NO:3)

5'-CATGT (A or G) G (T or A) AGGGAAICCCAG (G or C) A
(A or C) AI (A or G) A (A or G)(A or G) AA-3'
(SEQ ID NO:4)

The parentheses indicate the incorporation of a plurality of bases, leading to multiple oligonucleotides in the primer preparation. In other words, nucleotide residues in parentheses of the aforementioned DNAs were incorporated in the presence of a mixture of plural bases at the time of synthesis, provided that I denotes inosine.

Example 12

Cloning of Rabbit Gastropyrolic Part Smooth Muscle-Derived G Protein Coupled Receptor Protein cDNA

10 (1) Preparation of Poly(A)⁺ RNA Fraction from Rabbit

Gastropyrolic Part Smooth Muscle and Synthesis of cDNA

A total RNA was prepared from rabbit gastropyrolic part smooth muscles by the guanidine thiocyanate method (Kaplan B.B. et al., Biochem. J. 183, 181-184 (1979)) and, then, poly(A)⁺ RNA fractions were prepared with a mRNA purifying kit (Pharmacia Co.). Next, to 5 μ g of the poly(A)⁺ RNA fraction was added a random DNA hexamer (BRL Co.) as a primer, and the resulting mixture was subjected to reaction with MMLV reverse transcriptase (BRL Co.) in the buffer attached to the MMLV reverse transcriptase kit to synthesize complementary DNAs. The reaction product was extracted with phenol/chloroform (1:1), precipitated in ethanol, and was then dissolved in 30 μ l of TE.

25 (2) Amplification of Receptor cDNA by PCR Using Rabbit Gastropyrolic Part Smooth Muscle-Derived cDNA and Sequencing

By using, as a template, 1 μ l of cDNA prepared from the rabbit gastropyrolic part smooth muscle in the above step (1), PCR amplification using the DNA primer having a nucleotide sequence represented by SEQ ID NO: 3 and the DNA primer having a nucleotide sequence represented by SEQ ID NO: 4 synthesized in Example 11 was carried out.

A reaction solution was composed of the synthetic DNA primers (SEQ: 5' primer sequence and 3' primer sequence) each in an amount of 100 pM, 0.25 mM dNTPs, 1 μ l of Taq DNA polymerase

and 10 μ l of buffer attached to the enzyme kit, and the total amount of the reaction solution was made to be 100 μ l. The cycle for amplification including 96 °C for 30 sec., 45 °C for 1 min. and 60 °C for 3 min. was repeated 25 times by using
5 a Thermal Cycler (Perkin-Elmer Co.). The amplified products were confirmed relying upon 1.2% agarose gel electrophoresis and ethidium bromide staining.

(3) Subcloning of PCR Product into Plasmid Vector and
Selection of Novel Receptor Candidate Clone via Decoding
10 Nucleotide Sequence of Inserted cDNA Region

The PCR products obtained in the above step (2) were separated with a 1.0% low-melting temperature agarose gel, the band parts were excised from the gel with a razor blade, and were heat-melted, extracted with phenol and
15 precipitated in ethanol to recover DNAs. According to the protocol attached to a TA Cloning Kit (Invitrogen Co.), the recovered DNAs were subcloned to the plasmid vector, pCRTM II. The recombinant vectors were introduced into E. coli JM109 competent cells (Takara Shuzo Co., Japan) to
20 produce transformants. Then, transformant clones having a cDNA-inserted fragment were selected in an LB agar culture medium containing ampicillin, IPTG and X-gal. Only transformant clones exhibiting white color were picked with a sterilized toothstick to obtain transformant Escherichia
25 coli JM109/pMJ10.

The individual clones were cultured overnight in an LB culture medium containing ampicillin and treated with an automatic plasmid extracting machine (Kurabo Co., Japan) to prepare plasmid DNAs. An aliquot of the DNAs thus prepared
30 was cut by EcoRI to confirm the size of the cDNA fragment that was inserted. An aliquot of the remaining DNAs was further processed with RNase, extracted with phenol/chloroform, and precipitated in ethanol so as to be condensed. Sequencing was carried out by using a DyeDeoxy terminator cycle
35 sequencing kit (ABI Co.), the DNAs were decoded by using a fluorescent automatic sequencer, and the data of the

nucleotide sequences obtained were read by using DNASIS (Hitachi System Engineering Co., Japan). The determined nucleotide sequence was as shown in Figure 49.

Homology retrieval was carried out based upon the determined nucleotide sequence [Figure 49]. As a result, it was learned that a novel G protein coupled receptor protein was encoded by the cDNA fragment insert in the plasmid possessed by the transformant Escherichia coli JM109/pMJ10. To further confirm this fact, by using DNASIS (Hitachi System Engineering Co., Japan) the nucleotide sequence were converted into an amino acid sequence [Figure 49], and homology retrieval was carried out in view of hydrophobicity plotting [Figure 50] and at the amino acid sequence level to find homology relative to human ligand unknown receptor protein (B42009), human N-formyl peptide receptor protein (JC2014), rabbit N-formyl peptide receptor protein (A46520), mouse C5a anaphylatoxin receptor protein (A46525) and bovine neuropeptide Y receptor protein (S28787) [Figure 51]. Abbreviations in parentheses are reference numbers assigned when they are registered as data to NBRF-PIR/Swiss-PROT and are usually called "Accession Numbers".

Example 13

Preparation of Synthetic DNA Primer for Amplifying G Protein Coupled Receptor Protein-Encoding DNA

A comparison of nucleotide sequences coding for regions corresponding to or near the third membrane-spanning domain among known G protein coupled receptors, i.e., mouse-derived κ -opioid receptor protein (L11064), mouse-derived δ -opioid receptor protein (L11065), rat-derived μ -opioid receptor protein (D16349), mouse-derived bradykinin B2 receptor protein (X69676), rat-derived bradykinin B2 receptor protein (M599967), mouse-derived bombesin receptor protein (M35328), human-derived neuromedin B receptor protein (M73482), human-derived gastrin releasing peptide receptor protein (M73481), human-derived bombesin receptor protein subtype 3 (L08893), mouse-derived substance K receptor protein

(X62933), mouse-derived substance P receptor protein (X62934), rat-derived neurokinin 3 receptor protein (J05189), rat-derived endothelin receptor protein (M60786), rat-derived ligand unknown receptor proteins (L04672), (X61496), (X59249) and
5 (L09249), mouse-derived ligand unknown receptor protein (P30731), human-derived ligand unknown receptor proteins (M31210) and (U03642), etc. was made. In particular, the degenerate DNA primer having a nucleotide sequence
10 (3B in Figure 3; SEQ ID NO: 6) with highly common bases (highly homologous nucleotides) was synthesized to enhance base agreement of sequences with as many receptor cDNAs as possible even in other regions on the basis of nucleotide sequence regions that were in agreement with a large number of receptor cDNAs. Nucleotide synthesis was carried out by a DNA
15 synthesizer.

The nucleotide sequence represented by SEQ ID NO: 6 is:

5'-CTGAC (C or T) G (C or T) TCTI (A or G)(G or C) I
(A or G)(C or T) TGAC (A or C) G (A, C or G) TAT-3'

20 The parentheses indicate the incorporation of a plurality of bases, leading to multiple oligonucleotides in the primer preparation. In other words, nucleotide residues in parentheses of the aforementioned DNAs were incorporated in the presence of a mixture of plural bases at the time of synthesis,
25 provided that I denotes inosine.

Furthermore, a comparison of nucleotide sequences coding for regions corresponding to or near the sixth membrane-spanning domain among known G protein coupled receptors, i.e., mouse-derived κ -opioid receptor protein
30 (L11064), mouse-derived δ -opioid receptor protein (L11065), rat-derived μ -opioid receptor protein (D16349), mouse-derived bradykinin B2 receptor protein (X69676), rat-derived bradykinin B2 receptor protein (M59967), mouse-derived bombesin receptor protein (M35328), human-derived neuromedin B receptor protein

(M73482), human-derived gastrin releasing peptide receptor protein (M73481), human-derived bombesin receptor protein subtype 3 (L08893), mouse-derived substance K receptor protein (X62933), mouse-derived substance P receptor protein (X62934),
5 rat-derived neurokinin 3 receptor protein (J05189), rat-derived endothelin receptor protein (M60786), rat-derived ligand unknown receptor proteins (L04672), (X61496), (X59249) and (L09249), mouse-derived ligand unknown receptor protein (P30731), human-derived ligand unknown receptor proteins
10 (M31210) and (U03642), etc. was made. In particular, the degenerate DNA primer having a nucleotide sequence (SEQ ID NO: 8) which is complementary to the nucleotide sequence (6A in Figure 5) with highly common bases (highly homologous nucleotides) was synthesized to enhance base
15 agreement of sequences with as many receptor cDNAs as possible even in other portions on the basis of base regions that are in agreement with a large number of receptor cDNAs.

The nucleotide sequence represented by SEQ ID NO: 8 is:

20 5'-GATGTG (A or G) TA (A or G) GG (G or C)(A or G)
ICCAACAGAIG (A or G) (C or T) AAA-3'

The parentheses indicate the incorporation of a plurality of bases, leading to multiple oligonucleotides in the primer preparation. In other words, nucleotide residues in
25 parentheses of the aforementioned DNAs were incorporated in the presence of a mixture of plural bases at the time of synthesis, provided that I denotes inosine.

The aforementioned abbreviations in parentheses are reference numbers indicated when the GenBank/EMBL data base is
30 retrieved and are usually called "Accession Numbers".

Example 14Cloning of Rabbit Gastropyrolic Part Smooth Muscle-Derived G Protein Coupled Receptor Protein cDNA(1) Preparation of Poly(A)⁺ RNA Fraction from Rabbit

5 Gastropyrolic Part Smooth Muscle and Synthesis of cDNA

A total RNA was prepared from rabbit gastropyrolic part smooth muscles by the guanidine thiocyanate method (Kaplan B.B. et al., Biochem. J. 183, 181-184 (1979)) and, then, poly(A)⁺ RNA fractions were prepared with a mRNA
10 purifying kit (Pharmacia Co.). Next, to 5 μ g of the poly(A)⁺ RNA fraction was added a random DNA hexamer (BRL Co.) as a primer, and the resulting mixture was subjected to reaction with MMLV reverse transcriptase (BRL Co.)
15 in the buffer attached to the MMLV reverse transcriptase kit to synthesize complementary DNAs. The reaction product was extracted with phenol/chloroform (1:1), precipitated in ethanol, and was then dissolved in 30 μ l of TE.

(2) Amplification of Receptor cDNA by PCR Using Rabbit
20 Gastropyrolic Part Smooth Muscle-Derived cDNA and Sequencing

By using, as a template, 1 μ l of cDNA prepared from the rabbit gastropyrolic part smooth muscle in the above step (1), PCR amplification using the DNA primer having a nucleotide sequence represented by SEQ ID NO: 6 and the DNA
25 primer having a nucleotide sequence represented by SEQ ID NO: 8 synthesized in Example 13 was carried out. A reaction solution was composed of the synthetic DNA primers (SEQ: 5' primer sequence and 3' primer sequence) each in an amount of 100 pM, 0.25 mM dNTPs, 1 μ l of Taq DNA polymerase
30 and 10 μ l of buffer attached to the enzyme kit, and the total amount of the reaction solution was made to be 100 μ l. The cycle for amplification including 96 °C for 30 sec., 45 °C for 1 min. and 60 °C for 3 min. was repeated 25 times by using a Thermal Cycler (Perkin-Elmer Co.). The amplified products
35 were confirmed relying upon 1.2% agarose gel electrophoresis and ethidium bromide staining.

(3) Subcloning of PCR Product into Plasmid Vector and
Selection of Novel Receptor Candidate Clone via Decoding
Nucleotide Sequence of Inserted cDNA Region

The PCR products obtained in the above step (2)
5 were separated by using a 1.0% low-melting temperature agarose
gel, the band parts were excised from the gel with a razor
blade, and were heat-melted, extracted with phenol and
precipitated in ethanol to recover DNAs. According to the
protocol attached to a TA Cloning Kit (Invitrogen Co.),
10 the recovered DNAs were subcloned to the plasmid vector,
pCRTM II. The recombinant vectors were introduced into
E. coli JM109 competent cells (Takara Shuzo Co., Japan) to
produce transformants. Then, transformant clones having
a cDNA-inserted fragment were selected in an LB agar
15 culture medium containing ampicillin, IPTG and X-gal. Only
transformant clones exhibiting white color were picked with
a sterilized toothstick to obtain transformant Escherichia
coli JM109/pMH28.

The individual clones were cultured overnight in an
20 LB culture medium containing ampicillin and treated with an
automatic plasmid extracting machine (Kurabo Co., Japan) to
prepare plasmid DNAs. An aliquot of the DNAs thus prepared
was cut by EcoRI to confirm the size of the cDNA fragment
that was inserted. An aliquot of the remaining DNAs was
25 further processed with RNase, extracted with phenol/chloroform,
and precipitated in ethanol so as to be condensed. Sequencing
was carried out by using a DyeDeoxy terminator cycle
sequencing kit (ABI Co.), the DNAs were decoded by using a
fluorescent automatic sequencer, and the data of the
30 nucleotide sequences obtained were read by using DNASIS
(Hitachi System Engineering Co., Japan). The determined
nucleotide sequence was as shown in Figure 52.

Homology retrieval was carried out based upon the
determined nucleotide sequence [Figure 52]. As a result,
35 it was learned that a novel G protein coupled receptor
protein was encoded by the cDNA fragment insert in the
plasmid possessed by the transformant Escherichia coli

JM109/pMH28. To further confirm this fact, by using DNASIS (Hitachi System Engineering Co., Japan) the nucleotide sequence were converted into an amino acid sequence [Figure 52], and homology retrieval was carried out in view of hydrophobicity plotting [Figure 53] and at the amino acid sequence level to find homology relative to mouse IL-8 receptor protein (P35343), human somatostatin receptor protein 1 (A41795) and human somatostatin receptor protein 4 (A47457)[Figure 54]. The aforementioned abbreviations in parentheses are reference numbers assigned when they are registered as data to NBRF-PIR or SWISS-PROT and are usually called "Accession Numbers".

Example 15

Preparation of Synthetic DNA Primer for Amplifying G Protein Coupled Receptor Protein-Encoding DNA

A comparison of nucleotide sequences coding for regions corresponding to or near the second membrane-spanning domain among known G protein coupled receptors, i.e., human-derived galanin receptor (HUMGALAREC), rat-derived α -1B-adrenergic receptor (RATADR1B), human-derived β -1-adrenergic receptor (HUMADRB1), rabbit-derived IL-8 receptor (RABIL8RSB), human-derived opioid receptor (HUMOPIODRE), bovine-derived substance K receptor (BTSKR), human-derived somatostatin receptor-2 (HUMSTRI2A), human-derived somatostatin receptor-3 (HUMSSTR3Y), human-derived gastrin receptor (HUMGARE), human-derived cholecystokinin A receptor (HUMCCKAR), human-derived dopamine receptor-D5 (HUMD1B), human-derived serotonin receptor 5HT1E (HUM5HT1E), human-derived dopamine receptor D4 (HUMD4C), mouse-derived serotonin receptor-2 (MMSERO), rat-derived α -1A-adrenergic receptor (RATADRA1A), rat-derived histamine H2 receptor (S57565), etc. was made. In particular, the degenerate DNA primer having a nucleotide sequence (T2A in Figure 7, SEQ ID NO: 10) with highly common bases (highly homologous nucleotides) was synthesized to enhance base agreement of sequences with as many receptor cDNAs as possible even in other regions on the basis of nucleotide sequence

regions that were in agreement with a large number of receptor cDNAs. Nucleotide synthesis was carried out by a DNA synthesizer.

5 The nucleotide sequence represented by SEQ ID NO: 10 is:
5'-GYCACCAACN₂WSTTCATCCTSWN₂HCTG-3'

wherein S represents G or C; Y represents C or T; W represents A or T; H represents A, C or T and N₂ represents I.

10 The parentheses indicate the incorporation of a plurality of bases, leading to multiple oligonucleotides in the primer preparation. In other words, nucleotide residues in parentheses of the aforementioned DNAs were incorporated in the presence of a mixture of plural bases at the time of synthesis, provided that I denotes inosine.

15 Furthermore, a comparison of nucleotide sequences coding for regions corresponding to or near the seventh membrane-spanning domain among known G protein coupled receptors, i.e., human-derived galanin receptor (HUMGALAREC),
20 rat-derived A1 adenosine receptor (RAT1ADREC), porcine-derived angiotensin receptor (PIGA2R), rat-derived serotonin receptor (RAT5HTRTC), human-derived dopamine receptor (S58541), human-derived gastrin releasing peptide receptor (HUMGRPR), mouse-derived GRP/bombesin receptor (MUSGRPBOM), rat-derived vascular type 1 angiotensin receptor (RRVT1AIIR),
25 human-derived muscarinic acetylcholine receptor (HSHM4), human-derived β -1 adrenergic receptor (HUMDRB1), human-derived gastrin receptor (HUMGARE), rat-derived cholecystokinin receptor (RATCCKAR), rat-derived ligand unknown receptor (S59748), human-derived somatostatin receptor
30 (HUMSST28A), rat-derived ligand unknown receptor (RNGPROCR), mouse-derived somatostatin receptor 1 (MUSSRI1A), human-derived α -A1-adrenergic receptor (HUMA1AADR), mouse-derived delta-opioid receptor (S66181), human-derived somatostatin

receptor-3 (HUMSSTR3Y), etc. was made. In particular, the degenerate DNA primer having a nucleotide sequence (T7A in Figure 8, SEQ ID NO: 11) with highly common bases (highly homologous nucleotides) was synthesized to enhance base agreement of sequences with as many receptor cDNAs as possible even in other regions on the basis of nucleotide sequence regions that were in agreement with a large number of receptor cDNAs. Nucleotide synthesis was carried out by a DNA synthesizer.

10 The nucleotide sequence represented by SEQ ID NO: 11 is:

5'-ASN₂SAN₂RAAGSARTAGAN₂GAN₂RGGRIT-3'

wherein R represents A or G; S represents G or C and N₂ represents I.

15 The parentheses indicate the incorporation of a plurality of bases, leading to multiple oligonucleotides in the primer preparation. In other words, nucleotide residues in parentheses of the aforementioned DNAs were incorporated in the presence of a mixture of plural bases at the time of synthesis, provided that I denotes inosine.

20 The aforementioned abbreviations in parentheses are reference numbers indicated when the GenBank/EMBL data base is retrieved and are usually called "Accession Numbers".

Example 16

25 Cloning of Rabbit Gastropyrolic Part Smooth Muscle-Derived G Protein Coupled Receptor Protein cDNA

(1) Preparation of Poly(A)⁺ RNA Fraction from Rabbit

Gastropyrolic Part Smooth Muscle and Synthesis of cDNA

30 A total RNA was prepared from rabbit gastropyrolic part smooth muscles by the guanidine thiocyanate method (Kaplan B.B. et al., Biochem. J. 183, 181-184 (1979)) and, then, poly(A)⁺ RNA fractions were prepared with a mRNA purifying kit (Pharmacia Co.). Next, to 5 µg of the

poly(A)⁺ RNA fraction was added a random DNA hexamer (BRL Co.) as a primer, and the resulting mixture was subjected to reaction with MMLV reverse transcriptase (BRL Co.) in the buffer attached to the MMLV reverse transcriptase kit to synthesize complementary DNAs. The reaction product was extracted with phenol/chloroform (1:1), precipitated in ethanol, and was then dissolved in 30 μ l of TE.

(2) Amplification of Receptor cDNA by PCR Using Rabbit Gastropyrolic Part Smooth Muscle-Derived cDNA and Sequencing

By using, as a template, 1 μ l of cDNA prepared from the rabbit gastropyrolic part smooth muscle in the above step (1), PCR amplification using the DNA primer having a nucleotide sequence represented by SEQ ID NO: 10 and the DNA primer having a nucleotide sequence represented by SEQ ID NO: 11 synthesized in Example 15 was carried out.

A reaction solution was composed of the synthetic DNA primers (SEQ: 5' primer sequence and 3' primer sequence) each in an amount of 100 pM, 0.25 mM dNTPs, 1 μ l of Taq DNA polymerase and 10 μ l of buffer attached to the enzyme kit, and the total amount of the reaction solution was made to be 100 μ l. The cycle for amplification including 96 °C for 30 sec., 45 °C for 1 min. and 60 °C for 3 min. was repeated 25 times with a Thermal Cycler (Perkin-Elmer Co.). The amplified products were confirmed relying upon 1.2% agarose gel electrophoresis and ethidium bromide staining.

(3) Subcloning of PCR Product into Plasmid Vector and Selection of Novel Receptor Candidate Clone via Decoding Nucleotide Sequence of Inserted cDNA Region

The PCR products obtained in the above step (2) were separated with a 1.4% low-melting temperature agarose gel, the band parts were excised from the gel with a razor blade, and were eluted electrophoretically, extracted with phenol and precipitated in ethanol to recover DNAs. According to the protocol attached to a TA Cloning Kit (Invitrogen Co.),

the recovered DNAs were subcloned to the plasmid vector, PCRTM II. The recombinant vectors were introduced into E. coli JM109 competent cells (Takara Shuzo Co., Japan) to produce transformants. Then, transformant clones having
5 a cDNA-inserted fragment were selected in an LB agar culture medium containing ampicillin, IPTG and X-gal. Only transformant clones exhibiting white color were picked with a sterilized toothstick to obtain 100 transformant clones.

The individual clones were cultured overnight in an
10 LB culture medium containing ampicillin and treated with the automatic plasmid extracting machine PI-100 (Kurabo Co., Japan) to prepare plasmid DNAs. An aliquot of the DNA thus prepared was cut by EcoRI to confirm the size of the cDNA fragment that was inserted. An aliquot of the remaining DNA was
15 further processed with RNase, extracted with phenol/chloroform, and precipitated in ethanol so as to be condensed. Sequencing was carried out by using a DyeDeoxy terminator cycle sequencing kit (ABI Co.), the DNAs were decoded by using a fluorescent automatic sequencer.

20 Homology retrieval was carried out based upon the determined nucleotide sequence by using DNASIS (Hitachi System Engineering Co., Japan). As a result, it was learned that a novel G protein coupled receptor protein was encoded by the cDNA fragment insert in the plasmid possessed by
25 the transformant Escherichia coli JM109/pMN7. Figure 56 and Figure 56 show the nucleotide sequences of the cDNA fragments. To further confirm this fact, by using DNASIS (Hitachi System Engineering Co., Japan), the nucleotide sequences were converted into amino acid sequences [Figure 55] and [Figure 56],
30 and hydrophobicity plotting was carried out [Figure 57]. As a result, the presence of hydrophobic domains which prove that it is a G protein coupled receptor protein were confirmed. Furthermore, homology retrieval was carried out at the amino acid sequence level to find that the DNAs were novel receptor
35 proteins having 27% homology relative to rat-derived β_3 -adrenaline receptor protein (A41679), 29% homology relative to rat-derived serotonin (5-HT6) receptor protein (JN0591),

27% homology relative to dog-derived histamine H₂ receptor protein (A39008), 27% homology relative to human-derived somatostatin receptor (type 4) protein (JN0605), 24% homology relative to human-derived dopamine D₁ receptor protein (S11377), 23% homology relative to rat-derived neurotensin receptor protein (JH0164), 31% homology relative to human-derived cholecystokinin B receptor protein (JC1352), and 30% homology relative to rat-derived gastrin receptor protein (JQ1614). The aforementioned abbreviations in parentheses are reference numbers assigned when they are registered as data to NBRF-PIR and are usually called "Accession Numbers".

Example 17

Amplification of Receptor cDNA by PCR Using MIN6-Derived cDNA and Sequencing

By using, as a template, 5 μ l of cDNA prepared from the mouse pancreatic β -cell strain, MIN6 in Working Example 4 (1), PCR amplification using the DNA primers synthesized in Example 4 (2) as disclosed in Libert F. et al., "Science, 244:569-572, 1989", i.e., a synthetic primer represented by the following sequence:

5'-CTGTG (C or T) G (C or T) (G or C) AT (C or T) GCIIT
(G or T) GA (C or T) (A or C) G (G or C) TAC-3'

(SEQ ID NO: 60)

wherein I is inosine; and
a synthetic primer represented by the following sequence:

5'-A (G or T) G (A or T) AG (A or T) AGGGCAGCCAGCAGAI
(G or C) (A or G) (C or T) GAA-3'

(SEQ ID NO: 61)

wherein I is inosine, was carried out under the same conditions as in Example 3 (1). The resulting PCR product was subcloned to the plasmid vector, pCRTM II, in the same manner as in Example 3 (2) to obtain a plasmid, p5S38. The plasmid p5S38 was transfected into E. coli JM109 to obtain

transformant Escherichia coli JM109/p5S38.

The reaction for determining the nucleotide sequence (sequencing) was carried out with a DyeDeoxy terminator cycle sequencing kit (ABI Co.), the DNA was decoded with
5 the fluorescent automatic sequencer (ABI Co.), and the data of the nucleotide sequence obtained were read with DNASIS (Hitachi System Engineering Co., Japan).

Figure 62 shows a mouse pancreatic β -cell strain MIN6-derived G protein coupled receptor protein-encoding DNA
10 (SEQ ID NO: 33) and an amino acid sequence (SEQ ID NO: 28) encoded by the isolated DNA based upon the nucleotide sequence of plasmid, p5S38. The underlined portions represent regions corresponding to the synthetic primers.

Homology retrieval was carried out based upon the
15 determined nucleotide sequence [Figure 62]. As a result, it was learned that a novel G protein coupled receptor protein was encoded by the cDNA fragment obtained. To further confirm this fact, by using DNASIS (Hitachi System Engineering Co., Japan), the nucleotide sequence was converted into an
20 amino acid sequence [Figure 62], and hydrophobicity plotting was carried out to confirm the presence of four hydrophobic regions [Figure 64]. Upon comparing the amino acid sequence with those encoded by p19P2 obtained in Example 3 (2) and encoded by pG3-2 obtained in Example 4 (2), furthermore,
25 a high degree of homology was found as shown in Figure 63. As a result, it is strongly suggested that the mouse pancreatic β -cell strain, MIN6-derived G protein coupled receptor protein encoded by p5S38 recognizes the same ligand as the human pituitary gland-derived G protein coupled receptor
30 protein encoded by p19P2 does while the animal species from which the receptor protein encoded by p5S38 is derived is different from that from which the receptor protein encoded by p19P2 is. It is also strongly suggested that the mouse pancreatic β -cell strain, MIN6-derived G protein coupled
35 receptor protein encoded by p5S38 recognizes the same ligand as the mouse pancreatic β -cell strain, MIN6-derived G protein coupled receptor proteins encoded by pG3-2 and pG1-10 do and

they are analogous receptor proteins one another (so-called "subtype").

Example 18

Northern Hybridization with cDNA Fragment Included in

MIN6-Derived Receptor Protein-Encoding p3H2-17

Mouse cell line, MIN6, Neuro-2a, poly(A)⁺RNA (2.5 μ g) and mouse brain, spleen, thymus and pancreas poly(A)⁺RNAs (2.5 μ g) were used as poly(A) RNAs. The DNA fragment inserted into the plasmid, p3H2-17, obtained in Example 7 (3)

was recovered as a DNA fragment with about 400bp by cutting the plasmid with EcoRI and the resulting DNA fragment was labeled by incorporation of [³²P]dCTP (Dupont Co.) with a random prime DNA labeling kit (Amasham Co.). The about 400bp labeled DNA fragment was used as a probe for hybridization.

Nylon membrane (PaLL Biodyne, U.S.A.) was used as a filter for northern blotting and migration of the poly(A)⁺RNA and adsorption (sucking) thereof with the blotting filter was carried out according to the method as disclosed in Molecular Cloning, Cold Spring Harbor Laboratory Press, 1989.

The hybridization was carried out by incubating the above-mentioned filter and probe in a buffer containing 50% formamide, 5 x SSPE (20 x SSPE (pH 7.4) is 3 M NaCl, 0.2 M NaH₂PO₄·H₂O, 25 mM EDTA), 5 X Denhardt's solution (Nippon Gene, Japan), 0.1% SDS and 100 μ g/ml of salmon sperm DNA overnight at 42 °C. The filter was washed with 0.1 x SSC (20 x SSC is 3 M NaCl, 0.3 M sodium citrate), 0.1% SDS at 50 °C and, after drying with an air, was exposed to an X-ray film (XAR5, Kodak) for 15 days at -80 °C. The results were as shown in Figure 65.

It is considered from Figure 65 that mRNA for the the receptor gene encoded by the cDNA fragment included in p3H2-17 is expressed in the cell line, MIN6, Neuro-2a, and the mouse brain, pancreas, spleen and thymus and especially expressed in the mouse spleen and thymus intensely. The MIN6 signal position hybridizable in the northern hybridization plotting is different from those of other organ cells.

Example 19PCR Cloning of cDNA Comprising Whole Coding Regions of
Receptor Proteins from Mouse Spleen, Thymus-Derived
Poly(A)⁺ RNA and Sequencing5 (1) PCR Cloning of cDNA Comprising Whole Coding Region of
Receptor Protein

In order to obtain a full-length open reading frame
(coding region) of the receptor protein encoded by the cDNA
fragment included in p3H2-17, PCR amplification was carried out
10 by 5'RACE and 3'RACE wherein poly(A)⁺ RNA derived from mouse
spleen and thymus was used.

Based on the nucleotide sequence of 3H2-17 which was
disclosed, the following 4 primers were synthesized:

(Nucleotide sequence of synthesized primer)

- 15 ① 5'-TAGTGTGTGGAGTCGTGTGGCTGGCTG-3' (SEQ ID NO: 20)
- ② 5'-AGTCTTTGCTGCCACAGGCATCCAGCG-3' (SEQ ID NO: 21)
- ③ 5'-CAAGCCAGTAAGGCTATGAAGGGCAGCAAG-3' (SEQ ID NO: 22)
- 20 ④ 5'-ACAGGACCTGCTGGGCCATCCTGGCGACACA-3' (SEQ ID NO: 23)

The 5'RACE was carried out according to the protocol
of 5'Ampli Finder RACE kit from ClonTech Co. (ClonTech Co.).

25 In an embodiment, cDNA was prepared from 2 μ g each
of poly(A)⁺ RNAs derived from mouse spleen and thymus by using
the aforementioned primer ④ and ligated with an anchor
attached to the 5'RACE kit. A mixture of a 1/200 amount of the
cDNA thus prepared, the anchor and the aforementioned primer
30 ③ was subjected to PCR using 4 polymerases, Taq (Takara, Japan),
Ex Taq (Takara, Japan), Vent (New England Biolabs) and Pfu
(Stratagene) under the following conditions: 96 °C for 30 sec.,
60 °C for 60 sec., 72 °C for 90 sec. and 35 cycles. A 1/5
amount of the PCR product was subjected to agarose
35 electrophoresis and stained with ethidium bromide (EtBr).
The results are shown in Figure 66. The amplified DNA band

appeared at an about 1 kbp position and the isolated about 1 kbp DNA band which was synthesized from poly(A)⁺ RNAs derived from mouse spleen and thymus by the 5'RACE using Ex Taq polymerase was treated with SUPRECTM-01 (Takara, Japan) to recover cDNA.

The isolated DNA was subcloned into pCRTMII vector by using a TA Cloning Kit (Invitrogen Co.) and the vector was transfected into E. coli JM109 to obtain 3 transformant clones, N26, N64 and N75. The clone, N26, holds the thymus-derived cDNA which is amplified by the 5'RACE and the clone, N75, holds the spleen-derived cDNA which is amplified by the 5'RACE (Figure 68).

The 3'RACE was carried out according to the protocol of 3' RACE kit (GIBCO BRL Co.).

In an embodiment, cDNA was prepared from 1 μ g each of poly(A)⁺ RNAs derived from mouse spleen and thymus by using an adaptor primer attached to the 3' RACE kit. A mixture of the adaptor primer thus prepared and a 1/10 amount of cDNA which was prepared by using the aforementioned primer ① was subjected to 1st PCR using 4 polymerases, Taq (Takara, Japan), Ex Taq (Takara, Japan), Vent (NEB) and Pfu (Stratagene) under the following conditions: 96 °C for 30 sec., 55 °C for 60 sec., 72 °C for 120 sec. and 30 cycles. A mixture of a 1/50 amount of the 1st PCR product, the aforementioned primer ② and the adaptor primer was subjected to 2nd PCR using the aforementioned polymerases under the same conditions as aforementioned herein in the 5'RACE process. A 1/5 amount of the 2nd PCR product was subjected to agarose electrophoresis and stained with ethidium bromide. The results are shown in Figure 67.

The amplified DNA band appeared at an about 1 kbp position (which was synthesized from poly(A)⁺ RNAs derived from mouse thymus by the 3'RACE using Vent polymerase) and the amplified DNA band appeared at an about 1 kbp position (which was synthesized from poly(A)⁺ RNAs derived from mouse thymus by the 3'RACE using Pfu polymerase) were treated with

SUPRECTM-01 (Takara, Japan) to recover cDNA, respectively.

The isolated DNAs were treated with T4 polynucleotide kinase (Wako Pure Chemical Co., Japan) to add phosphate to the end thereof and the phosphorylated DNAs were ligated with pUC18 SmaI BAP (Pharmacia) by using DNA Ligation Kit (Takara, Japan) followed by transformation of E. coli JM109 to obtain 3 transformant clones, C2, C13 and C15. The clones, C13 and C15, hold the thymus-derived cDNA which is amplified by the 3'RACE and the clone, C2, holds the thymus-derived cDNA which is amplified by the 3'RACE (Figure 68).

Based on the nucleotide sequences of clones, N26, N64 and N75, which are considered to hold the N-terminal region of the open reading frame (ORF) of the cDNA fragment included in p3H2-17 and the nucleotide sequences of clones, C2, C13 and C15, which are considered to hold the C-terminal region of the open reading frame (ORF) of the cDNA fragment included in p3H2-17, the entire nucleotide sequence coding for the open reading frame and neighboring region of the receptor protein encoded by the cDNA included in p3H2-17 was determined. To be more specific, sequencing was carried out with the primers used in the 5'RACE and 3'RACE or synthetic primers for sequencing by using a DyeDeoxy Terminator Cycle Sequencing Kit (ABI Co.), the nucleotide sequences were decoded by using a fluorescent automatic sequencer. The obtained data of the DNA were analyzed by DNASIS (Hitachi System Engineering Co., Japan).

PCR errors which presumably happen to occur upon PCR have been corrected by a way of thinking that, when nucleotides between two clones which are independently produced by PCR are identical (e.g. those between clones, N75 and N64, are identical) each other, the identical base is considered as correct. The determined nucleotide sequence is shown in Figure 69. The amino acid sequence is deduced based on the determined nucleotide sequence (Figure 69). Hydrophobicity plotting was carried out based on the deduced amino acid sequence (Figure 70). As a result, it was learned that it was a seven transmembrane G protein coupled receptor, as

it is suggested from the cDNA fragment included in p3H2-17.

Homology retrieval at the amino acid level indicates that it is homologous to mouse P_{2U} purinoceptor and chicken P_{2Y} purinoceptor.

5 Further, the clone which are free of an error in the open reading frame (ORF) was selected and used to construct plasmids carrying the full-length ORF of the receptor protein encoded by p3H2-17. In an embodiment, the cDNA fragment held by the clone, N75, was digested with restriction enzymes,
10 DraIII and EcoRI, to obtain cDNA fragments which are the N-terminal region of the receptor protein held by p3H2-17. The C-terminal cDNA fragment encoded by C13 was digested with restriction enzymes, DraIII and EcoRI, to delete 5'-side regions from the DraIII site of the C-terminal and the long
15 fragment was obtained by the digestion of C13 with restriction enzymes, DraIII and EcoRI. The N75-derived N-terminal cDNA DraIII-EcoRI fragment was ligated with the long C13-derived DraIII-EcoRI fragment by using a DNA Ligation Kit (Takara, Japan) and transfected into Escherichia coli JM109 to obtain
20 transformant Escherichia coli JM109/pMAH2-17.

(2) Electrophysiological Measurement of Receptor Encoded by pMAH2-17

The receptor encoded by pMAH2-17 was examined electrophysiologically in Xenopus oocytes. The ORF of the
25 receptor encoded by pMAH2-17 was inserted into the XhoI-XbaI sites of pBluescriptTM II SK(+) (Stratagene) with directing the sequence thereof downstream from T7 promoter. The resulting plasmid as a template was treated with a mCAPTM mRNA Capping kit (Stratagene) to produce cRNA of this receptor
30 gene.

The cRNA was injected into Xenopus oocytes (50ng cRNA/50nl/oocyte), previously prepared according to the method disclosed in Nathan Dascal et al., Proc. Natl. Acad. Sci. USA, Vol. 90, pp.6596-6600 (1993). The cRNA-injected oocytes were
35 incubated at 20 °C for 2 to 3 days and subjected to electrophysiological measurements. The measurement was carried

out with a microelectrode-applicable high input resistance amplifier (MEz-8300, Nippon Koden, Co., Japan), and a voltage clamping amplifier (CEz -/200, Nippon Koden, Co., Japan). The initial membrane potential of oocytes was set to -60 mV and responses (current changes of the membrane) evoked by addition of ligands were recorded with a recorder (Thermal Array recorder, Nippon Koden, Co., Japan) (Nathan Dascal et al., Proc. Natl. Acad. Sci. USA, Vol. 90, pp.6596-6600 (1993)).

Typical inward currents elicited upon activation of phospholipase C-coupled receptors were observed in oocytes injected with pMAH2-17 cRNA via stimulation by 10 μ M ATP (Figure 75). In contrast, such a current was not observed in oocytes injected with H₂O, instead of pMAH2-17 cRNA, by the ATP stimulation.

In conclusion, it is considered that the receptor encoded by pMAH2-17 cRNA is classified into a subtype within the ATP receptor, P₂ purinoceptor.

Example 20

Cloning of Rabbit Gastropyrolic Part Smooth Muscle-Derived G Protein Coupled Receptor Protein cDNA

(1) Preparation of Poly(A)⁺ RNA Fraction from Rabbit

Gastropyrolic Part Smooth Muscle and Synthesis of cDNA

A total RNA was prepared from rabbit gastropyrolic part smooth muscles by the guanidine thiocyanate method (Kaplan B.B. et al., Biochem. J. 183, 181-184 (1979)) and, then, poly(A)⁺ RNA fractions were prepared with a mRNA purifying kit (Pharmacia Co.). Next, to 5 μ g of the poly(A)⁺ RNA fraction was added a random DNA hexamer (BRL Co.) as a primer, and the resulting mixture was subjected to reaction with MMLV reverse transcriptase (BRL Co.) in the buffer attached to the MMLV reverse transcriptase kit to synthesize complementary DNAs. The reaction product was extracted with phenol/chloroform (1:1), precipitated in ethanol, and was then dissolved in 30 μ l of TE.

(2) Amplification of Receptor cDNA by PCR Using Rabbit
Gastropyrolic Part Smooth Muscle-Derived cDNA and
Sequencing

By using, as a template, 1 μ l of cDNA prepared
from the rabbit gastropyrolic part smooth muscle in the above
step (1), PCR amplification using the DNA primer having a
nucleotide sequence represented by SEQ ID NO: 10 and the DNA
primer having a nucleotide sequence represented by SEQ ID
NO: 4 synthesized in Example 15 was carried out.
A reaction solution was composed of the synthetic DNA primers
(SEQ: 5' primer sequence and 3' primer sequence) each in an
amount of 100 pM, 0.25 mM dNTPs, 1 μ l of Taq DNA polymerase
and 10 μ l of buffer attached to the enzyme kit, and the
total amount of the reaction solution was made to be 100 μ l.
The cycle for amplification including 96 °C for 30 sec., 45 °C
for 1 min. and 60 °C for 3 min. was repeated 25 times by using
a Thermal Cycler (Perkin-Elmer Co.). The amplified products
were confirmed relying upon 1.2% agarose gel electrophoresis
and ethidium bromide staining.

(3) Subcloning of PCR Product into Plasmid Vector and
Selection of Novel Receptor Candidate Clone via Decoding
Nucleotide Sequence of Inserted cDNA Region

The PCR products obtained in the above step (2)
were separated by using a 1.0% low-melting temperature agarose
gel, the band parts were excised from the gel with a razor
blade, and were electro-eluted, extracted with phenol and
precipitated in ethanol to recover DNAs. According to the
protocol attached to a TA Cloning Kit (Invitrogen Co.),
the recovered DNAs were subcloned to the plasmid vector,
PCRTM II. The recombinant vectors were introduced into
E. coli JM109 competent cells (Takara Shuzo Co., Japan) to
produce transformants. Then, transformant clones having
a cDNA-inserted fragment were selected in an LB agar
culture medium containing ampicillin, IPTG and X-gal. Only
transformant clones exhibiting white color were picked with
a sterilized toothstick to obtain 100 transformant clones.

The individual clones were cultured overnight in an LB culture medium containing ampicillin and treated with the automatic plasmid extracting machine PI-100 (Kurabo Co., Japan) to prepare plasmid DNAs. An aliquot of the DNAs thus prepared was cut by EcoRI to confirm the size of the cDNA fragment that was inserted. An aliquot of the remaining DNAs was further processed with RNase, extracted with phenol/chloroform, and precipitated in ethanol so as to be condensed. Sequencing was carried out by using a DyeDeoxy terminator cycle sequencing kit (ABI Co.), the DNAs were decoded by using a fluorescent automatic sequencer.

Homology retrieval was carried out based upon the determined nucleotide sequence. As a result, it was learned that a novel G protein coupled receptor protein was encoded by the cDNA fragment insert in the plasmid possessed by the transformant Escherichia coli JM109/pMN128. The nucleotide sequences of the cDNA fragments are shown in Figures 71 and 72. To further confirm this fact, by using DNASIS (Hitachi System Engineering Co., Japan) the nucleotide sequences were converted into amino acid sequences [Figure 71 and Figure 72], and homology retrieval was carried out in view of hydrophobicity plotting [Figure 73] and at the amino acid sequence level to find a novel receptor protein which has 27% homology relative to hamster-derived β_2 -adrenaline receptor protein (A03159), 20% homology relative to rat-derived bradykinin receptor (type B_2) protein (A41283), 24% homology relative to human-derived dopamine D_1 receptor protein (S11377) and 23% homology relative to human-derived blue sensitive opsin receptor protein (A03156). The aforementioned abbreviations in parentheses are reference numbers assigned when they are registered as data to NBRF-PIR and are usually called "Accession Numbers".

Example 21

Cloning of cDNA Comprising Whole Coding Regions for Receptor Protein from Human-Derived DNA Library

The DNA library constructed by Clontech wherein
5 λ gt11 phage vector is used (CLONTECH Laboratories, Inc.;
CLH L1008b) was employed as a human placenta-derived cDNA
library. The human placenta cDNA library (1×10^5 pfu (plaque
forming units)) was thermally denatured. By using the human
placenta-derived cDNA library, PCR amplification using
10 the DNA primer having a nucleotide sequence represented by SEQ
ID NO: 20 and the DNA primer having a nucleotide sequence
represented by SEQ ID NO: 23 synthesized in Example 19 was
carried out.

(Nucleotide sequence of synthesized primer)

- 15 ① 5'-TAGTGTGTGGAGTCGTGTGGCTGGCTG-3'
(SEQ ID NO: 20)
② 5'-ACAGGACCTGCTGGGCCATCCTGGCGACACA-3'
(SEQ ID NO: 23)

The isolated DNA was subcloned using a TA Cloning Kit
20 (Invitrogen Co.) and sequencing was carried out. Figure 76
shows a nucleotide sequence of obtained cDNA fragment, ph3H2-17.
As a result, it was learned that ph3H2-17 is highly homologous
to the mouse purinoceptor cDNA fragment, p3H2-17. It is
strongly suggested that the human-derived cDNA fragment is a
25 partial nucleotide sequence of human purinoceptor.

Based on the nucleotide sequence of ph3H2-17 which
was sequenced, the following 2 primers were synthesized:

(Nucleotide sequence of synthesized primer)

- 30 ③ 5'-ACAGCCATCTTCGCTGCCACAGGCAT-3'
(SEQ ID NO: 58)
④ 5'-AGACAGTAGCAGGCCAGCAGGGCAGCAAA-3'
(SEQ ID NO: 59)

The above synthetic 2 primers were each used in combination with λ gt 11 primers (Takara, Japan; catalogue 3864) for obtaining full-length human purinoceptor cDNA. Thus, using thermally denatured, human placenta-derived λ gt 11 cDNA libraries (CLONTECH; CLHL 1008b), first RCR amplification using a combination of the DNA primer having a nucleotide sequence represented by SEQ ID NO: 20 with λ gt 11 Forward primer, of the DNA primer having a nucleotide sequence represented by SEQ ID NO: 20 with λ gt 11 Reverse primer, of the DNA primer having a nucleotide sequence represented by SEQ ID NO: 23 with λ gt 11 Forward primer, and of the DNA primer having a nucleotide sequence represented by SEQ ID NO: 23 with λ gt 11 Reverse primer was carried out with Ex Taq polymerase (Takara, Japan) (30 cycles; 95°C/30 seconds, 55 °C/60 seconds, and 72 °C/60 seconds), respectively.

Next, by using a 1/50 of the 1st PCR product, second RCR amplification was carried in the same manner as in the first PCR except for using the DNA primer having a nucleotide sequence represented by SEQ ID NO: 58 in place of SEQ ID NO: 20 and the DNA primer having a nucleotide sequence represented by SEQ ID NO: 59 in place of SEQ ID NO: 23 (30 cycles; 95°C/30 seconds, 65 °C/60 seconds and 72 °C/60 seconds). The amplified product DNA was subcloned using a TA Cloning Kit (Invitrogen Co.) and sequencing was carried out for three clones each of 5' and 3' sides (Figure 77).

Based on the amino acid sequence (Figure 77) deduced from the determined nucleotide sequence of human purinoceptor cDNA as shown in Figure 77, hydrophobicity plotting was carried out (Figure 78). As a result, it was learned that the human-derived receptor is a novel seven transmembrane G protein coupled receptor, similarly to the mouse type. It was also learned that the deduced amino acid sequence of human receptor has 87% homology relative to the amino acid sequence of mouse purinoceptor and its amino acid residues are well conserved (Figure 79).

Clones free of PCR errors which often occur in a PCR amplification were selected and restriction enzyme regions

comprising overlapping areas were obtained therefrom.

The restriction enzyme regions thus obtained were subjected to construction of plasmid pAH2-17 having a full-length open reading frame of human purinoceptor cDNA. The plasmid pAH2-17 is possessed by transformant Escherichia coli JM109/pAH2-17.

The DNA primers of the present invention allow efficient amplification of DNAs that encode G protein coupled receptor proteins. This makes it possible to efficiently screen for the DNAs coding for G protein coupled receptor proteins and to accomplish the cloning.

The G protein coupled receptor protein of the present invention and their G protein coupled receptor protein-encoding DNA are advantageously useful in:

- ① determining ligands,
- ② obtaining antibodies and an antisera,
- ③ constructing systems for expressing recombinant receptor proteins,
- ④ investigating or developing receptor-binding assay systems and screening for pharmaceutical candidate compounds, by using the above expression system
- ⑤ designing drugs based upon comparisons with ligands and receptors having a structure similar or analogous thereto,
- ⑥ preparing probes and/or PCR primers in gene diagnosis, and
- ⑦ gene manipulating therapy.

In particular, discovering the structure and properties of the G protein coupled receptor will lead to the development of unique pharmaceuticals acting upon these systems.

The practice of the present invention will employ, otherwise indicated, conventional techniques of molecular biology, microbiology, recombinant DNA, pharmacology, immunology, bioscience, and medical technology, which are within the skill of the art. All patents, patent applications, and publications mentioned herein, both supra and infra, are hereby incorporated herein by reference.

- 2 3 8 -

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT:

(A) NAME: Takeda Chemical Industries, Ltd.
(B) STREET: 1-1, Doshomachi 4-chome, Chuo-ku
(C) CITY: Osaka-shi
(D) STATE: Osaka
(E) COUNTRY: Japan
(F) POSTAL CODE (ZIP): 541

(ii) TITLE OF INVENTION: G Protein Coupled Receptor Protein,
Production, And Use Thereof

(iii) NUMBER OF SEQUENCES: 61

(iv) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version #1.25 (EPO)

(2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 25
(B) TYPE: Nucleic acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Other nucleic acid
Synthetic DNA

(iii) FEATURES: N is A, G, C, or T

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

CGTGGSCMTS STGGGCAACN YCCTG 25

(2) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 27
(B) TYPE: Nucleic acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Other nucleic acid
Synthetic DNA

(iii) FEATURES: N is A, G, C, or T

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

GTNGWRRGGC ANCCAGCAGA KGGCAAA 27

- 2 3 9 -

(2) INFORMATION FOR SEQ ID NO: 3:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 27
 - (B) TYPE: Nucleic acid
 - (C) STRANDEDNESS: Single
 - (D) TOPOLOGY: Linear
- (ii) MOLECULE TYPE: Other nucleic acid
Synthetic DNA
- (iii) FEATURES: N is inosine
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

CTCGCSGCMY TNRGYATGGA YCGNTAT 27

(2) INFORMATION FOR SEQ ID NO: 4:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 30
 - (B) TYPE: Nucleic acid
 - (C) STRANDEDNESS: Single
 - (D) TOPOLOGY: Linear
- (ii) MOLECULE TYPE: Other nucleic acid
Synthetic DNA
- (iii) FEATURES: N is inosine
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

CATGTRGWAG GGAANCCAGS AMANRARRAA 30

(2) INFORMATION FOR SEQ ID NO: 5:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 27
 - (B) TYPE: Nucleic acid
 - (C) STRANDEDNESS: Single
 - (D) TOPOLOGY: Linear
- (ii) MOLECULE TYPE: Other nucleic acid
Synthetic DNA
- (iii) FEATURES: N is inosine
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

CTGACYGYTC TNRSNRYTGA CMGVTAC 27

(2) INFORMATION FOR SEQ ID NO: 6:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 27
 - (B) TYPE: Nucleic acid
 - (C) STRANDEDNESS: Single

- 2 4 0 -

(D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Other nucleic acid
Synthetic DNA

(iii) FEATURES: N is inosine

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

CTGACYGYTC TNRSNRYTGA CMGVTAT 27

(2) INFORMATION FOR SEQ ID NO: 7:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 27
(B) TYPE: Nucleic acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear(ii) MOLECULE TYPE: Other nucleic acid
Synthetic DNA

(iii) FEATURES: N is inosine

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

CTCGCSGCMY TNRGYATGGA YCGNTAC 27

(2) INFORMATION FOR SEQ ID NO: 8:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 30
(B) TYPE: Nucleic acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear(ii) MOLECULE TYPE: Other nucleic acid
Synthetic DNA

(iii) FEATURES: N is inosine

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

GATGTGRTAR GGSRNCCAAC AGANGRYAAA 30

(2) INFORMATION FOR SEQ ID NO: 9:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 30
(B) TYPE: Nucleic acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear(ii) MOLECULE TYPE: Other nucleic acid
Synthetic DNA

- 2 4 1 -

(iii) FEATURES: N is inosine

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:

GATGTGRTAR GGSRNCCAAC AGANGRYGAA 30

(2) INFORMATION FOR SEQ ID NO: 10:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 27
(B) TYPE: Nucleic acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Other nucleic acid
Synthetic DNA

(iii) FEATURES: N is inosine

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:

GYCACCAACN WSTTCATCCT SWNHCTG 27

(2) INFORMATION FOR SEQ ID NO: 11:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 27
(B) TYPE: Nucleic acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Other nucleic acid
Synthetic DNA

(iii) FEATURES: N is inosine

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11:

ASNSANRAAG SARTAGANGA NRGGRIT 27

(2) INFORMATION FOR SEQ ID NO: 12:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 25
(B) TYPE: Nucleic acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Other nucleic acid
Synthetic DNA

(iii) FEATURES: N is inosine

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12:

TGNTSSTKMT NGSNGTKGTN GGNA 25

- 2 4 2 -

(2) INFORMATION FOR SEQ ID NO: 13:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 25
 - (B) TYPE: Nucleic acid
 - (C) STRANDEDNESS: Single
 - (D) TOPOLOGY: Linear
- (ii) MOLECULE TYPE: Other nucleic acid
Synthetic DNA
- (iii) FEATURES: N is inosine
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 13:

AYCKGTAYCK GTCCANKGWN ATKGC 25

(2) INFORMATION FOR SEQ ID NO: 14:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 24
 - (B) TYPE: Nucleic acid
 - (C) STRANDEDNESS: Single
 - (D) TOPOLOGY: Linear
- (ii) MOLECULE TYPE: Other nucleic acid
Synthetic DNA
- (iii) FEATURES: N is inosine
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 14:

CATKKCCSTG GASAGNTAYN TRGC 24

(2) INFORMATION FOR SEQ ID NO: 15:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 24
 - (B) TYPE: Nucleic acid
 - (C) STRANDEDNESS: Single
 - (D) TOPOLOGY: Linear
- (ii) MOLECULE TYPE: Other nucleic acid
Synthetic DNA
- (iii) FEATURES: N is inosine
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 15:

GWWGGGSAKC CAGCASANGG CRAA 24

(2) INFORMATION FOR SEQ ID NO: 16:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 18

- 2 4 3 -

(B) TYPE: Nucleic acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Other nucleic acid
Synthetic DNA

(iii) FEATURES: 15th N is A, G, C, or T
6th, 9th, 10th & 12th Ns are inosine

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 16:

ARYYTNGCNN TNGCNGAY 18

(2) INFORMATION FOR SEQ ID NO: 17:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 21
(B) TYPE: Nucleic acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Other nucleic acid
Synthetic DNA

(iii) FEATURES: 13th, 15th, 16th & 18th Ns are
each A, G, C, or T
1st, 4th, 6th Ns are inosine

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 17:

NGGNANCCAR CANANNRNRA A 21

(2) INFORMATION FOR SEQ ID NO: 18:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 27
(B) TYPE: Nucleic acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Other nucleic acid
Synthetic DNA

(iii) FEATURES: N is inosine

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 18:

GCCTSNTNRN SATGWSTGTG GANMGNT 27

(2) INFORMATION FOR SEQ ID NO: 19:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 27
(B) TYPE: Nucleic acid
(C) STRANDEDNESS: Single

- 2 4 4 -

(D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Other nucleic acid
Synthetic DNA

(iii) FEATURES: N is inosine

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 19:

GAWSNTGMYN ANRTGGWAGG GNANCCA 27

(2) INFORMATION FOR SEQ ID NO: 20:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 27
(B) TYPE: Nucleic acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear(ii) MOLECULE TYPE: Other nucleic acid
Synthetic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 20:

TAGTGTGTGG AGTCGTGTGG CTGGCTG 27

(2) INFORMATION FOR SEQ ID NO: 21:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 27
(B) TYPE: Nucleic acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear(ii) MOLECULE TYPE: Other nucleic acid
Synthetic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 21:

AGTCTTTGCT GCCACAGGCA TCCAGCG 27

(2) INFORMATION FOR SEQ ID NO: 22:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 30
(B) TYPE: Nucleic acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear(ii) MOLECULE TYPE: Other nucleic acid
Synthetic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 22:

CAAGCCAGTA AGGCTATGAA GGGCAGCAAG 30

- 2 4 5 -

(2) INFORMATION FOR SEQ ID NO: 23:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 31
 (B) TYPE: Nucleic acid
 (C) STRANDEDNESS: Single
 (D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Other nucleic acid
 Synthetic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 23:

ACAGGACCTG CTGGGCCATC CTGGCGACAC A 31

(2) INFORMATION FOR SEQ ID NO: 24:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 91
 (B) TYPE: Amino acid
 (C) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 24:

Leu Val Leu Val Ile Ala Arg Val Arg Arg Leu His Asn Val Thr Asn
 1 5 10 15

Phe Leu Ile Gly Asn Leu Ala Leu Ser Asp Val Leu Met Cys Thr Ala
 20 25 30

Cys Val Pro Leu Thr Leu Ala Tyr Ala Phe Glu Pro Arg Gly Trp Val
 35 40 45

Phe Gly Gly Gly Leu Cys His Leu Val Phe Phe Leu Gln Pro Val Thr
 50 55 60

Val Tyr Val Ser Val Phe Thr Leu Thr Thr Ile Ala Val Asp Arg Tyr
 65 70 75 80

Val Val Leu Val His Pro Leu Arg Arg Arg Ile
 85 90

(2) INFORMATION FOR SEQ ID NO: 25:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 59
 (B) TYPE: Amino acid
 (C) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 25:

Gly Leu Leu Leu Val Thr Tyr Leu Leu Pro Leu Leu Val Ile Leu Leu

- 2 4 6 -

1 5 10 15
 Ser Tyr Val Arg Val Ser Val Lys Leu Arg Asn Arg Val Val Pro Gly
 20 25 30
 Cys Val Thr Gln Ser Gln Ala Asp Trp Asp Arg Ala Arg Arg Arg Arg
 35 40 45
 Thr Phe Cys Leu Leu Val Val Val Val Val Val
 50 55

(2) INFORMATION FOR SEQ ID NO: 26:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 370
 (B) TYPE: Amino acid
 (C) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 26:

Met Ala Ser Ser Thr Thr Arg Gly Pro Arg Val Ser Asp Leu Phe Ser
 1 5 10 15
 Gly Leu Pro Pro Ala Val Thr Thr Pro Ala Asn Gln Ser Ala Glu Ala
 20 25 30
 Ser Ala Gly Asn Gly Ser Val Ala Gly Ala Asp Ala Pro Ala Val Thr
 35 40 45
 Pro Phe Gln Ser Leu Gln Leu Val His Gln Leu Lys Gly Leu Ile Val
 50 55 60
 Leu Leu Tyr Ser Val Val Val Val Val Gly Leu Val Gly Asn Cys Leu
 65 70 75 80
 Leu Val Leu Val Ile Ala Arg Val Arg Arg Leu His Asn Val Thr Asn
 85 90 95
 Phe Leu Ile Gly Asn Leu Ala Leu Ser Asp Val Leu Met Cys Thr Ala
 100 105 110
 Cys Val Pro Leu Thr Leu Ala Tyr Ala Phe Glu Pro Arg Gly Trp Val
 115 120 125
 Phe Gly Gly Gly Leu Cys His Leu Val Phe Phe Leu Gln Pro Val Thr
 130 135 140
 Val Tyr Val Ser Val Phe Thr Leu Thr Thr Ile Ala Val Asp Arg Tyr
 145 150 155 160
 Val Val Leu Val His Pro Leu Arg Arg Arg Ile Ser Leu Arg Leu Ser
 165 170 175
 Ala Tyr Ala Val Leu Ala Ile Trp Ala Leu Ser Ala Val Leu Ala Leu

- 2 4 7 -

	180							185						190				
Pro Ala Ala Val His Thr Tyr His Val Glu Leu Lys Pro His Asp Val 195 200 205																		
Arg Leu Cys Glu Glu Phe Trp Gly Ser Gln Glu Arg Gln Arg Gln Leu 210 215 220																		
Tyr Ala Trp Gly Leu Leu Leu Val Thr Tyr Leu Leu Pro Leu Leu Val 225 230 235 240																		
Ile Leu Leu Ser Tyr Val Arg Val Ser Val Lys Leu Arg Asn Arg Val 245 250 255																		
Val Pro Gly Cys Val Thr Gln Ser Gln Ala Asp Trp Asp Arg Ala Arg 260 265 270																		
Arg Arg Arg Thr Phe Cys Leu Leu Val Val Val Val Val Val Phe Ala 275 280 285																		
Val Cys Trp Leu Pro Leu His Val Phe Asn Leu Leu Arg Asp Leu Asp 290 295 300																		
Pro His Ala Ile Asp Pro Tyr Ala Phe Gly Leu Val Gln Leu Leu Cys 305 310 315 320																		
His Trp Leu Ala Met Ser Ser Ala Cys Tyr Asn Pro Phe Ile Tyr Ala 325 330 335																		
Trp Leu His Asp Ser Phe Arg Glu Glu Leu Arg Lys Leu Leu Val Ala 340 345 350																		
Trp Pro Arg Lys Ile Ala Pro His Gly Gln Asn Met Thr Val Ser Val 355 360 365																		
Val Ile 370																		

(2) INFORMATION FOR SEQ ID NO: 27:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 206
(B) TYPE: Amino acid
(C) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 27:

Leu Val Leu Val Ile Ala Arg Val Arg Arg Leu Tyr Asn Val Thr Asn
1 5 10 15
Phe Leu Ile Gly Asn Leu Ala Leu Ser Asp Val Leu Met Cys Thr Ala
20 25 30
Cys Val Pro Leu Thr Leu Ala Tyr Ala Phe Glu Pro Arg Gly Trp Val

- 2 4 8 -

[illegible]

(2) INFORMATION FOR SEO ID NO: 28:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 126
(B) TYPE: Amino acid
(C) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 28:

Val	Val	Leu	Val	His	Pro	Leu	Arg	Arg	Arg	Ile	Ser	Leu	Arg	Leu	Ser
1				5					10					15	
Ala	Tyr	Ala	Val	Leu	Gly	Ile	Trp	Ala	Leu	Ser	Ala	Val	Leu	Ala	Leu
			20					25					30		
Pro	Ala	Ala	Val	His	Thr	Tyr	His	Val	Glu	Leu	Lys	Pro	His	Asp	Val
		35					40					45			
Ser	Leu	Cys	Glu	Glu	Phe	Trp	Gly	Ser	Gln	Glu	Arg	Gln	Arg	Gln	Ile
	50					55					60				
Tyr	Ala	Trp	Gly	Leu	Leu	Leu	Gly	Thr	Tyr	Leu	Leu	Pro	Leu	Leu	Ala

- 2.4 9 -

65	70	75	80
Ile Leu Leu Ser Tyr Val Arg Val Ser Val Lys Leu Arg Asn Arg Val			
85		90	95
Val Pro Gly Ser Val Thr Gln Ser Gln Ala Asp Trp Asp Arg Ala Arg			
100	105		110
Arg Arg Arg Thr Phe Cys Leu Leu Val Val Val Val Val Val			
115	120		125

(2) INFORMATION FOR SEQ ID NO: 29:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 273
 (B) TYPE: Nucleic acid
 (C) STRANDEDNESS: Double
 (D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE

(C) IDENTIFICATION METHOD: S

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 29:

```

CTGGTGCTGG TGATCGCGCG GGTGCGCCGG CTGCACAACG TGACGAACTT CCTCATCGGC   60
AACCTGGCCT TGTCGACGT GTCATGTGC ACCGCTGCG TGCCGCTCAC GCTGGCCTAT   120
GCCTTCGAGC CACGCGGCTG GGTGTTCCGGC GGCGGCCTGT GCCACCTGGT CTTCTTCCTG   180
CAGCCGGTCA CCGTCTATGT GTCGGTGTTC ACGCTACCA CCATCGCAGT GGACCGGTAC   240
GTCGTGCTGG TGCACCCGCT GAGGCGGCGC ATC                               273

```

(2) INFORMATION FOR SEQ ID NO: 30:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 177
 (B) TYPE: Nucleic acid
 (C) STRANDEDNESS: Double
 (D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE

(C) IDENTIFICATION METHOD: S

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 30:

```

GGCCTGCTGC TGGTCACCTA CCTGCTCCCT CTGCTGGTCA TCCTCCTGTC TTACGTCCGG   60
GTGTCACTGA AGCTCCGCAA CCGCGTGGTG CCGGGCTGCG TGACCCAGAG CCAGGCCGAC   120
TGGGACCGCG CTCGGCGCCG GCGCACCTTC TGCTTGCTGG TGGTGGTCGT GGTGGTG   177

```

- 2 5 0 -

(2) INFORMATION FOR SEQ ID NO: 31:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1110
 (B) TYPE: Nucleic acid
 (C) STRANDEDNESS: Double
 (D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE

(C) IDENTIFICATION METHOD: S

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 31:

ATGGCCTCAT CGACCACTCG GGGCCCCAGG GTTTCTGACT TATTTTCTGG GCTGCCGCCG	60
GCGGTCACAA CTCCGCCAA CCAGAGCGCA GAGGCCTCGG CGGGCAACGG GTCGGTGGCT	120
GGCGCGGACG CTCCAGCCGT CACGCCCTTC CAGAGCCTGC AGCTGGTGCA TCAGCTGAAG	180
GGGCTGATCG TGCTGCTCTA CAGCGTCGTG GTGGTCGTGG GGCTGGTGGG CAACTGCCTG	240
CTGGTGCTGG TGATCGCGCG GGTGCGCCGG CTGCACAACG TGACGAACTT CCTCATCGGC	300
AACCTGGCCT TGTCCGACGT GCTCATGTGC ACCGCCTGCG TGCCGCTCAC GCTGGCCTAT	360
GCCTTCGAGC CACGCGGCTG GGTGTTTCGGC GGCGGCCTGT GCCACCTGGT CTTCTTCCTG	420
CAGCCGGTCA CCGTCTATGT GTCGGTGTTT ACGCTCACCA CCATCGCAGT GGACCGCTAC	480
GTCGTGCTGG TGCACCCGCT GAGGCGGCGC ATCTCGCTGC GCCTCAGCGC CTACGCTGTG	540
CTGGCCATCT GGGCGCTGTC CGCGGTGCTG GCGCTGCCCC CGCCCGTGCA CACCTATCAC	600
GTGGAGCTCA AGCCGCACGA CGTGCGCCTC TGCAGGAGT TCTGGGGCTC CCAGGAGCGC	660
CAGCGCCAGC TCTACGCCTG GGGGCTGCTG CTGGTCACCT ACCTGCTCCC TCTGCTGGTC	720
ATCCTCCTGT CTTACGTCCG GGTGTCAGTG AAGCTCCGCA ACCGCGTGGT GCCGGGCTGC	780
GTGACCCAGA GCCAGGCCGA CTGGGACCGC GCTCGGCGCC GGCGCACCTT CTGCTTGCTG	840
GTGGTGGTCG TGGTGGTGTT CGCCGTCTGC TGGCTGCCGC TGCACGTCTT CAACCTGCTG	900
CGGGACCTCG ACCCCCACGC CATCGACCCT TACGCCTTTG GGCTGGTGCA GCTGCTCTGC	960
CACTGGCTCG CCATGAGTTC GGCCTGCTAC AACCCTTCA TCTACGCCTG GCTGCACGAC	1020
AGCTTCCGCG AGGAGCTGCG CAAACTGTTG GTCGCTTGGC CCCGCAAGAT AGCCCCCAT	1080
GGCCAGAATA TGACCGTCAG CGTGGTCATC	1110

(2) INFORMATION FOR SEQ ID NO: 32:

- 2 5 1 -

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 618
(B) TYPE: Nucleic acid
(C) STRANDEDNESS: Double
(D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE

(C) IDENTIFICATION METHOD: S

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 32:

CTGGTGCTGG TGATCGCGCG GGTGCGCCGG CTGTACAACG TGACGAATTT CCTCATCGGC 60
AACCTGGCCT TGTCCGACGT GCTCATGTGC ACCGCCTGCG TGCCGCTCAC GCTGGCCTAT 120
GCCTTCGAGC CACGCGGCTG GGTGTTCTGGC GCGGGCCTGT GCCACCTGGT CTTCTTCCTG 180
CAGGCGGTCA CCGTCTATGT GTCGGTGTTT ACGCTACCA CCATCGCAGT GGACCGCTAC 240
GTCGTGCTGG TGCACCCGCT GAGGCGGCGC ATCTCGCTGC GCCTCAGCGC CTACGCTGTG 300
CTGGCCATCT GGGTGCTGTC CGCGGTGCTG GCGCTGCCCC CCGCCGTGCA CACCTATCAC 360
GTGGAGCTCA AGCCGCACGA CGTGCGCCTC TGCAGGAGT TCTGGGGCTC CCAGGAGCGC 420
CAGCGCCAGC TCTACGCCTG GGGGCTGCTG CTGGTCACCT ACCTGCTCCC TCTGCTGGTC 480
ATCCTCCTGT CTTACGCCCC GGTGTCAGTG AAGCTCCGCA ACCGCGTGGT GCCGGGCCGC 540
GTGACCCAGA GCCAGGCCGA CTGGGACCGC GCTCGGCGCC GGCGCACCTT CTGCTTGCTG 600
GTGGTGGTGG TGGTGGTG 618

(2) INFORMATION FOR SEQ ID NO: 33:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 378
(B) TYPE: Nucleic acid
(C) STRANDEDNESS: Double
(D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE

(C) IDENTIFICATION METHOD: S

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 33:

GTGGTTCTGG TGCACCCGCT ACGTCGGCGC ATTTCACTGA GGCTCAGCGC CTACGCGGTG 60
CTGGGCATCT GGGCTCTATC TGCAGTGCTG GCGCTGCCGG CCGCGGTGCA CACCTACCAT 120
GTGGAGCTCA AGCCCCACGA CGTGAGCCTC TGCAGGAGT TCTGGGGCTC GCAGGAGCGC 180

- 2 5 2 -

CAACGCCAGA TCTACGCCTG GGGGCTGCTT CTGGGCACCT ATTTGCTCCC CCTGCTGGCC 240
 ATCCTCCTGT CTTACGTACG GGTGTCAGTG AAGCTGAGGA ACCGCGTGGT GCCTGGCAGC 300
 GTGACCCAGA GTCAAGCTGA CTGGGACCGA GCGCGTCGCC GCCGCACTTT CTGTCTGCTG 360
 GTGGTGGTGG TGGTAGTG 378

(2) INFORMATION FOR SEQ ID NO: 34:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 70
 (B) TYPE: Amino acid
 (C) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 34:

Val Cys His Val Ile Phe Lys Asn Gln Arg Met His Ser Ala Thr Ser
 1 5 10 15
 Leu Phe Ile Val Asn Leu Ala Val Ala Asp Ile Met Ile Thr Leu Ile
 20 25 30
 Asn Thr Pro Phe Thr Leu Val Arg Phe Val Asn Ser Thr Trp Ile Phe
 35 40 45
 Gly Lys Gly Met Cys His Val Ser Arg Phe Ala Gln Tyr Cys Ser Leu
 50 55 60
 His Val Ser Ala Leu Thr
 65 70

(2) INFORMATION FOR SEQ ID NO: 35:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 71
 (B) TYPE: Amino acid
 (C) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 35:

Glu Pro Ala Asp Leu Phe Trp Lys Asn Leu Asp Leu Pro Thr Phe Ile
 1 5 10 15
 Leu Leu Asn Ile Leu Pro Leu Leu Ile Ile Ser Val Ala Tyr Val Arg
 20 25 30
 Val Thr Lys Lys Leu Trp Leu Cys Asn Met Ile Val Asp Val Thr Thr
 35 40 45
 Glu Gln Tyr Phe Ala Leu Arg Pro Lys Lys Lys Lys Thr Ile Lys Met
 50 55 60

- 2 5 3 -

Leu Met Leu Val Val Val Leu
65 70

(2) INFORMATION FOR SEQ ID NO: 36:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 210
(B) TYPE: Nucleic acid
(C) STRANDEDNESS: Double
(D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE

(C) IDENTIFICATION METHOD: S

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 36:

GTCTGTCATG TCATCTTCAA GAACCAGCGA ATGCACTCGG CCACCAGCCT CTTTCATCGTC 60
AACCTGGCAG TTGCCGACAT AATGATCAG CTGCTCAACA CCCCCTTCAC TTTGGTTCGC 120
TTTGTGAACA GCACATGGAT ATTTGGAAG GGCATGTGCC ATGTCAGCCG CTTTGCCCAG 180
TACTGCTCAC TGCACGTCTC AGCACTGACA 210

(2) INFORMATION FOR SEQ ID NO: 37:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 213
(B) TYPE: Nucleic acid
(C) STRANDEDNESS: Double
(D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE

(C) IDENTIFICATION METHOD: S

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 37:

GAGCCAGCTG ACCTCTTCTG GAAGAACCTG GACTTGCCCA CCTTCATCCT GCTCAACATC 60
CTGCCCCCTCC TCATCATCTC TGTGGCCTAC GTTCGTGTGA CCAAGAAACT GTGGCTGTGT 120
AATATGATTG TCGATGTGAC CACAGAGCAG TACTTTGCCC TCGGGCCCAA AAAGAAGAAG 180
ACCATCAAGA TGTTGATGCT GGTGGTAGTC CTC 213

(2) INFORMATION FOR SEQ ID NO: 38:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 115
(B) TYPE: Amino acid
(C) TOPOLOGY: Linear

- 2 5 4 -

(ii) MOLECULE TYPE: Peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 38:

Ala Ser Trp His Lys Arg Gly Gly Arg Arg Ala Ala Trp Val Val Cys
 1 5 10 15

Gly Val Val Trp Leu Ala Val Thr Ala Gln Cys Leu Pro Thr Ala Val
 20 25 30

Phe Ala Ala Thr Gly Ile Gln Arg Asn Arg Thr Val Cys Tyr Asp Leu
 35 40 45

Ser Pro Pro Ile Leu Ser Thr Arg Tyr Leu Pro Tyr Gly Met Ala Leu
 50 55 60

Thr Val Ile Gly Phe Leu Leu Pro Phe Ile Ala Leu Leu Ala Cys Tyr
 65 70 75 80

Cys Arg Met Ala Arg Arg Leu Cys Arg Gln Asp Gly Pro Ala Gly Pro
 85 90 95

Val Ala Gln Glu Arg Arg Ser Lys Ala Ala Arg Met Ala Val Val Val
 100 105 110

Ala Ala Val
 115

(2) INFORMATION FOR SEQ ID NO: 39:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 328
 (B) TYPE: Amino acid
 (C) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 39:

Met Glu Gln Asp Asn Gly Thr Ile Gln Ala Pro Gly Leu Pro Pro Thr
 1 5 10 15

Thr Cys Val Tyr Arg Glu Asp Phe Lys Arg Leu Leu Leu Thr Pro Val
 20 25 30

Tyr Ser Val Val Leu Val Val Gly Leu Pro Leu Asn Ile Cys Val Ile
 35 40 45

Ala Gln Ile Cys Ala Ser Arg Arg Thr Leu Thr Arg Ser Ala Val Tyr
 50 55 60

Thr Leu Asn Leu Ala Leu Ala Asp Leu Met Tyr Ala Cys Ser Leu Pro
 65 70 75 80

- 2 5 5 -

Leu Leu Ile Tyr Asn Tyr Ala Arg Gly Asp His Trp Pro Phe Gly Asp
 85 90 95

Leu Ala Cys Arg Phe Val Arg Phe Leu Phe Tyr Ala Asn Leu His Gly
 100 105 110

Ser Ile Leu Phe Leu Thr Cys Ile Ser Phe Gln Arg Tyr Leu Gly Ile
 115 120 125

Cys His Pro Leu Ala Ser Trp His Lys Arg Gly Gly Arg Arg Ala Ala
 130 135 140

Trp Val Val Cys Gly Val Val Trp Leu Ala Val Thr Ala Gln Cys Leu
 145 150 155 160

Pro Thr Ala Val Phe Ala Ala Thr Gly Ile Gln Arg Asn Arg Thr Val
 165 170 175

Cys Tyr Asp Leu Ser Pro Pro Ile Leu Ser Thr Arg Tyr Leu Pro Tyr
 180 185 190

Gly Met Ala Leu Thr Val Ile Gly Phe Leu Leu Pro Phe Ile Ala Leu
 195 200 205

Leu Ala Cys Tyr Cys Arg Met Ala Arg Arg Leu Cys Arg Gln Asp Gly
 210 215 220

Pro Ala Gly Pro Val Ala Gln Glu Arg Arg Ser Lys Ala Ala Arg Met
 225 230 235 240

Ala Val Val Val Ala Ala Val Phe Ala Ile Ser Phe Leu Pro Phe His
 245 250 255

Ile Thr Lys Thr Ala Tyr Leu Ala Val Arg Ser Thr Pro Gly Val Ser
 260 265 270

Cys Pro Val Leu Glu Thr Phe Ala Ala Ala Tyr Lys Gly Thr Arg Pro
 275 280 285

Phe Ala Ser Val Asn Ser Val Leu Asp Pro Ile Leu Phe Tyr Phe Thr
 290 295 300

Gln Gln Lys Phe Arg Arg Gln Pro His Asp Leu Leu Gln Arg Leu Thr
 305 310 315 320

Ala Lys Trp Gln Arg Gln Arg Val
 325

(2) INFORMATION FOR SEQ ID NO: 40:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 345
- (B) TYPE: Nucleic acid
- (C) STRANDEDNESS: Double
- (D) TOPOLOGY: Linear

- 2 5 6 -

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE

(C) IDENTIFICATION METHOD: S

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 40:

```
GCTTCCTGGC ACAAGCGTGG AGGTCGCCGT GCTGCTTGGG TAGTGTGTGG AGTCGTGTGG 60
CTGGCTGTGA CAGCCCAGTG CCTGCCCACG GCAGTCTTTG CTGCCACAGG CATCCAGCGC 120
AACCGCACTG TGTGCTACGA CCTGAGCCCA CCCATCCTGT CTACTCGCTA CCTGCCCTAT 180
GGTATGGCCC TCACGGTCAT CGGCTTCTTG CTGCCCTTCA TAGCCTTACT GGCTTGTTAT 240
TGTCGCATGG CCCGCCGCCT GTGTCGCCAG GATGGCCCAG CAGGTCCTGT GGCCCAAGAG 300
CGGCGCAGCA AGGCGGCTCG TATGGCTGTG GTGGTGGCAG CTGTC 345
```

(2) INFORMATION FOR SEQ ID NO: 41:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 984
(B) TYPE: Nucleic acid
(C) STRANDEDNESS: Double
(D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE

(C) IDENTIFICATION METHOD: S

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 41:

```
ATGGAGCAGG ACAATGGCAC CATCCAGGCT CCAGGCTTGC CGCCCACCAC CTGCGTCTAC 60
CGTGAGGATT TCAAGCGACT GCTGCTAACC CCGGTATACT CGGTGGTGCT GGTGGTCGGC 120
CTGCCACTGA ACATCTGCGT CATTGCCCAG ATCTGCGCAT CCCGCCGGAC CCTGACCCGT 180
TCCGCTGTGT ACACCCTGAA CCTGGCACTG GCGGACCTGA TGTATGCCTG TTTACTACCC 240
CTACTTATCT ATAACTACGC CAGAGGGGAC CACTGGCCCT TCGGAGACCT CGCCTGCCGC 300
TTTGTACGCT TCCTCTTCTA TGCCAATCTA CATGGCAGCA TCCTGTTTCT CACCTGCATT 360
AGCTTCCAGC GCTACCTGGG CATCTGCCAC CCCCTGGCTT CCTGGCACAA GCGTGGAGGT 420
CGCCGTGCTG CTGGGGTAGT GTGTGGAGTC GTGTGGCTGG CTGTGACAGC CCAGTGCCTG 480
CCCACGGCAG TCTTTGCTGC CACAGGCATC CAGCGCAACC GCACTGTGTG CTACGACCTG 540
AGCCCACCCA TCCTGTCTAC TCGCTACCTG CCCTATGGTA TGGCCCTCAC GGTTCATCGGC 600
TTCTTGCTGC CCTTCATAGC CTTACTGGCT TGTTATTGTC GCATGGCCCC CCGCCTGTGT 660
```

- 2 5 7 -

CGCCAGGATG GCCCAGCAGG TCCTGTGGCC CAAGAGCGGC GCAGCAAGGC GGCTCGTATG 720
 GCTGTGGTGG TGGCAGCTGT CTTTGCCATC AGCTTCCTGC CTTTCCACAT CACCAAGACA 780
 GCCTACTTGG CTGTGCGCTC CACGCCCCGT GTCTCTTGCC CTGTGCTGGA GACCTTCGCT 840
 GCTGCCTACA AAGGCACTCG GCCCTTCGCC AGTGTCAACA GTGTTCTGGA CCCCATTCTC 900
 TTCTACTTCA CACAACAGAA GTTCCGGCGG CAACCCACG ATCTCTTACA GAGGCTCACA 960
 GCCAAGTGGC AGAGGCAGAG AGTC 984

(2) INFORMATION FOR SEQ ID NO: 42:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 128
 (B) TYPE: Amino acid
 (C) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 42:

Ala Ala Met Ser Val Asp Arg Tyr Val Ala Ile Val His Ser Arg Arg
 1 5 10 15
 Ser Ser Ser Leu Arg Val Ser Arg Asn Ala Leu Leu Gly Val Gly Phe
 20 25 30
 Ile Trp Ala Leu Ser Ile Ala Met Ala Ser Pro Val Ala Tyr His Gln
 35 40 45
 Arg Leu Phe His Arg Asp Ser Asn Gln Thr Phe Cys Trp Glu Gln Trp
 50 55 60
 Pro Asn Lys Leu His Lys Lys Ala Tyr Val Val Cys Thr Phe Val Phe
 65 70 75 80
 Gly Tyr Leu Leu Pro Leu Leu Leu Ile Cys Phe Cys Tyr Ala Lys Val
 85 90 95
 Leu Asn His Leu His Lys Lys Leu Lys Asn Met Ser Lys Lys Ser Glu
 100 105 110
 Ala Ser Lys Lys Lys Thr Ala Gln Thr Val Leu Val Val Val Val Val
 115 120 125

(2) INFORMATION FOR SEQ ID NO: 43:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 384
 (B) TYPE: Nucleic acid
 (C) STRANDEDNESS: Double
 (D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: cDNA

- 2 5 8 -

(ix) FEATURE

(C) IDENTIFICATION METHOD: S

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 43:

```

GCCGCGATGT CTGTGGATCG CTACGTGGCC ATTGTGCACT CGCGGCGCTC CTCCTCCCTC      60
AGGGTGTCCC GCAACGCACT GCTGGGCGTG GGCTTCATCT GGGCGCTGTC CATCGCCATG      120
GCCTCGCCGG TGGCCTACCA CCAGCGTCTT TTCCATCGGG ACAGCAACCA GACCTTCTGC      180
TGGGAGCAGT GGCCCAACAA GCTCCACAAG AAGGCTTACG TGGTGTGCAC TTTCGTCTTT      240
GGGTACCTTC TGCCCTTACT GTCATCTGCG TTTTGCTATG CCAAGGTCCT TAATCATCTG      300
CATAAAAAGC TGAAAAACAT GTCAAAAAG TCTGAAGCAT CCAAGAAAAA GACTGCACAG      360
ACCGTCCTGG TGGTCGTTGT AGTA                                           384

```

(2) INFORMATION FOR SEQ ID NO: 44:

(i) SEQUENCE CHARACTERISTICS:

```

(A) LENGTH:      71
(B) TYPE:         Amino acid
(C) TOPOLOGY:     Linear

```

(ii) MOLECULE TYPE: Peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 44:

```

Val Leu Trp Phe Phe Gly Phe Ser Ile Lys Arg Thr Pro Phe Ser Val
 1               5               10               15
Tyr Phe Leu His Leu Ala Ser Ala Asp Gly Ala Tyr Leu Phe Ser Lys
          20               25               30
Ala Val Phe Ser Leu Leu Asn Ala Gly Gly Phe Leu Gly Thr Phe Ala
          35               40               45
His Tyr Val Arg Ser Val Ala Arg Val Leu Gly Leu Cys Ala Phe Val
          50               55               60
Ala Gly Val Ser Leu Leu Pro
65               70

```

(2) INFORMATION FOR SEQ ID NO: 45:

(i) SEQUENCE CHARACTERISTICS:

```

(A) LENGTH:      215
(B) TYPE:         Nucleic acid
(C) STRANDEDNESS: Double
(D) TOPOLOGY:     Linear

```

(ii) MOLECULE TYPE: cDNA

- 2 5 9 -

(ix) FEATURE

(C) IDENTIFICATION METHOD: S

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 45:

```

GTGCTCTGGT TCTTCGGCTT CTCCATCAAG AGGACCCCCT TCTCCGTCTA CTCCTGCAC      60
CTGGCCAGCG CCGACGGCGC CTACCTCTTC AGCAAGGCCG TGTTCTCCCT GCTGAACGCC      120
GGCGGCTTCC TGGGCACCTT CGCCCACTAT GTGCGCAGCG TGGCCCGGGT GCTGGGGCTC      180
TGC GCCTTCG TGGCGGGCGT GAGCCTCCTG CCGGC                                215

```

(2) INFORMATION FOR SEQ ID NO: 46:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 348
 (B) TYPE: Amino acid
 (C) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 46:

```

Met Glu Leu Ala Met Val Asn Leu Ser Glu Gly Asn Gly Ser Asp Pro
 1           5           10           15
Glu Pro Pro Ala Pro Glu Ser Arg Pro Leu Phe Gly Ile Gly Val Glu
          20           25           30
Asn Phe Ile Thr Leu Val Val Phe Gly Leu Ile Phe Ala Met Gly Val
      35           40           45
Leu Gly Asn Ser Leu Val Ile Thr Val Leu Ala Arg Ser Lys Pro Gly
      50           55           60
Lys Pro Arg Ser Thr Thr Asn Leu Phe Ile Leu Asn Leu Ser Ile Ala
      65           70           75           80
Asp Leu Ala Tyr Leu Leu Phe Cys Ile Pro Phe Gln Ala Thr Val Tyr
          85           90           95
Ala Leu Pro Thr Trp Val Leu Gly Ala Phe Ile Cys Lys Phe Ile His
      100           105           110
Tyr Phe Phe Thr Val Ser Met Leu Val Ser Ile Phe Thr Leu Ala Ala
      115           120           125
Met ser Val Asp Arg Tyr Val Ala Ile Val His Ser Arg Arg Ser Ser
      130           135           140
Ser Leu Arg Val Ser Arg Asn Ala Leu Leu Gly Val Gly Phe Ile Trp
      145           150           155           160
Ala Leu Ser Ile Ala Met Ala Ser Pro Val Ala Tyr His Gln Arg Leu
          165           170           175

```

- 2 6 0 -

Phe His Arg Asp Ser Asn Gln Thr Phe Cys Trp Glu Gln Trp Pro Asn
 180 185 190
 Lys Leu His Lys Lys Ala Tyr Val Val Cys Thr Phe Val Phe Gly Tyr
 195 200 205
 Leu Leu Pro Leu Leu Leu Ile Cys Phe Cys Tyr Ala Lys Val Leu Asn
 210 215 220
 His Leu His Lys Lys Leu Lys Asn Met Ser Lys Lys Ser Glu Ala Ser
 225 230 235 240
 Lys Lys Lys Thr Ala Gln Thr Val Leu Val Val Val Val Val Phe Gly
 245 250 255
 Ile Ser Trp Leu Pro His His Val Val His Leu Trp Ala Glu Phe Gly
 260 265 270
 Ala Phe Pro Leu Thr Pro Ala Ser Phe Phe Phe Arg Ile Thr Ala His
 275 280 285
 Cys Leu Ala Tyr Ser Asn Ser Ser Val Asn Pro Ile Ile Tyr Ala Phe
 290 295 300
 Leu Ser Glu Asn Phe Arg Lys Ala Tyr Lys Gln Val Phe Lys Cys His
 305 310 315 320
 Val Cys Asp Glu Ser Pro Arg Ser Glu Thr Lys Glu Asn Lys Ser Arg
 325 330 335
 Met Asp Thr Pro Pro Ser Thr Asn Cys Thr His Val
 340 345

(2) INFORMATION FOR SEQ ID NO: 47:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1044
 (B) TYPE: Nucleic acid
 (C) STRANDEDNESS: Double
 (D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE

(C) IDENTIFICATION METHOD: S

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 47:

ATGGAAGTGG CTATGGTGAA CCTCAGTGAA GGAATGGGA GCGACCCAGA GCCGCCAGCC 60
 CCGGAGTCCA GGCCGCTCTT CGGCATTGGC GTGGAGAACT TCATTACGCT GGTAGTGT TT 120
 GGCCTGATTT TCGCGATGGG CGTGCTGGGC AACAGCCTGG TGATCACCGT GCTGGCGCGC 180
 AGCAAACCAG GCAACCCCCG CAGCACCACC AACCTGTTTA TCCTCAATCT GAGCATCGCA 240

- 2 6 1 -

GACCTGGCCT ACCTGCTCTT CTGCATCCCT TTTCAGGCCA CCGTGTATGC ACTGCCCACC 300
 TGGGTGCTGG GCGCCTTCAT CTGCAAGTTT ATACACTACT TCTTCACCGT GTCCATGCTG 360
 GTGAGCATCT TCACCCTGGC CGCGATGTCT GTGGATCGCT ACGTGGCCAT TGTGCACTCG 420
 CGGCGCTCCT CCTCCCTCAG GGTGTCCCGC AACGCACTGC TGGGCGTGGG CTTCATCTGG 480
 GCGCTGTCCA TCGCCATGGC CTCGCCGGTG GCCTACCACC AGCGTCTTTT CCATCGGGAC 540
 AGCAACCAGA CCTTCTGCTG GGAGCAGTGG CCCAACAAGC TCCACAAGAA GGCTTACGTG 600
 GTGTGCACTT TCGTCTTTGG GTACCTTCTG CCCTTACTGC TCATCTGCTT TTGCTATGCC 660
 AAGGTCCTTA ATCATCTGCA TAAAAAGCTG AAAAACATGT CAAAAAAGTC TGAAGCATCC 720
 AAGAAAAAGA CTGCACAGAC CGTCCTGGTG GTCGTTGTAG TATTTGGCAT ATCCTGGCTG 780
 CCCCATCATG TCGTCCACCT CTGGGCTGAG TTTGGAGCCT TCCCACTGAC GCCAGCTTCC 840
 TTCTTCTTCA GAATCACCGC CCATTGCCTG GCATACAGCA ACTCCTCAGT GAACCCCATC 900
 ATATATGCCT TTCTCTCAGA AAAGTCCGG AAGGCGTACA AGCAAGTGTT CAAGTGTCAT 960
 GTTTGCATG AATCTCCACG CAGTGAACT AAGGAAAACA AGAGCCGGAT GGACACCCCG 1020
 CCATCCACCA ACTGCACCCA CGTG 1044

(2) INFORMATION FOR SEQ ID NO: 48:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 125
 (B) TYPE: Amino acid
 (C) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 48:

Leu Leu Thr Leu His Pro Val Trp Ser Gln Lys His Arg Thr Ser His
 1 5 10 15
 Trp Ala Ser Arg Val Val Leu Gly Val Trp Leu Ser Ala Thr Ala Phe
 20 25 30
 Ser Val Pro Tyr Leu Val Phe Arg Glu Thr Tyr Asp Asp Arg Lys Gly
 35 40 45
 Arg Val Thr Cys Arg Asn Asn Tyr Ala Val Ser Thr Asp Trp Glu Ser
 50 55 60
 Lys Glu Met Gln Thr Val Arg Gln Trp Ile His Ala Thr Cys Phe Ile
 65 70 75 80
 Ser Arg Phe Ile Leu Gly Phe Leu Leu Pro Phe Leu Val Ile Gly Phe

95

Ala Val Pro Asn Val Ile Leu Thr Asn Gln Arg Val Lys Asp Val Thr
35 40 45

- 2 6 3 -

Gln Ile Lys Cys Met Glu Leu Lys Asn Glu Leu Gly Arg Gln Trp His
 50 55 60

Lys Ala Ser Asn Tyr Ile Phe Val Gly Ile Phe Trp Leu Val Phe Leu
 65 70 75 80

Leu Leu Ile Ile Phe Tyr Thr Ala Ile Thr Arg Lys Ile Phe Lys Ser
 85 90 95

His Leu Lys Ser Arg Lys Asn Ser Ile Ser Val Lys Lys Lys Ser Ser
 100 105 110

Arg Asn Ile Phe Ser Ile Val
 115

(2) INFORMATION FOR SEQ ID NO: 51:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 357
 (B) TYPE: Nucleic acid
 (C) STRANDEDNESS: Double
 (D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE

- (C) IDENTIFICATION METHOD: S

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 51:

TTCAAGATTG TGAAGCCCCT TTCCACGTCC TTCATCCAGT CTGTGAACTA CAGCAAAC TC 60
 GTCTCGCTGG TGGTCTGGTT GCTCATGCTC CTCCTCGCCG TCCCCAACGT CATTCTCACC 120
 AACCAGAGAG TTAAGGACGT GACGCAGATA AAATGCATGG AACTTAAAAA CGAACTGGGC 180
 CGCCAGTGGC ACAAGGCGTC AAATACATC TTTGTGGGCA TTTTCTGGCT TGTGTTCCCTT 240
 TTGCTAATCA TTTTCTACAC TGCTATCACC AGGAAAATCT TTAAGTCCCA CCTGAAATCC 300
 AGAAAGAATT CCATCTCGGT CAAAAGAAA TCTAGCCGCA ACATCTTCAG CATCGTG 357

(2) INFORMATION FOR SEQ ID NO: 52:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 252
 (B) TYPE: Amino acid
 (C) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 52:

Val Asp Leu Leu Ala Ala Leu Thr Leu Met Pro Leu Ala Met Leu Ser
 1 5 10 15

- 2 6 4 -

Ser Ser Ala Leu Phe Asp His Ala Leu Phe Gly Glu Val Ala Cys Arg
 20 25 30
 Leu Tyr Leu Phe Leu Ser Val Cys Phe Val Ser Leu Ala Ile Leu Ser
 35 40 45
 Val Ser Ala Ile Asn Val Glu Arg Tyr Tyr Tyr Val Val His Pro Met
 50 55 60
 Arg Tyr Glu Val Arg Met Lys Leu Gly Leu Val Ala Ser Val Leu Val
 65 70 75 80
 Gly Val Trp Val Lys Ala Leu Ala Met Ala Ser Val Pro Val Leu Gly
 85 90 95
 Arg Val Ser Trp Glu Glu Gly Pro Pro Ser Val Pro Pro Gly Cys Ser
 100 105 110
 Leu Gln Trp Ser His Ser Ala Tyr Cys Gln Leu Phe Val Val Val Phe
 115 120 125
 Ala Val Leu Tyr Phe Leu Leu Pro Leu Leu Leu Ile Leu Val Val Tyr
 130 135 140
 Cys Ser Met Phe Arg Val Ala Arg Val Ala Ala Met Gln His Gly Pro
 145 150 155 160
 Leu Pro Thr Trp Met Glu Thr Pro Arg Gln Arg Ser Glu Ser Leu Ser
 165 170 175
 Ser Arg Ser Thr Met Val Thr Ser Ser Gly Ala Pro Gln Thr Thr Pro
 180 185 190
 His Arg Thr Phe Gly Gly Gly Lys Ala Ala Val Val Leu Leu Ala Val
 195 200 205
 Gly Gly Gln Phe Leu Leu Cys Trp Leu Pro Tyr Phe Ser Phe His Leu
 210 215 220
 Tyr Val Ala Leu Ser Ala Gln Pro Ile Ala Ala Gly Gln Val Glu Asn
 225 230 235 240
 Val Val Thr Trp Ile Gly Tyr Phe Cys Phe Thr Ser
 245 250

(2) INFORMATION FOR SEQ ID NO: 53:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 756
- (B) TYPE: Nucleic acid
- (C) STRANDEDNESS: Double
- (D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: cDNA

- 2 6 5 -

(ix) FEATURE

(C) IDENTIFICATION METHOD: S

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 53:

```

GTGGACCTGC TGGCTGCCCT GACCCTCATG CCTCTGGCCA TGCTCTCCAG CTCCGCCCTC   60
TTTGACCACG CCCTCTTTGG GGAGGTGGCC TGCCGCCTCT ACTTGTTCTT GAGCGTCTGC   120
TTTGTACAGC TGGCCATCCT CTCGGTGTCC GCCATCAATG TGGAGCGCTA CTATTATGTG   180
TGCCACCCCA TGCCTATGA GGTGCGCATG AAAGTGGGGC TGGTGGCCTC TGTGCTGGTG   240
GGCGTGTGGG TGAAGGCCCT GGCCATGGCT TCTGTGCCAG TGTGGGAAG GGTGTCCTGG   300
GAGGAAGGCC CTCCAGTGT CCCCCAGGC TGTCACTCC AATGGAGCCA CAGTGCCTAC   360
TGCCAGCTTT TCGTGGTGGT CTTGCGCGTC CTCTACTTCC TGCTGCCCCT GCTCCTCATC   420
CTTGTGGTCT ACTGCAGCAT GTTCCGGGTG GTCGTGTGG CTGCCATGCA GCACGGGGCCG   480
CTGCCCACGT GGATGGAGAC GCCCCGGCAA CGCTCCGAGT CTCTCAGCAG CCGCTCCACT   540
ATGGTCACCA GCTCGGGGGC CCCGCAGACC ACCCCTCACC GGACGTTTGG CGGAGGGAAG   600
GCAGCAGTGG TCCTCCTGGC TGTGGGAGGA CAGTTCCTGC TCTGTTGGTT GCCCTACTTC   660
TCCTTCCACC TCTATGTGGC CCTGAGCGCT CAGCCCATTG CAGCGGGGCA GGTGGAGAAC   720
GTGGTGACCT GGATTGGCTA CTTCTGCTTC ACCTCC                               756

```

(2) INFORMATION FOR SEQ ID NO: 54:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 263
 (B) TYPE: Amino acid
 (C) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 54:

```

Ala Asp Val Leu Val Thr Ala Ile Cys Leu Pro Ala Ser Leu Leu Val
 1           5           10           15
Asp Ile Thr Glu Ser Trp Leu Phe Gly His Ala Leu Cys Lys Val Ile
          20           25           30
Pro Tyr Leu Gln Ala Val Ser Val Ser Val Val Val Leu Thr Leu Ser
          35           40           45
Ser Ile Ala Leu Asp Arg Trp Tyr Ala Ile Cys His Pro Leu Leu Phe
          50           55           60
Lys Ser Thr Ala Arg Arg Ala Arg Gly Ser Ile Leu Gly Ile Trp Ala
          65           70           75           80

```

- 2 6 6 -

Val Ser Leu Ala Val Met Val Pro Gln Ala Ala Val Met Glu Cys Ser
 85 90 95

Ser Val Leu Pro Glu Leu Ala Asn Arg Thr Arg Leu Leu Ser Val Cys
 100 105 110

Asp Glu Arg Trp Ala Asp Asp Leu Tyr Pro Lys Ile Tyr His Ser Cys
 115 120 125

Phe Phe Ile Val Thr Tyr Leu Ala Pro Leu Gly Leu Met Ala Met Ala
 130 135 140

Tyr Phe Gln Ile Phe Arg Lys Leu Trp Gly Arg Gln Ile Pro Gly Thr
 145 150 155 160

Thr Ser Ala Leu Val Arg Asn Trp Lys Arg Pro Ser Asp Gln Leu Asp
 165 170 175

Asp Gln Gly Gln Gly Leu Ser Ser Glu Pro Gln Pro Arg Ala Arg Ala
 180 185 190

Phe Leu Ala Glu Val Lys Gln Met Arg Ala Arg Arg Lys Thr Ala Lys
 195 200 205

Met Leu Met Val Val Leu Leu Val Phe Ala Leu Cys Tyr Leu Pro Ile
 210 215 220

Ser Val Leu Asn Val Leu Lys Arg Val Phe Gly Met Phe Arg Gln Ala
 225 230 235 240

Ser Asp Arg Glu Ala Ile Tyr Ala Cys Phe Thr Phe Ser His Trp Leu
 245 250 255

Val Tyr Ala Asn Ser Ala Ala
 260

(2) INFORMATION FOR SEQ ID NO: 55:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 789
 (B) TYPE: Nucleic acid
 (C) STRANDEDNESS: Double
 (D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE

- (C) IDENTIFICATION METHOD: S

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 55:

GCCGATGTGC TGGTGACAGC CATCTGCCTG CCGGCCAGTC TGCTGGTAGA CATCACGGAA 60

TCCTGGCTCT TTGGCCATGC CCTCTGCAAG GTCATCCCCT ATCTACAGGC CGTGTCCGTG 120

- 2 6 7 -

TCAGTGGTCG TGCTGACTCT CAGCTCCATC GCCCTGGACC GCTGGTACGC CATCTGCCAC 180
 CCGCTGTTGT TCAAGAGCAC TGCCCGGCGC GCCCGCGGCT CCATCCTCGG CATCTGGGCG 240
 GTGTCGCTGG CTGTCATGGT GCCTCAGGCT GCTGTCATGG AGTGTAGCAG CGTGCTGCCC 300
 GAGCTGGCCA ACCGCACCCG CCTCCTGTCT GTCTGTGATG AGCGCTGGGC AGACGACCTG 360
 TACCCCAAGA TCTACCACAG CTGCTTCTTC ATTGTCACCT ACCTGGCCCC ACTGGGCCTC 420
 ATGGCCATGG CCTATTTCCA GATCTTCCGC AAGCTCTGGG GCCGCCAGAT CCCCGGCACC 480
 ACCTCGGCCC TGGTGCGCAA CTGGAAGCGG CCCTCAGACC AGCTGGACGA CCAGGGCCAG 540
 GGCCTGAGCT CAGAGCCCCA GCCCCGGGCT CGCGCCTTCC TGGCCGAGGT GAAACAGATG 600
 CGAGCCCGGA GGAAGACGGC CAAGATGCTG ATGGTGGTGC TGCTGGTCTT CGCCCTCTGC 660
 TACCTGCCCCA TCAGTGTCTT CAACGTCTC AAGAGGGTCT TCGGGATGTT CCGCCAAGCC 720
 AGCGACCGAG AGGCCATCTA CGCCTGCTTC ACCTTCTCCC ACTGGCTGGT GTACGCCAAC 780
 AGCGCCGCC 789

(2) INFORMATION FOR SEQ ID NO: 56:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 328
 (B) TYPE: Amino acid
 (C) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 56:

Met Glu Trp Asp Asn Gly Thr Gly Gln Ala Leu Gly Leu Pro Pro Thr
 1 5 10 15
 Thr Cys Val Tyr Arg Glu Asn Phe Lys Gln Leu Leu Leu Pro Pro Val
 20 25 30
 Tyr Ser Ala Val Leu Ala Ala Gly Leu Pro Leu Asn Ile Cys Val Ile
 35 40 45
 Thr Gln Ile Cys Thr Ser Arg Arg Ala Leu Thr Arg Thr Ala Val Tyr
 50 55 60
 Thr Leu Asn Leu Ala Leu Ala Asp Leu Leu Tyr Ala Cys Ser Leu Pro
 65 70 75 80
 Leu Leu Ile Tyr Asn Tyr Ala Gln Gly Asp His Trp Pro Phe Gly Asp
 85 90 95
 Phe Ala Cys Arg Leu Val Arg Phe Leu Phe Tyr Ala Asn Leu His Gly
 100 105 110

- 2 6 8 -

Ser Ile Leu Phe Leu Thr Cys Ile Ser Phe Gln Arg Tyr Leu Gly Ile
 115 120 125
 Cys His Pro Leu Ala Pro Trp His Lys Arg Gly Gly Arg Arg Ala Ala
 130 135 140
 Trp Leu Val Cys Val Thr Val Trp Leu Ala Val Thr Thr Gln Cys Leu
 145 150 155 160
 Pro Thr Ala Ile Phe Ala Ala Thr Gly Ile Gln Arg Asn Arg Thr Val
 165 170 175
 Cys Tyr Asp Leu Ser Pro Pro Ala Leu Ala Thr His Tyr Met Pro Tyr
 180 185 190
 Gly Met Ala Leu Thr Val Ile Gly Phe Leu Leu Pro Phe Ala Ala Leu
 195 200 205
 Leu Ala Cys Tyr Cys Leu Leu Ala Cys Arg Leu Cys Arg Gln Asp Gly
 210 215 220
 Pro Ala Glu Pro Val Ala Gln Glu Arg Arg Gly Lys Ala Ala Arg Met
 225 230 235 240
 Ala Val Val Val Ala Ala Ala Phe Ala Ile Ser Phe Leu Pro Phe His
 245 250 255
 Ile Thr Lys Thr Ala Tyr Leu Ala Val Gly Ser Thr Pro Gly Val Pro
 260 265 270
 Cys Thr Val Leu Glu Ala Phe Ala Ala Ala Tyr Lys Gly Thr Arg Pro
 275 280 285
 Phe Ala Ser Ala Asn Ser Val Leu Asp Rro Ile Leu Phe Tyr Phe Thr
 290 295 300
 Gln Lys Lys Phe Arg Arg Arg Pro His Glu Leu Leu Gln Lys Leu Thr
 305 310 315 320
 Ala Lys Trp Gln Arg Gln Gly Arg
 325

(2) INFORMATION FOR SEQ ID NO: 57:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 984
- (B) TYPE: Nucleic acid
- (C) STRANDEDNESS: Double
- (D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE

- (C) IDENTIFICATION METHOD: S

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 57:

- 2 6 9 -

ATGGAATGGG ACAATGGCAC AGGCCAGGCT CTGGGCTTGC CACCCACCAC CTGTGTCTAC 60
 CGCGAGAACT TCAAGCAACT GCTGCTGCCA CCTGTGTATT CGGCGGTGCT GGC GGCTGGC 120
 CTGCCGCTGA ACATCTGTGT CATTACCCAG ATCTGCACGT CCCGCCGGGC CCTGACCCGC 180
 ACGGCCGTGT ACACCCTAAA CCTTGCTCTG GCTGACCTGC TATATGCCTG CTCCTGCCC 240
 CTGCTCATCT ACAACTATGC CCAAGGTGAT CACTGGCCCT TTGGCGACTT CGCCTGCCGC 300
 CTGGTCCGCT TCCTCTTCTA TGCCAACCTG CACGGCAGCA TCCTCTTCCT CACCTGCATC 360
 AGCTTCCAGC GCTACCTGGG CATCTGCCAC CCGCTGGCCC CCTGGCACAAC ACGTGGGGGC 420
 CGCCGGGCTG CCTGGCTAGT GTGTGTAACC GTGTGGCTGG CCGTGACAAC CCAGTGCTG 480
 CCCACAGCCA TCTTCGCTGC CACAGGCATC CAGCGTAACC GCACTGTCTG CTATGACCTC 540
 AGCCCGCCTG CCCTGGCCAC CCACTATATG CCCTATGGCA TGGCTCTCAC TGTCATCGGC 600
 TTCCTGCTGC CCTTTGCTGC CCTGCTGGCC TGCTACTGTC TCCTGGCCTG CCGCCTGTGC 660
 CGCCAGGATG GCCCGGCAGA GCCTGTGGCC CAGGAGCGGC GTGGCAAGGC GGCCCGCATG 720
 GCCGTGGTGG TGGCTGCTGC CTTTGCCATC AGCTTCCTGC CTTTTCACAT CACCAAGACA 780
 GCCTACCTGG CAGTGGGCTC GACGCCGGGC GTCCCCTGCA CTGTATTGGA GGCCTTTGCA 840
 GCGGCCTACA AAGGCACGCG GCCGTTTGCC AGTGCCAACA GCGTGCTGGA CCCCATCCTC 900
 TTCTACTTCA CCCAGAAGAA GTTCCGCCGG CGACCACATG AGCTCCTACA GAAACTCACA 960
 GCCAAATGGC AGAGGCAGGG TCGC 984

(2) INFORMATION FOR SEQ ID NO: 58:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26
 (B) TYPE: Nucleic acid
 (C) STRANDEDNESS: Single
 (D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Other nucleic acid
Synthetic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 58:

ACAGCCATCT TCGCTGCCAC AGGCAT 26

(2) INFORMATION FOR SEQ ID NO: 59:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 29
 (B) TYPE: Nucleic acid
 (C) STRANDEDNESS: Single

- 2 7 0 -

(D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Other nucleic acid
Synthetic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 59:

AGACAGTAGC AGGCCAGCAG GGCAGCAAA 29

(2) INFORMATION FOR SEQ ID NO: 60:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 27
(B) TYPE: Nucleic acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear(ii) MOLECULE TYPE: Other nucleic acid
Synthetic DNA

(iii) FEATURES: N is inosine

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 60:

CTGTGYGYSA TYGCNNTKGA YMGSTAC 27

(2) INFORMATION FOR SEQ ID NO: 61:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 29
(B) TYPE: Nucleic acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear(ii) MOLECULE TYPE: Other nucleic acid
Synthetic DNA

(iii) FEATURES: N is inosine

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 61:

AKGWAGWAGG GCAGCCAGCA GANSRYGAA 29

CLAIMS

1. A DNA which comprises a nucleotide sequence represented by a SEQ ID NO selected from the group consisting of SEQ ID NO: 1 to SEQ ID NO: 19.

5 2. A method for amplifying a DNA coding for a G protein coupled receptor protein by polymerase chain reaction techniques, which comprises:

(i) carrying out a polymerase chain reaction in the presence of a mixture of

- 10 ① a DNA coding for a G protein coupled receptor protein, said DNA being capable of acting as a template,
- ② at least one DNA primer selected from the group consisting of DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 1, DNA primers comprising a nucleotide sequence represented by SEQ ID
- 15 NO: 3, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 5, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 6, DNA primers comprising a nucleotide sequence represented by
- 20 SEQ ID NO: 7, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 10, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 14, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 16 and DNA primers comprising
- 25 a nucleotide sequence represented by SEQ ID NO: 18, and
- ③ at least one DNA primer selected from the group consisting of DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 2, DNA primers comprising a nucleotide sequence represented by SEQ ID
- 30 NO: 4, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 8, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 9, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 11, DNA primers comprising a nucleotide

sequence represented by SEQ ID NO: 15, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 17 and DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 19; or

5 (ii) carrying out a polymerase chain reaction in the presence of a mixture of

- ① a DNA coding for G protein coupled receptor protein, said DNA being capable of acting as a template,
- ② at least one DNA primer selected from the group
10 consisting of DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 1 and DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 12, and
- ③ at least one DNA primer selected from the group
15 consisting of DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 13.

3. A method for screening a DNA library for a DNA coding for a G protein coupled receptor protein, which comprises:

- 20 (i) carrying out a polymerase chain reaction in the presence of a mixture of
- ① said DNA library,
 - ② at least one DNA primer selected from the group
25 consisting of DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 1, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 3, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 5, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 6, DNA
30 primers comprising a nucleotide sequence represented by SEQ ID NO: 7, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 10, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 14, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 16 and DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 18, and
35
 - ③ at least one DNA primer selected from the group

consisting of DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 2, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 4, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 8, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 9, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 11, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 15, DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 17 and DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 19,

under conditions to amplify selectively a template DNA coding for the G protein coupled receptor protein, contained in the DNA library and selecting said DNA; or

(ii) carrying out a polymerase chain reaction in the presence of a mixture of

① said DNA library

② at least one DNA primer selected from the group consisting of DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 1 and DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 12, and

③ at least one DNA primer selected from the group consisting of DNA primers comprising a nucleotide sequence represented by SEQ ID NO: 13,

under conditions to amplify selectively a DNA coding for the G protein coupled receptor protein, contained in the DNA library and selecting said DNA.

4. A DNA coding for a G protein coupled receptor protein or a fragment thereof, which is obtained by the method according to claim 2 to 3.

5. A G protein coupled receptor protein encoded by the DNA according to claim 4, a peptide segment or fragment thereof or a salt thereof.

6. A G protein coupled receptor protein comprising an amino acid sequence selected from the group consisting of

an amino acid sequence represented by SEQ ID NO: 24, an amino acid sequence represented by SEQ ID NO: 25, an amino acid sequence represented by SEQ ID NO: 26, an amino acid sequence represented by SEQ ID NO: 27, an amino acid sequence represented by SEQ ID NO: 28, an amino acid sequence represented by SEQ ID NO: 34, an amino acid sequence represented by SEQ ID NO: 35, an amino acid sequence represented by SEQ ID NO: 38, an amino acid sequence represented by SEQ ID NO: 39, an amino acid sequence represented by SEQ ID NO: 56, and substantial equivalents to the amino acid sequence represented by SEQ ID NO: 24, SEQ ID NO: 25, SEQ ID NO: 26, SEQ ID NO: 27, SEQ ID NO: 28, SEQ ID NO: 34, SEQ ID NO: 35, SEQ ID NO: 38, SEQ ID NO: 39, or SEQ ID NO: 56; a peptide segment (or fragment) thereof, a modified peptide derivative thereof or a salt thereof.

7. The G protein coupled receptor protein according to claim 6, comprising an amino acid sequence selected from the group consisting of an amino acid sequence represented by SEQ ID NO: 38, an amino acid sequence represented by SEQ ID NO: 39, an amino acid sequence represented by SEQ ID NO: 56 and substantial equivalents to the amino acid sequence represented by SEQ ID NO: 38, SEQ ID NO: 39, or SEQ ID NO: 56.

8. The G protein coupled receptor protein according to claims 6 or 7, wherein said receptor is a purinoceptor.

9. The G protein coupled receptor protein according to any of claims 6 to 8, wherein an agonist to said receptor is useful as an immunomodulator or an antitumor agent, in addition it is useful in therapeutically or prophylactically treating hypertension, diabetes or cystic fibrosis, and an antagonist to said receptor is useful as a hypotensive agent, an analgesic, or an agent for therapeutically or prophylactically treating incontinence of urine.

10. A DNA which comprises a nucleotide sequence coding for a G protein coupled receptor protein of claim 6.

11. The DNA according to claim 10 comprising a nucleotide sequence coding for the G protein coupled receptor

protein according to claim 7.

12. The DNA according to claim 11 comprising a nucleotide sequence represented by SEQ ID NO: 40, SEQ ID NO: 41, or SEQ ID NO: 57.

5 13. A transformant containing a vector comprising the DNA according to claim 4 or 10; or an expression system comprising an open reading frame (ORF) of DNA derived from a G protein coupled receptor protein DNA according to claim 4 or 10, wherein the ORF is operably linked
10 to a control sequence compatible with a desired host cell.

14. A method for determining a ligand to the G protein coupled receptor protein according to any of claims 5 to 8, which comprises contacting

15 (i) at least one component selected from the group consisting of G protein coupled receptor proteins or salts thereof according to any of claims 5 to 8, peptide segments or salts thereof, and mixtures thereof,
with

20 (ii) at least one compound to be tested and determining whether said compound to be tested bound to the component of (i).

25 15. A screening method for a compound capable of inhibiting the binding of a G protein coupled receptor protein according to any of claims 5 to 8 with a ligand, which comprises carrying out a comparison between:

30 (i) at least one case where said ligand is contacted with at least one component selected from the group consisting of G protein coupled receptor proteins or salts thereof according to any of claims 5 to 8, peptide segments or salts thereof, and mixtures thereof,
and

35 (ii) at least one case where said ligand together with a compound to be tested is contacted with at least one component selected from the group consisting of G protein coupled receptor proteins or salts thereof according to any of claims 5 to 8, peptide segments or salts thereof, and mixtures thereof.

16. A compound which is determined through the method according to claim 15 or a salt thereof.

17. The compound according to claim 16, which is an agonist or antagonist to a G protein coupled receptor protein
5 according to any of claims 5 to 8.

18. A ligand to a G protein coupled receptor protein according to any of claims 5 to 8, which is determined through the method according to claim 14.

10

15

1 / 7 9

FIGURE 1

OLIGODEOXYNUCLEOTIDE SEQUENCE FOR 5' SIDE PRIMER

Primer HS-1

CGTGGCCATCCTGGGCAACACCCTG

G C G G

C T

G

T

HTRHR	CCTGGGCATTGTAGGCAACATCATGGT
HUMRANTES	CATTGGCCTGGTTGGAAACATCCTGGT
HSBLR1A	CCTGGGCGTGATCGGCAACGTCCTGGT
HUMSOMAT	GGTGGGGCTGGTGGGCAACGCCCTGGT
RNU02083	AGTGGGCCTCTTCGGAAACTTCCTGGT
U00442	GGTGGGCTTAGTGGGCAATTCCTGGT
HUMNMBR	CGTGGGCTTGCTGGGCAACATCATGCT
HSHM4	GGTGACCATCATCGGCAACATCCTGGT
RATAADRE01	CTTTGCCATCGTGGGCAACATCTTGGT
HUMSSTR3X	GGTGGGCCTGCTGGGTAACCTCGCTGGT
HUMC5AAR	GGTGGGAGTGCTGGGCAATGCCCTGGT
HUMRDC1A	CATCGGCATGATTGCCAACTCCGTGGT
HUMOPIODRE	CGTGGCGGTGCTCGGCAACCTCGTGGT
RATA2BAR	GCTGGCAGTGGCGGGCAACGTGCTGGT

2 / 7 9

FIGURE 2

OLIGODEOXYNUCLEOTIDE SEQUENCE FOR 3' SIDE PRIMER

Complementary Sequence
to Primer HS-2

TTTGCCATCTGCTGGATGCCCCACAAC

C	C	TTT	C
	G		G
	T		T

HUMSGIR

TTTGCCCTCTGCTGGTTCCCTCTCAAC

HUMBOMB3S

TTTGCCCTCTGCTGGTTGCCAAATCAC

S46950

TTTGCCCTCTGCTGGCTGCCCCTACAC

MUSGPCR

TTTGCCCTCGTCTGGTGCCCTCTCAAC

S43387

TTTGCCCTTTTATGGATGCCCTACAGG

RATNEURA

TTTGCCATCTGCTGGCTGCCCTATCAC

RATA1ARA

TTTGCCCTCAGCTGGCTGCCGCTGCAT

HUMOPIODRE

TTTGCCATCTGCTGGCTGCCCTATCAC

HUMNEKAR

TTTGCCATCTGCTGGCTGCCCTACCAC

RATADENREC

TTTGCCCTTGCTGGCTGCCTTTGTCC

HUMSRI1A

TTTGTCATCTGCTGGATGCCTTTCTAC

S8637154

TTTGCTATCTGCTGGCTGCCCTATCAT

RNCGPCR

TTTGCCGCCTGCTGGATGCCTTTTACC

HUMSSTR4Z

TTTGCTGCTCTGCTGGATGCCTTTCTAC

RATGNRHA

TTTGCACTGCTCGAAGCCAGACAAA

3 / 7 9

FIGURE 3

OLIGODEOXYNUCLEOTIDE SEQUENCE FOR 5' SIDE PRIMER

Primer 3A CTGACCGCTCTIACIACTGACCGATAC
 T T GG GT A C
 G

Primer 3B CTGACCGCTCTIACIACTGACCGATAT
 T T GG GT A C
 G

L11064	CTCACCATGATGAGCGTGGACCGCTAC
L11065	TTGACCATGATGGAGTGTGACCGCTAC
D16349	CTCTGCACCATGAGCGTGGACCGCTAC
X69676	CTGATGCTCGTGAGTATCGACCGCTAC
M35328	CTTACGGCACTGTCAGCTGACAGGTAC
M73482	CTCACTGCCCTCAGCGCCGACAGGTAC
M73481	CTCACGGCGCTCTCGGCAGACAGATAC
L08893	TTAACAATTCTCAGCGCTGACAGATAC
X62933	ATGACCGCCATCGCCGCTGACAGGTAC
X62934	ATGACAACTGTGGCCTTTGACAGATAC
J05189	ATGACAGCCATTGCAGTGGACAGGTAT
M60786	CTCTGCGCTCTCAGTGTGGACAGGTAC
L04672	CTCACCTGCCTCAGCATTGACCGCTAC
X61496	TTGCTGGCTATCACTGTGGACCGCTAC
X59249	TTGCTGGCCATTGCTGTAGACCGATAC
L09249	CTCACCTGCCTCAGCATTGACCGCTAC
P30731	CTGACAGCTATCGCAGTGGACCGCCAC
M31210	CTCCTCGCCATCGCCATTGAGCGCTAT
U03642	CTCACCGGCCTCAGCTTCGACCGCTAC

4 / 7 9

FIGURE 4

OLIGODEOXYNUCLEOTIDE SEQUENCE FOR 5' SIDE PRIMER

Primer 3C	CTCGCCGCTATIAGCATGGACCGITAC
	G CC G T T

Primer 3D	CTCGCCGCTATIAGCATGGACCGITAT
	G CC G T T

L32840	ATTACCTGCATGAGTGTGATAGGTAC
X64052	CTCACGTGTCTCAGCATCGATCGCTAC
M90065	CTCACGTGTCTCAGCATCGATCGCTAC
M91464	CTCACGTGTCTCAGCATTGATCGATAC
M88096	CTGGTAGCCATCTCTCTGGAGAGATAT
M99418	CTCGTGGCCATAGCCCTGGAGCGATAC
L04473	CTCGTGGCCATCGCACTGGAGCGGTAC
M73969	CTGGCCTGCATCAGTGTGGACCGTTAC
X65858	TTGGCCTGCATCAGTGTGGACCGTTAC
S46665	CTGGCTACCATTAGTGCCGACCGTTTC
M60626	ATCGCCCTCATTGCTCTGGACCGCTGT

FIGURE 5

OLIGODEOXYNUCLEOTIDE SEQUENCE FOR 3' SIDE PRIMER

Complementary Sequence TTTACCITCTGTTGGICGCCCTACCACATC
to Primer 6A GT TC T T

Complementary Sequence TTCACCITCTGTTGGICGCCCTACCACATC
to Primer 6B GT TC T T

L11064	TTCGTGGTGTGCTGGGCGCCCATCCACATC
L11065	TTCATCATCTGTTGGACCCCCATTACATC
D16349	TTTATCGTCTGCTGGACCCCCATCCACATC
X69676	TTTGTGCTGTGTTGGGTGCCTTTCCAGATC
M35328	TTTGCCTTCTGCTGGCTCCCCAACCATGTC
M73482	TTCATCTTCTGTTGGTTTCCAAACCACATC
M73481	TTGCGCTTCTGCTGGCTCCCCAATCATGTC
L08893	TTTGCCCTCTGCTGGTTGCCAAATCACCTC
X62933	TTTGCCATCTGCTGGCTGCCCTACCACCTC
X62934	TTGCCATCTGCTGGCTGCCCTTCCACATC
J05189	TTTGCCATCTGCTGGCTGCCCTATCACGTG
M60786	TTGCGCCTGTGCTGGTTCCCTCTTCACTTA
L04672	TTTGTCACTCTGCTGGCTGCCCTACCACGTG
X61496	TTTGGCGCCTGCTGGATGCCTTTTACCCTC
X59249	TTTGCCTTGTGCTGGCTGCCTTTGTCCATC
L09249	TTTGCCATCTGCTGGCTGCCCTACCACGTG
P30731	TTTGCCCTCTGCTGGTTCCCTCTCAACTGC
M31210	TTCATCGCCTGCTGGGCACCGCTCTTCATC
U03642	TTTGCCCTGTGCTGGATGCCCTACCACCTG

FIGURE 6

OLIGODEOXYNUCLEOTIDE SEQUENCE FOR 3' SIDE PRIMER

Complementary Sequence TTTTTCITTTGCTGGITTCCTACCATG
to Primer 6C CC T G C T T

L32840	TTCATCATTTGCTGGCTTCCCTTCCATGTT
X64052	TTCTTCTTTTCCTGGGTCCCCACCAAATA
M90065	TTCTTCTTTTCCTGGGTCCCCACCAAATA
M91464	TTTTTCTTTTCCTGGATTCCCCACCAAATA
M88096	TTCTTCTGTGCTGGATGCCCATCTTCAGC
M99418	TTCTTCTGTGTTGGCTGCCAGGTGTACAGC
L04473	TTTTTCTGTGTTGGTTGCCAGTTTATAGT
M73969	TTCTTGCTTTGCTGGCTGCCCTACAACCTG
X65858	TTCTTGCTTTGCTGGCTGCCCTACAACCTG
S46665	TTCTTTATCTTCTGGCTGCCCTATCAGGTG
M60626	TTTTTCTCTGCTGGTCCCCATATCAGGTG

7 / 7 9

FIGURE 7

OLIGODEOXYNUCLEOTIDE SEQUENCE FOR 5' SIDE PRIMER

Primer T2A

GTCACCAACITGTTTCATCCTCAICCTG

C	AC	GT T
		A

HUMGALAREC	ACCACCAACCTGTTTCATCCTCAACCTG
RATADRA1B	CCCACCAACTACTTTATCGTCAACCTG
HUMADRB1	ACCACCAACCTGTTTCATCCTCAACCTG
RABIL8RSB	GTCACCGACGTCTACCTGCTGAACCTG
HUMOPIODRE	GTCACCAACTCCTTCCTCGTGAACCTG
BTSKR	GTGACCAACTACTTCATCGTCAACCTG
HUMSRI2A	ATCACCAACATTTACATCCTCAACCTG
HUMSSTR3Y	GTCACCAACGTCTACATCCTCAACCTG
HUMGARE	GTCACCAACGCCTTCCTCCTCTCACTG
HUMCCKAR	GTCACCAACATCTTCCTCCTCTCCCTG
HUMSHTR	CCCTCCAACCTACCTGATCGTGTCCCTG
HUMD1B	ATGACCAACGTCTTCATCGTGTCTCTG
HUM5HT1E	CCTGCCAACTACCTAATCTGTTCTCTG
HUMD4C	CCCACCAACTCCTTCATCGTGAGCCTG
MMSERO	GCCACCAACTATTTCTGATGTCACTT
RATADRA1A	GTCACCAACTATTTTCATCGTGAACCTG
S57565	CTGACCAATTGCTTCATTGTGTCCCTG

FIGURE 8

COMPLEMENTARY OLIGODEOXYNUCLEOTIDE SEQUENCE TO 3' SIDE PRIMER

Complementary Sequence	AACCCCTCITCTATTGCTTTITCICT
to Primer T7A	T T C C C G G

HUMGALAREC	AATCCTATCATTTATGCATTTCTCTCT
RATA1ADREC	AACCCCATCGTCTATGCCTTCCGGATC
PIGA2R	AATCCTCTCTTTTATGGCTTTCTGGGG
RAT5HTRTC	AACCCTATCATCTACCCGCTCTTTATG
S58541	AACCCCATCATTTATGCCTTTAATGCT
HUMGRPR	AACCCCTTTGCCCTCTACCTGCTGAGC
MUSGRPBOM	AACCCCTTTGCTCTTTATCTGCTGAGC
RRVT1AIIR	AACCCTCTGTTCTACGGCTTTCTGGGG
HUMADRB1	AACCCCATCATCTACTGCCGCAGCCCC
HSHM4	AACCCCGTGTGCTATGCTCTGTGCAAC
HUMGARE	AACCCCTGGTCTACTGCTTCATGCAC
RATCCKAR	AACCCCATCATCTATTGCTTCATGAAC
S59749	AATCCCATGCTCTACACCTTCGCTGGC
HUMSST28A	AACCCCGTCCTCTACGGCTTCCTCTCG
RNGPROCR	AACCCCATCCTCTACGGCTTCCTCTCC
MUSSSRI1A	AACCCCATACTCTACGGCTTCCTGTGC
HUMA1AADR	AACCCGCTCATCTACCCCTGTTCCAGC
S66181	AACCCGGTTCTCTACGCCTTCCTGGAC
HUMSSTR3Y	AACCCCATCCTTTATGGCTTCCTCTCC

9 / 7 9

FIGURE 9

OLIGODEOXYNUCLEOTIDE SEQUENCE FOR 5' SIDE PRIMER

Primer TM1-A2 TGITGGTTATIGGIGTTGTIGGIAA
 CC GC C G

MUSBB2R	TGGTGGTGGTGGTGGTGGTGGGCAA
BTSKR	TGGTGCTGGTGGCTGTGATGGGCAA
BOVEETBR	TGTTTCGTGCTGGGCATCATCGGAAA
HUMNEUYREC	TGATCATTCTTGGTGTCTCTGGAAA
MMSUBKREC	TGGTGCTGGTGGCTGTAACAGGCAA
HUMPGE2R	TGTTTCATCTTCGGGGTGGTGGGCAA
HUMPIR	TGTTTCGTGGCCGGTGTGGTGGGCAA
HSU11053	TGTTTCGTGCTGGGCTTGGTGGGCAA
RRMC3RA	TGGTGATCCTGGCTGTGGTGAGGAA
HUMMR	TGGTTATCCTGGCCGTGGTCAGGAA
MUSGRPBO	TCATCGTGATAGGTCTTATTGGCAA
RATCHOLREC	TCTTTCTGATGAGTGTTGGCGGAAA
RATCCKAR	TATTCCTTCTCAGTGTGCGGGGAA

1 0 / 7 9

FIGURE 10

COMPLEMENTARY OLIGODEOXYNUCLEOTIDE SEQUENCE TO 3' SIDE PRIMER

Complementary Sequence	GCCAT	IACCITGGACAGATACCGAT
to Primer TM3-B2	A	T A C G A G

HUMCCKR	GCCATCGCACTGGAGCGGTACAG
HUMCCKBGR	GCCATCGCACTGGAGCGGTACAG
MMGMC5R	GCCATTGCGGTGGACAGGTACA
HUMV2R	GCCATGACGCTGGACCGCCACCG
RATNEURA	GCCATTGCAGTGGACAGGTA
DOGGSTRN	GCCATCGCCCTGGAGCGATACAG
RAT5HT5A	GCAATAGCTTTGGACCGCTACTGGT
MUSALP2ADA	GCCATTAGTCTGGACCGCTACTGGT
HUMADORA1X	GCAATTGCTGTGGACCGCTACC
HUMOPIODRE	GCCATCGCGGTGGACAGATACA
MUSGRPBOM	GCACTGTCAGCTGACAGGTACAAA
RATCCKAR	GCCATCTCTCTGGAGAGATATGG
HSTRHREC	GCCTTTACCATTGAGAGGTACATA

1 1 / 7 9

FIGURE 11

OLIGODEOXYNUCLEOTIDE SEQUENCE FOR 5' SIDE PRIMER

Primer TM3-C2

CATGGCCGTGGAGAGITACITGGC

TT C C T A

HUMNK3R

CATTGCGGTGGACAGGTATATGGC

HSMRNOXY

CATGTCCCTGGACCGCTGCCTGGC

S68242

CATATCGCTGGAGAGATACGGAGC

CFGPCR4

CATCGCTCTGGACAGGTACTGGGC

MMSUBPREC

TGGCCTTTGACAGATACATGGC

HUMOPIODRE

CATCGCGGTGGACAGATACATGGC

HUMGALAREC

ATGTCCGTGGACCGCTACGTGGC

HSS31G

CATTGCCCTGGACAGGTACTGGGC

HUMARB3A

CCTGGCCGTGGACCGCTACCTGGC

HUMHPR

CATGGCCGTGGAGCGCTGCCTGGC

RATCCKAR

CATCTCTCTGGAGAGATATGGCGC

1 2 / 7 9

FIGURE 12

COMPLEMENTARY OLIGODEOXYNUCLEOTIDE SEQUENCE TO 3' SIDE PRIMER

Complementary Sequence	TTTGCCITCTGCTGGATCCCCAAC
to Primer TM6-E2	C G C G TT

HUMNEKAR	TTTGCCATCTGCTGGCTGCCCTAC
HUMSUBPRA	TTCGCCATCTGCTGGCTGCCCTTC
RATSKR	TTTGCCATCTGCTGGCTGCCCTAC
MUSGRPBOM	TTTGCCCTTCTGCTGGCTCCCCAAC
HUMOPIODRE	TTTGCCATCTGCTGGCTGCCCTA
HUMA2XXX	TTTGCCCTCTGCTGGCTGCCCT
HUMADRBR	TTCACCCTCTGCTGGCTGCCCTTC
CFGPCR8	TTCGCCCCCTCTGTGGCTGCCCT
HUMETSR	TTTGCCCTCTGCTGGCTTCCCCT
MMNPY1CDS	TTCGCCGTCTGCTGGCTGCCCT
HSMRNAOXY	TTCATCGTGTGCTGGACGCCTTTC
RATCCKAR	TTCTTCCTGTGCTGGATGCCCATC

1 3 / 7 9

FIGURE 13

OLIGODEOXYNUCLEOTIDE SEQUENCE FOR 5' SIDE PRIMER

Primer TM2F18

ARYYTIGCIITIGCNGAY

HUMTSHX	AACCTGGCCTTTGCGGAT
HUMNEKAR	AATCTGGCGCTGGCTGAC
HUMFMLP	AACCTGGCCGTGGCTGAC
HUMINTLEU8	AACCTAGCCTTGGCCGAC
HUMA1AADR	AACCTGGCCGTGGCCGAC
HUMIL8RA	AACCTGGCCTTGGCCGAC
HSDD2	AGCCTCGCAGTGGCCGAC
HUMANTIR	AATTTAGCACTGGCTGAC
HUMSOMAT	AACCTGGCCGTAGCCGAC
HUMEL4REC	AGCTTGGCTGTGGCTGAT
HSTRHREC	AGCCTGGCAGTAGCTGAT
HSU07882	AACCTGGCCTTAGCCGAT

(R = A or G, Y = C or T, N = A, C, G or T, and
I = Inosine)

1 4 / 7 9

FIGURE 14

COMPLEMENTARY OLIGODEOXYNUCLEOTIDE SEQUENCE TO 3' SIDE PRIMER

Complementary Sequence TTYNNNTNTGYTGGITICCI
to Primer TM6R21

HSBAR	TTCACCCTCTGCTGGCTGCCC
HUMNEKAR	TTTGCCATCTGCTGGCTGCCC
HUMETN1R	TTTGCTCTTTGCTGGTTCCCT
HUMHISH2R	TTCATCATCTGCTGGTTTCCC
HUMA1AADR	TTCGTGCTCTGCTGGTTCCCT
HUMIL8RA	TTCTGCTTTGCTGGCTGCCC
HUMNMBR	TTCATCTTCTGTTGGTTTCCT
HUMNKIRX	TTGCCATCTGCTGGCTGCCC
HUMSUBPRA	TTGCCATCTGCTGGCTGCCC
HUM5HT1DA	TTTATCATCTGCTGGCTGCCC
HUMPFPR2A	TTCTTCATCTGTTGGTTTCCC
HSDD2	TTCATCATCTGCTGGCTGCCC
HUMNEUYREC	TTTGCACTCTGCTGGCTCCCT
HUM2XXX	TTTGCCCTCTGCTGGCTGCCC
HUMBK2A	TTCATCATCTGCTGGCTGCCC
HUMFMLPX	TTCTTCATCTGTTGGTTTCCC
HUMSSTR3X	TTCGTGCTCTGCTGGATGCCC
HUMCCKR	TTTTTTCTGTGTTGGTTGCCA
HSNEURA	TTTGTGGTCTGCTGGCTGCCC

(Y = C or T, N = A, C, G or T, and I = Inosine)

1 5 / 7 9

FIGURE 15

OLIGODEOXYNUCLEOTIDE SEQUENCE FOR 5' SIDE PRIMER

Primer S3A	GCCTGITIAIGATGAGTGTGGAIAGIT
	C G C TC C

HUMGALAREC	CCCTGGCCGCGATGTCCGTGGACCGCT
S70057	GCCTCGTGGCCATCGCACTGGAGCGGT
S67127	ACCTCTGCGCTCTTAGTGTTGACAGGT
S44866	GTCTATGTGCTCTGAGTATTGACAGAT
HUMC5AAR	TCCTGGCCACCATCAGCGCCGACCGCT
HUMANTIR	TACTCACGTGTCTCAGCATTGATCGAT
HUMBK2A	TCCTGATGCTGGTGAGCATCGACCGCT
HSNEURA	ACGTGGCCAGCCTGAGTGTGGAGCGCT
HUMGRPR	CACTCACGGCGCTCTCGGCAGACAGAT
HUMFSRS	GCCTGACAGTCATGAGCGTGGACCGCT
HUMIL8RA	TGTTGGCCTGCATCAGTGTGGACCGTT
HUMNEKAR	CCATGACCGCCATTGCTGCCGACAGGT

1 6 / 7 9

FIGURE 16

COMPLEMENTARY OLIGODEOXYNUCLEOTIDE SEQUENCE TO 3' SIDE PRIMER

Complementary Sequence	TGGITICCTACCACITIAATCAICATC
to Primer S6A	T T GG GT

HUMGALAREC	TGGCTGCCGCACCACATCATCCATCTC
S70057	TGGTTGCCAGTTTATAGTGCCAACACG
S67127	TGGTTCCCTCTTCATTTAAGCCGTATA
S44866	TGGCTTCCCTTCACCTCAGCAGGATT
HUMC5AAR	TGGTTGCCCTACCAGGTGACGGGGATA
HUMANTIR	TGGATTCCCCACCAAATATTCACTTTT
HUMBK2A	TGGCTGCCCTTCCAGATCAGCACCTTC
HSNEURA	TGGACTCCGTTCTCTATGACTTCTAC
HUMGRPR	TGGCTCCCCAATCATGTCATCTACCTG
HUMFSRS	TGGCTGCCCTTCTTCACCGTCAACATC
HUMIL8RA	TGGCTGCCCTACAACCTGGTCCTGCTG
HUMNEKAR	TGGCTGCCCTACCACCTCTACTTCATC

17/79

FIGURE 17



FIGURE 18

	10	20	30	40	50
A58-T7-2	GTGGGCATG	TGGGCAAC	CCCTGGT	CATCTTC	TGTCGATCCTTCGCTACGC
HUMSOMAT	X:::	:::	:::	:::	:::
	GTGGGCTG	TGGGCAAC	CGCCCTGGT	CATCTTC	TGTCGATCCTTCGCTACGC
	285	295	305	315	325
A58-T7-2	60	70	80	90	100
HUMSOMAT	CAAGATGA	AGACGGCT	TACCAACA	TCTACCT	TGCTCAACCTGGCCGTAGCCG
	:::	:::	:::	:::	:::
	CAAGATGA	AGACGGCT	TACCAACA	TCTACCT	TGCTCAACCTGGCCGTAGCCG
	335	345	355	365	375
A58-T7-2	110	120	130	140	150
HUMSOMAT	ACGAGCTC	TTCATGCT	AGCGTG	CGCCCTTC	TGTCGGCCTCGTCGGCCGCCCTG
	:::	:::	:::	:::	:::
	ACGAGCTC	TTCATGCT	AGCGTG	CGCCCTTC	TGTCGGCCTCGTCGGCCGCCCTG
	385	395	405	415	425
A58-T7-2	160	170	180	190	200
HUMSOMAT	CGCCACTG	GGCCCTTC	GGCTCCG	TGCTGTG	CCGCGGTCGTCAGCGTCGA
	:::	:::	:::	:::	:::
	CGCCACTG	GGCCCTTC	GGCTCCG	TGCTGTG	CCGCGGTCGTCAGCGTCGA
	435	445	455	465	475
A58-T7-2	210	220	230	240	
HUMSOMAT	CGGCCTCA	ACATGTT	CACACG	GTCTTCT	TGTCACCGTGCTCAGCGT
	:::	:::	:::	:::	:::
	CGGCCTCA	ACATGTT	CACACG	GTCTTCT	TGTCACCGTGCTCAGCGT
	485	495	505	515	

2 0 / 7 9

FIGURE 20

	10	20	30	40	50
57-A-2	<u>GTGGGCATGCTGGGCAACCTCCTGGAAGGCAGTCGCCGAGGTGGCCGGTT</u>				
HUMDRD5A	X::: ::::: :: : ::				
	424	434	444	454	
	60	70	80	90	100
57-A-2	ACTGGCCCTTTGGAGCGTTCTGCGACGTCTGGGTGGCCTTCGACATCATG				
HUMDRD5A	::				
	464	474	484	494	504
	110	120	130	140	150
57-A-2	TGCTCCACTGCCTCCATCCTGAACCTGTGCGTCATCAGCGTGGACCGCTA				
HUMDRD5A	::				
	514	524	534	544	554
	160	170	180	190	200
57-A-2	CTGGGCCATCTCCAGGCCCTTCCGCTACAAGCGCAAGATGACTCAGCGCA				
HUMDRD5A	::				
	564	574	584	594	604
	210	220	230	240	250
57-A-2	TGGCCTTGGTCATGGTCGGCCTGGCATGGACCTTGTCCATCCTCATCTCC				
HUMDRD5A	::				
	614	624	634	644	654
	260	270	280	290	300
57-A-2	TTCATTCCGGTCCAGGTCAACTGGGACAGGGACCAGGCGGGCTCTTGGGG				
HUMDRD5A	::::::::::::::::: ::::: ::::::::::::::::::: :::::::::::				
	664	674	684	694	704
	310				
57-A-2	GGGGCTGGACCTGCCAAA				
HUMDRD5A	:::::::::::::X				
	714	724			


2 2 / 7 9

FIGURE 22

5'	9	18	27	36	45	54			
GTG	GGC	ATG	GTG	GGC	ATC	GTG	CGC	CGG	
---	---	---	---	---	---	---	---	---	
Val	Gly	Met	Val	Gly	Asn	Val	Leu	Leu	
63	72	81	90	99	108				
CTG	CAC	AAC	GTG	ACG	AAC	CTG	GGC	ATC	CTC
---	---	---	---	---	---	---	---	---	---
Leu	His	Asn	Val	Thr	Asn	Phe	Leu	Ile	Gly
117	126	135	144	153	162				
ATG	TGC	ACC	GCC	TGC	GTG	ACG	CTC	CTC	COG
---	---	---	---	---	---	---	---	---	---
Met	Cys	Thr	Ala	Cys	Val	Pro	Leu	Thr	Leu
171	180	189	198	207	216				
TGG	GTG	TTC	GGC	GGC	CTG	TGC	CTG	TGC	CTG
---	---	---	---	---	---	---	---	---	---
Trp	Val	Phe	Gly	Gly	Leu	Cys	His	Leu	Val
225	234	243	252	261	270				
GTC	TAT	GTG	TCG	GTG	TTC	ACG	CTC	CTC	CTC
---	---	---	---	---	---	---	---	---	---
Val	Tyr	Val	Ser	Val	Phe	Thr	Leu	Thr	Thr
279	288	297							
GCT	GGT	GCA	CCC	GCT	GAG	GGG	GGG	CAT	3'
---	---	---	---	---	---	---	---	---	---
Ala	Gly	Ala	Pro	Ala	Glu	Ala	Gly	His	

2 4 / 7 9

FIGURE 24



2 5 / 7 9

FIGURE 25





FIGURE 27

	9			18			27			36			45			54		
5'	GTG	GGC	ATG	GTG	GGC	AAC	ATC	CTG	CTG	GTG	CTG	GTG	ATC	GCG	CGG	GTG	CGC	CGG
	Val	Gly	Met	Val	Gly	Asn	Ile	Leu	Leu	Val	Leu	Val	Ile	Ala	Arg	Val	Arg	Arg
	63			72			81			90			99			108		
	CTG	TAC	AAC	GTG	ACG	AAT	TTC	CTC	ATC	GGC	AAC	CTG	GCC	TTG	TCC	GAC	GTG	CTC
	Leu	Tyr	Asn	Val	Thr	Asn	Phe	Leu	Ile	Gly	Asn	Leu	Ala	Leu	Ser	Asp	Val	Leu
	117			126			135			144			153			162		
	ATG	TGC	ACC	GCC	TGC	GTG	CCG	CTC	ACG	CTG	GCC	TAT	GCC	TTC	GAG	CCA	CGC	GGC
	Met	Cys	Thr	Ala	Cys	Val	Pro	Leu	Thr	Leu	Ala	Tyr	Ala	Phe	Glu	Pro	Arg	Gly
	171			180			189			198			207			216		
	TGG	GTG	TTC	GGC	GGC	GGC	CTG	TGC	CAC	CTG	GTC	TTC	TTC	CTG	CAG	GCG	GTC	ACC
	Trp	Val	Phe	Gly	Gly	Gly	Leu	Cys	His	Leu	Val	Phe	Phe	Leu	Gln	Ala	Val	Thr
	225			234			243			252			261			270		
	GTC	TAT	GTG	TGG	GTG	TTC	ACG	CTC	ACC	ACC	ATC	GCA	GTG	GAC	GCG	TAC	GTC	GTG
	Val	Tyr	Val	Ser	Val	Phe	Thr	Leu	Thr	Thr	Ile	Ala	Val	Asp	Arg	Tyr	Val	Val
	279			288			297			306			315			324		
	CTG	GTG	CAC	CCG	CTG	AGG	CGG	CGC	ATC	TGG	CTG	CGC	CTC	AGC	GCC	TAC	GCT	GTG
	Leu	Val	His	Pro	Leu	Arg	Arg	Arg	Ile	Ser	Leu	Arg	Leu	Ser	Ala	Tyr	Ala	Val
	333			342			351			360			369			378		
	CTG	GCC	ATC	TGG	GTG	CTG	TCC	GCG	GTG	CTG	GCG	CTG	CCC	GCC	GCC	GTG	CAC	ACC
	Leu	Ala	Ile	Trp	Val	Leu	Ser	Ala	Val	Leu	Ala	Leu	Pro	Ala	Ala	Val	His	Thr
	387			396			405			414			423			432		
	TAT	CAC	GTG	GAG	CTC	AAG	CCG	CAC	GAC	GTG	CGC	CTC	TGC	GAG	GAG	TTC	TGG	GGC
	Tyr	His	Val	Glu	Leu	Lys	Pro	His	Asp	Val	Arg	Leu	Cys	Glu	Glu	Phe	Trp	Gly
	441			450			459			468			477			486		
	TCC	CAG	GAG	CGC	CAG	CGC	CAG	CTC	TAC	GCC	TGG	GGG	CTG	CTG	CTG	GTC	ACC	TAC
	Ser	Gln	Glu	Arg	Gln	Arg	Gln	Leu	Tyr	Ala	Trp	Gly	Leu	Leu	Leu	Val	Thr	Tyr
	495			504			513			522			531			540		
	CTG	CTC	CCT	CTG	CTG	GTC	ATC	CTC	CTG	TCT	TAC	GCC	CGG	GTG	TCA	GTG	AAG	CTC
	Leu	Leu	Pro	Leu	Leu	Val	Ile	Leu	Leu	Ser	Tyr	Ala	Arg	Val	Ser	Val	Lys	Leu
	549			558			567			576			585			594		
	CGC	AAC	CGC	GTG	GTG	CCG	GGC	CGC	GTG	ACC	CAG	AGC	CAG	GCC	GAC	TGG	GAC	CGC
	Arg	Asn	Arg	Val	Val	Pro	Gly	Arg	Val	Thr	Gln	Ser	Gln	Ala	Asp	Trp	Asp	Arg
	603			612			621			630			639			648		
	GCT	CGG	CGC	CGG	CGC	ACC	TTC	TGC	TTC	CTG	GTG	GTG	GTC	GTG	GTG	GTG	TTC	ACC
	Ala	Arg	Arg	Arg	Arg	Thr	Phe	Cys	Leu	Leu	Val	Val	Val	Val	Val	Val	Phe	Thr
	657			666														
	CTC	TGC	TGG	CTG	CCC	TTC	TTC	3'										
	Leu	Cys	Trp	Leu	Pro	Phe	Phe											

28 / 79

FIGURE 28



5'	10	19	28	37	46	55
GTG GGC ATG CTG GGC AAC GCC CTG GTC TGT CAT GTC ATC TTC AAG AAC CAG CGA						
Val Cys His Val Ile Phe Lys Asn Gln Arg						
64	73	82	91	100	109	
ATG CAC TCG GCC ACC AGC CTC TTC ATC GTC AAC CTG GCA GTT GCC GAC ATA ATG						
Met His Ser Ala Thr Ser Leu Phe Ile Val Asn Leu Ala Val Ala Asp Ile Met						
118	127	136	145	154	163	
ATC ACG CTG CTC AAC ACC CCC TTC ACT TTG GTT CGC TTT GTG AAC AGC ACA TGG						
Ile Thr Leu Leu Asn Thr Pro Phe Thr Leu Val Arg Phe Val Asn Ser Thr Trp						
172	181	190	199	208	217	
ATA TTT GGG AAG GGC ATG TGC CAT GTC AGC CGC TTT GCC CAG TGC TCA CTG						
Ile Phe Gly Lys Gly Met Cys His Val Ser Arg Phe Ala Gln Tyr Cys Ser Leu						
226	235					
CAC GTC TCA GCA CTG ACA 3'						
His Val Ser Ala Leu Thr						


3 0 / 7 9

FIGURE 30

5'	9	18	27	36	45	54
GAG CCA GCT GAC CTC TTC TGG AAG AAC CTG GAC TTG CCC ACC TTC ATC CTC						
Glu Pro Ala Asp Leu Phe Trp Lys Asn Leu Asp Leu Pro Thr Phe Ile Leu Leu						
63	72	81	90	99	108	
AAC ATC CTG CCC CTC CTC ATC ATC TCT GTG GCC TAC GTT CGT GTG ACC AAG AAA						
Asn Ile Leu Pro Leu Leu Ile Ile Ser Val Ala Tyr Val Arg Val Thr Lys Lys						
117	126	135	144	153	162	
CTG TGG CTG TGT AAT ATG ATT GTC GAT GTG ACC ACA GAG CAG TAC TTT GCC CTG						
Leu Trp Leu Cys Asn Met Ile Val Asp Val Thr Thr Glu Gln Tyr Phe Ala Leu						
171	180	189	198	207	216	
CGG CCC AAA AAG AAG ACC ATC AAG ATG ATG TTG ATG CTG GTG GTA GTC CTC TTT						
Arg Pro Lys Lys Lys Thr Ile Lys Met Leu Met Leu Val Val Val Leu Phe						
225	234					
GCC CTC TGC TGG TTG CCT CTC GAC 3'						
Ala Leu Cys Trp Leu Pro Leu Asp						


3 1 / 7 9

FIGURE 31



3 2 / 7 9

FIGURE 32




	10	20	30	40	50
p63A2	1	MHSATSLFLV	NLKVADIMLV	LINTPFTLVR	FUNSTNIPGR
P30731	1	MHSATSLFLV	NLKVADIMLV	LINTPFTLVR	FVNSTNIPGR
p63A2	51	GCCHVSRPAG	YCSLHVSRAL		
P30731	51	GCCHVSRPAG	YCSLHVSRAL		
p63A2	101				
P30731	101	VIVVMATFFS	LPHALCQKLF	TFKYSEDIVR	SLCLPDPFPH
p63A2	150				
P30731	150				
p63A2	151				
P30731	151				
p63A2	201				
P30731	201				

3 4 / 7 9

1	CATCGTCAAGCAGATGAAGATCATCAAGAGGATGGCTACTCCGAGGGCCAGCAGAAATT	60
1		1
61	CTGCCCTTCTTCCCGGAGTGCTTTCCCGCTCTCCAAACCCCACTCCAGGTGGCCATG	120
1	Met	1
121	GCCTCATGGACCACTGGGGCCCCAGGGTTTCTGACTTATTTCTGGGCTGCGCGGGCG	180
1	AlaSerSerThrThrArgGlyProArgValSerAspLeuPheSerGlyLeuProProAla	21
181	GTCACAACTCCCGCCAAACCAGAGCGCAGAGGCTGGGGGCCAACGGGTGGGTGGCTGGC	240
21	ValThrThrProAlaAsnGlnSerAlaGluAlaSerAlaGlyAsnGlySerValAlaGly	41
241	GCGGACGCTCCAGCGTCAAGGCTTCCAGAGCTGCAGCTGGTGCATCAGCTGAAGGGG	300
41	AlaAspAlaProAlaValThrProPheGlnSerLeuGlnLeuValHisGlnLeuLysGly	61
301	CTGATGTGCTGCTCTACAGCGTGGTGGTGGTGGGGCTGGTGGGCAACTGCTGCTG	360
61	LeuIleValLeuLeuTyrSerValValValValValGlyLeuValGlyAsnCysLeuLeu	81
361	GTGCTGGTGATCGCGGGGTGGCGGGCTGCACAACTGACGAACCTTCTCATCGGCAAC	420
81	ValLeuValIleAlaArgValArgArgLeuHisAsnValThrAsnPheLeuIleGlyAsn	101
421	CTGGCTGTGTCGAAGTGTCTATGTGCACCGCTGGTGGCGCTCAGCTGGCTATGCC	480
101	LeuAlaLeuSerAspValLeuMetCysThrAlaCysValProLeuThrLeuAlaTyrAla	121
481	TTGAGCCACGGGCTGGGTGTTCGGCGGGCGGCTGTGCCACCTGGTCTTCTTCTGCGAG	540
121	PheGluProArgGlyTrpValPheGlyGlyGlyLeuCysHisLeuValPhePheLeuGln	141
541	CGGTACCGCTCTATGTGTGGGTGTTCACGCTCACCACCATGCGAGTGGACCGCTACGTC	600
141	ProValThrValTyrValSerValPheThrLeuThrThrIleAlaValAspArgTyrVal	161
601	GTGCTGGTGCACCGCTGAGGCGGGCATCTGGCTGGCGCTCAGCGCTACGCTGTGCTG	660
161	ValLeuValHisProLeuArgArgArgIleSerLeuArgLeuSerAlaTyrAlaValLeu	181
661	GCCATCTGGGCGCTGTCCGGGTGCTGGCGCTGGCGCGCGCTGCACACCTATCACGTG	720
181	AlaIleTrpAlaLeuSerAlaValLeuAlaLeuProAlaAlaValHisThrTyrHisVal	201
721	GAGCTCAAGCCGACGACGTGGCGCTCTGGAGGAGTTCTGGGGCTCCAGGAGCGCCAG	780
201	GluLeuLysProHisAspValArgLeuCysGluGluPheTrpGlySerGlnGluArgGln	221
781	CGCCAGCTCTACCGCTGGGGCTGCTGCTGGTCACTTACCTGCTCCCTCTGCTGGTCATC	840
221	ArgGlnLeuTyrAlaTrpGlyLeuLeuLeuValThrTyrLeuLeuProLeuLeuValIle	241
841	CTCTGTCTTAAGTCCGGGTGTCACTGAAGCTCCGCAACCGGTGGTGGCGGCTGGGTG	900
241	LeuLeuSerTyrValArgValSerValIleLeuArgAsnArgValValProGlyCysVal	261
901	ACCCAGAGCCAGGCGACTGGGACCGGCTGGGGCGGGCGCACCTTCTGCTTGGTGGTG	960
261	ThrGlnSerGlnAlaAspTrpAspArgAlaArgArgArgThrPheCysLeuLeuVal	281
961	GTGGTGGTGGTGGTGTTCGGCGTCTGCTGGCTGGCGCTGCAGTCTTCAACCTGCTGGG	1020
281	ValValValValValPheAlaValCysTrpLeuProLeuHisValPheAsnLeuLeuArg	301
1021	GACCTGACCCCCACGCCATCGACCTTACGGCTTGGGCTGGTGCAGCTGCTCTGCCAC	1080
301	AspLeuAspProHisAlaIleAspProTyrAlaPheGlyLeuValGlnLeuLeuCysHis	321
1081	TGGCTGGCCATGAGTTGGGCTGCTACAAACCCCTTCATCTACCGCTGGCTGCACGACAGC	1140
321	TrpLeuAlaMetSerSerAlaCysTyrAsnProPheIleTyrAlaTrpLeuHisAspSer	341
1141	TTCCGGAGGAGCTGGCAAACCTGTGGTGGCTGGGGCGGGCAAGATAGCCCCCATGGC	1200
341	PheArgGluGluLeuArgLysLeuLeuValAlaTrpProArgLysIleAlaProHisGly	361
1201	CAGAAATGACCGTCAGGTGGTCACTGTGATGCCACTTAGCCAGGCCTTGGTCAAGGAGC	1260
361	GlnAsnMetThrValSerValValIle***	371
1261	TCCACTTCAACTGGCTCTTAGGGCACCACTCGAGGTCAATCTGGTGGCTATTCTCAGCA	1320
371		371
1321	CCAGAGCTAGC	1331
371		371


3 5 / 7 9

FIGURE 35



3 6 / 7 9

FIGURE 36




3 7 / 7 9

FIGURE 37

5'	9			18			27			36			45			54			
	GTC	GGC	CTG	GTG	GGC	AAC	ATC	CTG	GCT	TCC	TGG	CAC	AAG	CGT	GGA	GGT	CGC	CGT	
	Val	Gly	Leu	Val	Gly	Asn	Ile	Leu	Ala	Ser	Trp	His	Lys	Arg	Gly	Gly	Arg	Arg	
<hr/>																			
	63			72			81			90			99			108			
	GCT	GCT	TGG	GTA	GTG	TGT	GGA	GTC	GTG	TGG	CTG	GCT	GTG	ACA	GCC	CAG	TGC	CTC	
	Ala	Ala	Trp	Val	Val	Cys	Gly	Val	Val	Trp	Leu	Ala	Val	Thr	Ala	Gln	Cys	Leu	
<hr/>																			
	117			126			135			144			153			162			
	CCC	ACG	GCA	GTC	TTT	GCT	GCC	ACA	GGC	ATC	CAG	CGC	AAC	CGC	ACT	GTG	TGC	TAC	
	Pro	Thr	Ala	Val	Phe	Ala	Ala	Thr	Gly	Ile	Gln	Arg	Asn	Arg	Thr	Val	Cys	Tyr	
<hr/>																			
	171			180			189			198			207			216			
	GAC	CTG	AGC	CCA	CCC	ATC	CTG	TCT	ACT	CGC	TAC	CTG	CCC	TAT	GGT	ATG	GCC	CTC	
	Asp	Leu	Ser	Pro	Pro	Ile	Leu	Ser	Thr	Arg	Tyr	Leu	Pro	Tyr	Gly	Met	Ala	Leu	
<hr/>																			
	225			234			243			252			261			270			
	ACG	GTC	ATC	GGC	TTC	TTG	CTG	CCC	TTC	ATA	GCC	TTA	CTG	GCT	TGT	TAT	TGT	CGC	
	Thr	Val	Ile	Gly	Phe	Leu	Leu	Pro	Phe	Ile	Ala	Leu	Leu	Ala	Cys	Tyr	Cys	Arg	
<hr/>																			
	279			288			297			306			315			324			
	ATG	GCC	CGC	CGC	CTG	TGT	CGC	CAG	GAT	GGC	CCA	GCA	GGT	CCT	GTG	GCC	CAA	GAG	
	Met	Ala	Arg	Arg	Leu	Cys	Arg	Gln	Asp	Gly	Pro	Ala	Gly	Pro	Val	Ala	Gln	Glu	
<hr/>																			
	333			342			351			360			369			378			
	CGG	CGC	AGC	AAG	GCG	GCT	CGT	ATG	GCT	GTG	GTG	GTG	GCA	GCT	GTC	TTT	GCC	CTC	
	Arg	Arg	Ser	Lys	Ala	Ala	Arg	Met	Ala	Val	Val	Val	Ala	Ala	Val	<u>Phe</u>	<u>Ala</u>	<u>Leu</u>	
<hr/>																			
	387			396															
	TGC	TGG	CTG	OCT	CTC	TAC	3'												
	Cys	Trp	Leu	Pro	Leu	Tyr													

3 8 / 7 9

FIGURE 38



3 9 / 7 9

FIGURE 39

	10	20	30	40	50
P3H2-17	1 VGLVGNILAS	WHKRGRRRAA	WVVCGVVWLA	VTACCLPTAV	FAATG QRN-
P34996	1 RYTGVVHPLK	SLGRLLKKNA	VYVSSLVWAL	VVAVIAPILF	YSGIGVRRN-
A46226	1 RYLAVVHPTIR	SARWETAPVA	RTVSAADVVA	SAVVVLPPVV	F--SGVPRG-
JN0605	1 RYVAVVHPLP	AATYERPSVA	KLINLGVWLA	SLVLTPIAI	FADIRPARGG
S28787	1 RYLAI VHATN	SQKPRKLLAE	KVVYVGVWLP	AVLLTIPDLI	FADIKEVDE-
	60	70	80	90	100
P3H2-17	51 RIV-CYDL--	SEPI STRYL	PYGMALIVIG	FLLPFIALLA	CYCRMARRLC
P34996	51 KITTGYDT--	TAD EYLBSYF	VYSMCTIVFM	ECIPEFIVILG	CYGLIVKALI
A46226	51 MST-CHMOWP	EPAAAWPAGF	I IY--IAALG	FFGPLLVIICL	CYLLIVVKVR
JN0605	51 QAVAGNLQWP	HPAWSAVFVV	YTF----LGG	FLPVLAIIGL	CYLLIVGKVR
S28787	51 RYI-CDRF--	YFSDLWLVVE	QFQ--HIVVG	LLLPGLVILS	CYGLIISKLS
	110	120	130	140	150
P3H2-17	101 RQDGP A-GPV	AOE-RRS--K	AARMAVVVAA	VFALCWLP LY
P34996	101 YKDLDN-SPL	---RR---K	SIYLVII VLT	VFAVSYP PH
A46226	101 SAGR VWAP S	CORRRRSERR	VTRVVA VVA	LEVL CWMPFY
JN0605	101 AVALRA---G	WQRRRSEKK	I TRLVLMVVV	VFVL CWMPFY
S28787	101 HSKG-----	YQKR-----K	ALKT TVILIL	TEFACWLP PY


40 / 79

	10			19			28			37			46			55		
5'	GTG	GGC	CTG	GTG	GGC	AAC	TTC	CTG	GCC	GCG	ATG	TCT	GTG	GAT	CGC	TAC	GTG	GCC
	Val	Gly	Leu	Val	Gly	Asn	Phe	Leu	Ala	Ala	Met	Ser	Val	Asp	Arg	Tyr	Val	Ala
	64			73			82			91			100			109		
	ATT	GTG	CAC	TGG	CGG	CGC	TCC	TCC	TCC	CTC	AGG	GTG	TCC	CGC	AAC	GCA	CTG	CTG
	Ile	Val	His	Ser	Arg	Arg	Ser	Ser	Ser	Leu	Arg	Val	Ser	Arg	Asn	Ala	Leu	Leu
	118			127			136			145			154			163		
	GGC	GTG	GGC	TTC	ATC	TGG	GCG	CTG	TCC	ATC	GCC	ATG	GCC	TGG	CCG	GTG	GCC	TAC
	Gly	Val	Gly	Phe	Ile	Trp	Ala	Leu	Ser	Ile	Ala	Met	Ala	Ser	Pro	Val	Ala	Tyr
	172			181			190			199			208			217		
	CAC	CAG	CGT	CTT	TTC	CAT	CGG	GAC	AGC	AAC	CAG	ACC	TTC	TGC	TGG	GAG	CAG	TGG
	His	Gln	Arg	Leu	Phe	His	Arg	Asp	Ser	Asn	Gln	Thr	Phe	Cys	Trp	Glu	Gln	Trp
	226			235			244			253			262			271		
	CCC	AAC	AAG	CTC	CAC	AAG	AAG	GCT	TAC	GTG	GTG	TGC	ACT	TTC	GTC	TTT	GGG	TAC
	Pro	Asn	Lys	Leu	His	Lys	Lys	Ala	Tyr	Val	Val	Cys	Thr	Phe	Val	Phe	Gly	Tyr
	280			289			298			307			316			325		
	CTT	CTG	CCC	TTA	CTG	CTC	ATC	TGC	TTT	TGC	TAT	GCC	AAG	GTC	CTT	AAT	CAT	CTG
	Leu	Leu	Pro	Leu	Leu	Leu	Ile	Cys	Phe	Cys	Tyr	Ala	Lys	Val	Leu	Asn	His	Leu
	334			343			352			361			370			379		
	CAT	AAA	AAG	CTG	AAA	AAC	ATG	TCA	AAA	AAG	TCT	GAA	GCA	TCC	AAG	AAA	AAG	ACT
	His	Lys	Lys	Leu	Lys	Asn	Met	Ser	Lys	Lys	Ser	Glu	Ala	Ser	Lys	Lys	Lys	Thr
	388			397			406			415			424			433		
	GCA	CAG	ACC	GTC	CTG	GTG	GTC	GTT	GTA	GTA	TTT	GCC	CTC	TGC	TGG	CTG	CCT	TTC
	Ala	Gln	Thr	Val	Leu	Val	Val	Val	Val	Val	Phe	Ala	Leu	Cys	Trp	Leu	Pro	Phe
	TAC 3'																	

	Tyr																	

4 1 / 7 9

FIGURE 41



4 2 / 7 9

FIGURE 42

p3H2-34	1	VGLVGNFLAA	MSYDRYVAVV	HSRFSSSLFV	SRNALGVGF	IAALSLAMAS	50
JN0605	1	HTTSVECLTV	LSVDRYVAVV	HELRAITVRR	ESVAKLINIG	IAALSLAMAS	50
B41795	1	QETSIFCLTV	MSIDRYLAVV	HPIKSEKWRP	PRDAKMIIMA	IAALSLAMAS	50
A39297	1	HTTSIVLTV	LSVDRYVAVV	HPIKSEKWRP	PRDAKMIIMA	IAALSLAMAS	50
p3H2-34	51	MA-YHORLF	HSISNQTF	EQHNKHK-	-KAYVVCITV	FYLLPLLI	100
JN0605	51	PIATIEDIRP	ALGQOAVA	LONHPEWS-	-AVFVWYTEL	LCHELLFVLA	100
B41795	51	PIMIVGLAS	QWGRSS-CT	INWIGESGA	YTGELIKTEL	LCHELLFVLA	100
A39297	51	PIWESRIPAA	SEGIVA-CT	MLMEERAR	LVGEVLKTEL	MSFILLINGAI	100
p3H2-34	101	FLY----	VLNHLHGKLG	NMSKKSEASK	KKTAQTVLV	AVVEALCHLF	150
JN0605	101	GLCYILLIAG	HAVALRAG	QOPRSE---	KKITRLVLA	AVVEVLCIML	150
B41795	101	GLCYIFILLI	VKSSGIFVCS	SKRKSE---	KKVTRVSI	AVVEVLCIML	150
A39297	101	FLCYVALLIAK	HYVALKAG	QOKRSE---	RKITLAVAV	AVVEVLCIML	150
p3H2-34	151	FY.....	200
JN0605	151	FY.....	200
B41795	151	FY.....	200
A39297	151	FY.....	200


4 3 / 7 9

FIGURE 43

5'	GTG	GCC	ATG	GTG	GCC	AAC	GTC	CTG	GTG	CTC	TGG	TTC	TTC	GCC	TTC	TCC	ATC	AAG	55
	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
	Val	Gly	Met	Val	Gly	Asn	Val	Leu	Val	Leu	Trp	Phe	Phe	Gly	Phe	Ser	Ile	Lys	
	64	ACC	CCC	TTC	TCC	GTC	TAC	TTC	CTG	CAC	CTG	GCC	AGC	GCC	GAC	GCC	GCC	TAC	109
	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
	Arg	Thr	Pro	Phe	Ser	Val	Tyr	Phe	Leu	His	Leu	Ala	Ser	Ala	Asp	Gly	Ala	Tyr	
	118	TTC	AGC	AAG	GCC	GTC	TTC	TCC	CTG	CTG	AAC	GCC	GCC	GCC	TTC	CTG	GCC	ACC	163
	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
	Leu	Phe	Ser	Lys	Ala	Val	Phe	Ser	Leu	Leu	Asn	Ala	Gly	Gly	Phe	Leu	Gly	Thr	
	172	GCC	CAC	TAT	GTG	CGC	AGC	GTG	GCC	CGG	GTG	CTG	GCG	CTC	TGC	GCC	TTC	GTG	217
	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
	Phe	Ala	His	Tyr	Val	Arg	Ser	Val	Ala	Arg	Val	Leu	Gly	Leu	Cys	Ala	Phe	Val	
	226	GCC	GTG	AGC	CTC	CTG	CCG	GCC	GTG	AGC	ATG	GAG	CGC	TGC	GCG	TCT	G	3'	
	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
	Ala	Gly	Val	Ser	Leu	Leu	Pro	Ala	Val	Ser	Met	Glu	Arg	Cys	Ala	Ser			
						235			244			253			262				

4 4 / 7 9

FIGURE 44




4 6 / 7 9

FIGURE 46

1	CAAAGCAACAGGTGCAACCTCAAGGCACTGAAAGCAAGGGGACGCAGCTCACAAGGGCCAAGGGATTGAACC	72
1		1
73	CATAACCGCTCAGAAGATTCTCCGCCTGCGGAGAGCTCCGGAGGAGTCCCACCCGTCCAGCTTGCTGACTGC	144
1		1
145	GAGCAGTGAGAGTCCGCTAGACCCGTACCTCTGTGTCTCTGGAGCCTGCCGCCCCCGCACGGGAAAGGCTTAG	216
1		1
217	CTCGGGACTTGCAGCACCGCCTCCTCTTTAGCCAGGCCAGGCACGAGGATAGTGTGATCGGGCACAGCCAGG	288
1		1
289	GTCGCTCTTCCAGGCTTTCTTGCGGGTTGCGGGAGGTACTAGTTGGAGACGCGCGGCTCGCTCTCGCCGCT	360
1		1
361	CTGTCTTGGGCCACTCCGTGATCCTAGGCTACCTCCAGAGCCAGTTTTCCTGGCTGGCACAACCTCTCCAGG	432
1		1
433	GCGCTCCGGTCCGTTGCACAGCGCCCCAAGGGGGTATCCCAAGTAAGTGATGGAACCTGGCTATGGTGAACCTC	504
1	MetGluLeuAlaMetValAsnLeu	8
505	AGTGAAGGGAATGGGAGCGACCCAGAGCCGCCAGCCCCGAGTCCAGGCCGCTCTTCGGCATTGGCGTGGAG	576
8	SerGluGlyAsnGlySerAspProGluProProAlaProGluSerArgProLeuPheGlyIleGlyValGlu	32
577	AACTTCATTACGCTGGTAGTGTTCGGCTGATTTTCGCGATGGGCGTGCTGGGCAACAGCCTGGTGATCACC	648
32	AsnPheIleThrLeuValValPheGlyLeuIlePheAlaMetGlyValLeuGlyAsnSerLeuValIleThr	56
649	GTGCTGGCGCGCAGCAAACCAGGCAAGCCGCGCAGCACCACCAACCTGTTTATCTCAATCTGAGCATCGCA	720
56	ValLeuAlaArgSerLysProGlyLysProArgSerThrThrAsnLeuPheIleLeuAsnLeuSerIleAla	80
721	GACCTGGCCTACCTGCTCTTCTGCATCCCTTTTCAGGCCACCGTGATGCACTGCCCACCTGGGTGCTGGGC	792
80	AspLeuAlaTyrLeuLeuPheCysIleProPheGlnAlaThrValTyrAlaLeuProThrTrpValLeuGly	104
793	GCCTTCATCTGCAAGTTTATACACTACTTCTTCACCGTGCCATGCTGGTGAGCATCTTCACCTGGCCGCG	864
104	AlaPheIleCysLysPheIleHisTyrPhePheThrValSerMetLeuValSerIlePheThrLeuAlaAla	128
865	ATGCTGTGGATCGCTACGTGGCCATTGTGCACTCGGGCGCTCCTCCTCCCTCAGGGTGTCCCGCAACGCA	936
128	MetSerValAspArgTyrValAlaIleValHisSerArgArgSerSerSerLeuArgValSerArgAsnAla	152
937	CTGCTGGGCGTGGGCTTCATCTGGGCGCTGTCCATCGCCATGGCCTCGCCGTTGGCCTACCACAGCGTCTT	1008
152	LeuLeuGlyValGlyPheIleTrpAlaLeuSerIleAlaMetAlaSerProValAlaTyrHisGlnArgLeu	176
1009	TTCCATCGGGACAGCAACCAGACCTTCTGCTGGGAGCAGTGGCCCAACAAGCTCCACAAGAAGGCTTACGTG	1080
176	PheHisArgAspSerAsnGlnThrPheCysTrpGluGlnTrpProAsnLysLeuHisLysLysAlaTyrVal	200
1081	GTGTGCACTTTCGTCTTTGGGTACCTTCTGCCCTTACTGCTCATCTGCTTTTGCTATGCCAAGGTCCTTAAT	1152
200	ValCysThrPheValPheGlyTyrLeuLeuProLeuLeuLeuIleCysPheCysTyrAlaLysValLeuAsn	224
1153	CATCTGCATAAAAAGCTGAAAAACATGTCAAAAAGTCTGAAGCATCCAAGAAAAGACTGCACAGACCGTC	1224
224	HisLeuHisLysLysLeuLysAsnMetSerLysLysSerGluAlaSerLysLysLysThrAlaGlnThrVal	248
1225	CTGGTGGTCTGTGTAGTATTGGCATATCCTGGCTGCCCATCATGTCTGTCACCTCTGGGCTGAGTTTGGGA	1296
248	LeuValValValValValPheGlyIleSerTrpLeuProHisHisValValHisLeuTrpAlaGluPheGly	272
1297	GCCTTCCCACTGACGCCAGCTTCCTTCTTCTTCAGAATCACCGCCCATTGCTGGCATACAGCAACTCCTCA	1368
272	AlaPheProLeuThrProAlaSerPhePhePheArgIleThrAlaHisCysLeuAlaTyrSerAsnSerSer	296
1369	GTGAACCCCATCATATATGCGTTTCTCTCAGAAAACCTCCGGAAGGGGTACAAGCAAGTGTTCAGTGTAT	1440
296	ValAsnProIleIleTyrAlaPheLeuSerGluAsnPheArgLysAlaTyrLysGlnValPheLysCysHis	320
1441	GTTTGGATGAATCTCCACGCAGTGAACCTAAGGAAAACAAGAGCCGGATGGACACCCCGCCATCCACCAAC	1512
320	ValCysAspGluSerProArgSerGluThrLysGluAsnLysSerArgMetAspThrProProSerThrAsn	344
1513	TGCACCCACGTGTGAAGGTTTTCGGGAGCCTCCCGACTTCCAGCTCCCATGTGTGTTAGAGAGAGGAGGGCG	1584
344	CysThrHisVal***	349
1585	GAGCGAATTATCAAGTAACATGG	1607
349		349

47 / 79

FIGURE 47



4 9 / 7 9


FIGURE 49

5'	CTC	GCG	GCT	CTG	GGT	ATG	GAT	CGG	TAT	CTT	CTC	ACC	CTT	CAC	CCA	GTG	TGG	TCC
										Leu	Leu	Thr	Leu	His	Pro	Val	Trp	Ser
	CAA	AAG	CAC	CGA	ACC	TCA	CAC	TGG	GCT	TCC	AGA	GTC	GTT	CTG	GGA	GTC	TGG	CTC
	Gln	Lys	His	Arg	Thr	Ser	His	Trp	Ala	Ser	Arg	Val	Val	Leu	Gly	Val	Trp	Leu
	TCT	GCC	ACT	GCC	TTC	AGC	GTG	CCC	TAT	TTC	GTT	TTC	AGG	GAG	ACA	TAT	GAT	GAC
	Ser	Ala	Thr	Ala	Phe	Ser	Val	Pro	Tyr	Leu	Val	Phe	Arg	Glu	Thr	Tyr	Asp	Asp
	CGT	AAA	GGA	AGA	GTG	ACC	TGC	AGA	AAT	AAC	TAC	GCT	GTG	TCC	ACT	GAC	TGG	GAA
	Arg	Lys	Gly	Arg	Val	Thr	Cys	Arg	Asn	Asn	Tyr	Ala	Val	Ser	Thr	Asp	Trp	Glu
	AGC	AAA	GAG	ATG	CAA	ACA	GTA	AGA	CAA	TGG	ATT	CAT	GCC	ACC	TGT	TTC	ATC	AGC
	Ser	Lys	Glu	Met	Gln	Thr	Val	Arg	Gln	Trp	Ile	His	Ala	Thr	Cys	Phe	Ile	Ser
	CGC	TTC	ATA	CTG	GGC	TTC	CTT	CTG	CCT	TTC	TTA	GTC	ATT	GGC	TTT	TGT	TAT	GAA
	Arg	Phe	Ile	Leu	Gly	Phe	Leu	Leu	Pro	Phe	Leu	Val	Ile	Gly	Phe	Cys	Tyr	Glu
	AGA	GTA	GCC	CGC	AAG	ATG	AAA	GAG	AGG	GGC	CTC	TTT	AAA	TCC	AGC	AAA	CCC	TTC
	Arg	Val	Ala	Arg	Lys	Met	Lys	Glu	Arg	Gly	Leu	Phe	Lys	Ser	Ser	Lys	Pro	Phe
	AAA	GTC	ACG	ATG	ACT	GCT	GTT	ATC	TCT	TTT	TTC	TGT	CCT	GGC	TTC	CCT	ACC	ACA
	Lys	Val	Thr	Met	Thr	Ala	Val	Ile										

TG 3'

5 0 / 7 9

FIGURE 50



5 1 / 7 9

FIGURE 51


PMJ10	1	10	20	30	40	50	
B42009	1	1	1	1	1	1	50
JC2014	1	1	1	1	1	1	50
A46520	1	1	1	1	1	1	50
A46525	1	1	1	1	1	1	50
S28787	1	1	1	1	1	1	50
PMJ10	51	60	70	80	90	100	
B42009	51	51	51	51	51	51	100
JC2014	51	51	51	51	51	51	100
A46520	51	51	51	51	51	51	100
A46525	51	51	51	51	51	51	100
S28787	51	51	51	51	51	51	100
PMJ10	101	110	120	130	140	150	
B42009	101	101	101	101	101	101	150
JC2014	101	101	101	101	101	101	150
A46520	101	101	101	101	101	101	150
A46525	101	101	101	101	101	101	150
S28787	101	101	101	101	101	101	150

FIGURE 52

	9			18			27			36			45			54		
5'	CTG	ACT	GCT	CTG	GGG	ACT	GAC	CGG	TAT	TTC	AAG	ATT	GTG	AAG	CCC	CTT	TCC	ACG
	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
										Phe	Lys	Ile	Val	Lys	Pro	Leu	Ser	Thr
	63			72			81			90			99			108		
	TCC	TTC	ATC	CAG	TCT	GTG	AAC	TAC	AGC	AAA	CTC	GTC	TCG	CTG	GTG	GTC	TGG	TTG
	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
	Ser	Phe	Ile	Gln	Ser	Val	Asn	Tyr	Ser	Lys	Leu	Val	Ser	Leu	Val	Val	Trp	Leu
	117			126			135			144			153			162		
	CTC	ATG	CTC	CTC	CTC	GCC	GTC	CCC	AAC	GTC	ATT	CTC	ACC	AAC	CAG	AGA	GTT	AAG
	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
	Leu	Met	Leu	Leu	Leu	Ala	Val	Pro	Asn	Val	Ile	Leu	Thr	Asn	Gln	Arg	Val	Lys
	171			180			189			198			207			216		
	GAC	GTG	ACG	CAG	ATA	AAA	TGC	ATG	GAA	CTT	AAA	AAC	GAA	CTG	GGC	CGC	CAG	TGG
	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
	Asp	Val	Thr	Gln	Ile	Lys	Cys	Met	Glu	Leu	Lys	Asn	Glu	Leu	Gly	Arg	Gln	Trp
	225			234			243			252			261			270		
	CAC	AAG	GCG	TCA	AAC	TAC	ATC	TTT	GTG	GGC	ATT	TTC	TGG	CTT	GTG	TTC	CTT	TTG
	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
	His	Lys	Ala	Ser	Asn	Tyr	Ile	Phe	Val	Gly	Ile	Phe	Trp	Leu	Val	Phe	Leu	Leu
	279			288			297			306			315			324		
	CTA	ATC	ATT	TTC	TAC	ACT	GCT	ATC	ACC	AGG	AAA	ATC	TTT	AAG	TCC	CAC	CTG	AAA
	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
	Leu	Ile	Ile	Phe	Tyr	Thr	Ala	Ile	Thr	Arg	Lys	Ile	Phe	Lys	Ser	His	Leu	Lys
	333			342			351			360			369			378		
	TCC	AGA	AAG	AAT	TCC	ATC	TGG	GTC	AAA	AAG	AAA	TCT	AGC	CGC	AAC	ATC	TTC	AGC
	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
	Ser	Arg	Lys	Asn	Ser	Ile	Ser	Val	Lys	Lys	Lys	Ser	Ser	Arg	Asn	Ile	Phe	Ser
	387			396			405			414								
	ATC	GTG	TTT	ATC	CTC	TGT	TGG	CCC	CCC	TAC	CAC	ATC	3'					
	---	---	---	---	---	---	---	---	---	---	---	---						
	Ile	Val																

53 / 79

FIGURE 53



5 4 / 7 9

FIGURE 54

PMH28	1	10	20	30	40	50
P35343	1	FKIVKPLSTS	FIOSVNYSKL	VSIVVNLML	LIAVPNVILT	NQVKDVTQI
A41795	1	LALVHATST-	LQKRHLVKE	VCIAMNLSV	ILALPILIER	NEVKVNLSTL
A47457	1	VAVVHPEIKAA	RYRRFTVAKV	VNLGVVLES	LVTLPINWFS	RITAANSDGIV
	1	VAVVHPELRAA	TYRRPSVAKI	INLGVVLSL	LVTLPINWFS	DIRPARGGC-
PMH28	51	60	70	80	90	100
P35343	51	KCME-LKNEI	GROHKAANY	IFVGIWLV	FLLIIFYTA	IT-RKIFKSH
A41795	51	VCYEDVGNT	SRL--RVLR	ILPQTEGFLV	PLLLMLFCYG	FTLRITLFAH
A47457	51	ACNM-LVPEH	AQRVLGFEV	LYTFMLGFL	PVGAICLCYV	LITAKMRMVA
	51	AVAC-NLQWE	HPA-SAVEV	VYTFELGELL	EVLAIGLCYL	LIVGKBRVA
PMH28	101	110	120	130	140	150
P35343	101	LKSRKNSI-S	VKKKSSRNIF	S--IV....
A41795	101	MG----QKH	AMR----VIF	AWLV....
A47457	101	LKAGNCQKR	SEKKTLMVM	WVWV....
	101	LRAGNCQRR	SEKKTIRLVL	WVWV....

5 5 / 7 9

FIGURE 55

```

      9      18      27      36      45      54
5'  GCC ACC AAC GTG TTC ATC CTG TGT CTG GTG GAC CTG CTG GCT GCC CTG ACC CTC
    -----
                                Val Asp Leu Leu Ala Ala Leu Thr Leu

      63      72      81      90      99      108
ATG CCT CTG GCC ATG CTC TCC AGC TCC GCC CTC TTT GAC CAC GCC CTC TTT GGG
    -----
Met Pro Leu Ala Met Leu Ser Ser Ser Ala Leu Phe Asp His Ala Leu Phe Gly

      117      126      135      144      153      162
GAG GTG GCC TGC CGC CTC TAC TTG TTC CTG AGC GTC TGC TTT GTC AGC CTG GCC
    -----
Glu Val Ala Cys Arg Leu Tyr Leu Phe Leu Ser Val Cys Phe Val Ser Leu Ala

      171      180      189      198      207      216
ATC CTC TCG GTG TCC GCC ATC AAT GTG GAG CGC TAC TAT TAT GTG GTC CAC CCC
    -----
Ile Leu Ser Val Ser Ala Ile Asn Val Glu Arg Tyr Tyr Tyr Val Val His Pro

      225      234      243      252      261      270
ATG CGC TAT GAG GTG CGC ATG AAA CTG GGG CTG GTG GCC TCT GTG CTG GTG GGC
    -----
Met Arg Tyr Glu Val Arg Met Lys Leu Gly Leu Val Ala Ser Val Leu Val Gly

      279      288      297      306      315      324
GTG TGG GTG AAG GCC CTG GCC ATG GCT TCT GTG CCA GTG TTG GGA AGG GTG TCC
    -----
Val Trp Val Val Lys Ala Leu Ala Met Ala Ser Val Pro Val Leu Gly Arg Val Ser

      333      342      351      360      369      378
TGG GAG GAA GGC CCT CCC AGT GTC CCC CCA GGC TGT TCA CTC CAA TGG AGC CAC
    -----
Trp Glu Glu Gly Pro Pro Ser Val Pro Pro Gly Cys Ser Leu Gln Trp Ser His

      387      396      405      414      423      432
AGT GCC TAC TGC CAG CTT TTC GTG GTG GTC TTC GCC GTC CTC TAC TTC CTG CTG
    -----
Ser Ala Tyr Cys Gln Leu Phe Val Val Val Phe Ala Val Leu Tyr Phe Leu Leu

      441      450      459      468      477      486
CCC CTG CTC CTC ATC CTT GTG GTC TAC TGC AGC ATG TTC CGG GTG GCT CGT GTG
    -----
Pro Leu Leu Leu Ile Leu Val Val Tyr Cys Ser Met Phe Arg Val Ala Arg Val

      495      504      513      522      531      540
GCT GCC ATG CAG CAC GGG CCG CTG CCC ACG TGG ATG GAG ACG CCC CGG CAA CGC
    -----
Ala Ala Met Gln His Gly Pro Leu Pro Thr Trp Met Glu Thr Pro Arg Gln Arg

```


5 6 / 7 9

FIGURE 56

549	558	567	576	585	594
TCC GAG TCT CTC AGC AGC CGC TCC ACT ATG GTC ACC AGC TCG GGG GCC CCG CAG					
---	---	---	---	---	---
Ser Glu Ser Leu Ser Ser Arg Ser Thr Met Val Thr Ser Ser Gly Ala Pro Gln					
603	612	621	630	639	648
ACC ACC CCT CAC CGG ACG TTT GGC GGA GGG AAG GCA GCA GTG GTC CTC CTG GCT					
---	---	---	---	---	---
Thr Thr Pro His Arg Thr Phe Gly Gly Gly Lys Ala Ala Val Val Leu Leu Ala					
657	666	675	684	693	702
GTG GGA GGA CAG TTC CTG CTC TGT TGG TTG CCC TAC TTC TCC TTC CAC CTC TAT					
---	---	---	---	---	---
Val Gly Gly Gln Phe Leu Leu Cys Trp Leu Pro Tyr Phe Ser Phe His Leu Tyr					
711	720	729	738	747	756
GTG GCC CTG AGC GCT CAG CCC ATT GCA GCG GGG CAG GTG GAG AAC GTG GTG ACC					
---	---	---	---	---	---
Val Ala Leu Ser Ala Gln Pro Ile Ala Ala Gly Gln Val Glu Asn Val Val Thr					
765	774	783	792	801	810
TGG ATT GGC TAC TTC TGC TTC ACC TCC <u>AAC CCT CTC CTC TAT TCC TTC CTC CCT</u> 3'					
---	---	---	---	---	---
Trp Ile Gly Tyr Phe Cys Phe Thr Ser					


57 / 79

FIGURE 57




58 / 79

FIGURE 58



59 / 79

FIGURE 59



6 0 / 7 9

FIGURE 60

p19P2	10	20	30	40	50
S12863	1 VGMVGNLIV LVIVARVRRLH NVINFLIGNL ALSDVLMCTA CVELILAYAF				50
	1 LGVSGNLAII IIILKQKEMR NVINILIVNL SFSDLLVAVM CLFFIFVYTL				50
p19P2	60	70	80	90	100
S12863	51 EPRGIVFCGG LCHLVFELQP VIVVVSVEITL TTIAVDRIYV LVHFLRRRI-				100
	51 MDH-IVFGET MCKLNPVQC VSITVSIESTL VLIAVERHQL IINIRGWRPN				100
p19P2	110	120	130	140	150
S12863	101 ----- IWVLAVASSL PFVIYQILTD EPFQNVSLAA FKDKYVCFDK				150
p19P2	160	170	180	190	200
S12863	151 -----GLLVV TYLPLLVIL LS-----V VRVSVKLRNE VVPGCVTQSQ				200
	151 FPSDSHRISTY ITLLVLOYF GPLCFIFICY FKIIYIRLKR NNMMDKIRDS				200
p19P2	210	220	230	240	250
S12863	201 ADWDRARRRR TFCLLVWVV VFAICVLEFY				250
	201 KYRSSETKRI NVMLLSIWA -EAVCHLELT				250

6 2 / 7 9

FIGURE 62

```

5'  CTG  TGT  GTC  ATC  GCG  GTG  GAT  AGG  TAC  GTG  GTT  CTG  GTG  CAC  CCG  CTA  CGT  CCG
    ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---
    Leu  Cys  Val  Ile  Ala  Val  Asp  Arg  Tyr  Val  Val  Leu  Val  His  Pro  Leu  Arg  Arg

      9      18      27      36      45      54
    CGC  ATT  TCA  CTG  AGG  CTC  AGC  GCC  TAC  GCG  GTG  CTG  GGC  ATC  TGG  GCT  CTA  TCT
    ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---
    Arg  Ile  Ser  Leu  Arg  Leu  Ser  Ala  Tyr  Ala  Val  Leu  Gly  Ile  Trp  Ala  Leu  Ser

      63      72      81      90      99      108
    GCA  GTG  CTG  GCG  CTG  CCG  GCC  GCG  GTG  CAC  ACC  TAC  CAT  GTG  GAG  CTC  AAG  CCC
    ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---
    Ala  Val  Leu  Ala  Leu  Pro  Ala  Ala  Val  His  Thr  Tyr  His  Val  Glu  Leu  Lys  Pro

      117     126     135     144     153     162
    CAC  GAC  GTG  AGC  CTC  TGC  GAG  GAG  TTC  TGG  GGC  TGG  CAG  GAG  CGC  CAA  CGC  CAG
    ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---
    His  Asp  Val  Ser  Leu  Cys  Glu  Glu  Phe  Trp  Gly  Ser  Gln  Glu  Arg  Gln  Arg  Gln

      171     180     189     198     207     216
    ATC  TAC  GCC  TGG  GGG  CTG  CTT  CTG  GGC  ACC  TAT  TTG  CTC  CCC  CTG  CTG  GCC  ATC
    ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---
    Ile  Tyr  Ala  Trp  Gly  Leu  Leu  Leu  Gly  Thr  Tyr  Leu  Leu  Pro  Leu  Leu  Ala  Ile

      225     234     243     252     261     270
    CTC  CTG  TCT  TAC  GTA  CCG  GTG  TCA  GTG  AAG  CTG  AGG  AAC  CGC  GTG  GTG  CCT  GGC
    ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---
    Leu  Leu  Ser  Tyr  Val  Arg  Val  Ser  Val  Lys  Leu  Arg  Asn  Arg  Val  Val  Pro  Gly

      279     288     297     306     315     324
    AGC  GTG  ACC  CAG  AGT  CAA  GCT  GAC  TGG  GAC  CGA  GCG  CGT  CGC  CGC  CGC  ACT  TTC
    ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---
    Ser  Val  Thr  Gln  Ser  Gln  Ala  Asp  Trp  Asp  Arg  Ala  Arg  Arg  Arg  Arg  Thr  Phe

      323     342     351     360     369     378
    TGT  CTG  CTG  GTG  GTG  GTG  GTG  GTA  GTG  TTC  ACG  CTC  TGC  TGG  CTG  CCC  TTC  TAC
    ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---
    Cys  Leu  Leu  Val  Val  Val  Val  Val  Val  Phe  Thr  Leu  Cys  Trp  Leu  Pro  Phe  Tyr

```

CT 3'


6 3 / 7 9

FIGURE 63

p19P2	1	10	20	30	40	50	
pg3-2/pg1-10	1	VGMVGNVLLV	LVIARVRRLH	NVTNFIIGNL	ALSDVLMCTA	CVPLTLAYAF	50
p5S38	-79	VGMVGNVLLV	LVIRVRRLY	NVTNFIIGNL	ALSDVLMCTA	CVPLTLAYAF	50
							-30
p19P2	51	60	70	80	90	100	
pg3-2/pg1-10	51	EPRGVAFGGG	LCHLVFFLCP	VTVYVSVFTI	TTIAVDRYAV	LVHPLRRRI-	100
p5S38	-29	EPRGVAFGGG	LCHLVFFLCA	VTVYVSVFTI	TTIAVDRYAV	LVHPLRRRIIS	100
					CVIAVDRYAV	LVHPLRRRIIS	21
p19P2	101	110	120	130	140	150	
pg3-2/pg1-10	101	LRLSAYAVIA	IRLSAVIAL	PAVHTYHVE	LKPHDVRLCE	EFWGSQERQF	150
p5S38	22	LRLSAYAVIG	IRLSAVIAL	PAVHTYHVE	LKPHDVSLCE	EFWGSQERQF	150
							71
p19P2	151	160	170	180	190	200	
pg3-2/pg1-10	151	QLYANGLLVA	FYLLPLLVIL	LSYVRVSVKI	RNRVVFPGCVT	QSQADHIDRAF	200
p5S38	72	QLYANGLLIG	FYLLPLLVIL	LSYVRVSVKI	RNRVVFPGCVT	QSQADHIDRAF	200
					RNRVVFPGSVI	QSQADHIDRAF	121
p19P2	201	210	220	230	240	250	
pg3-2/pg1-10	201	RRRTFCCLVA	VVVFATCML	FVY			250
p5S38	122	RRRTFCCLVA	VVVFATCML	PFE			250
				PFY			171


6 4 / 7 9

FIGURE 64



6 5 / 7 9


FIGURE 65



6 6 / 7 9


FIGURE 66

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17




67 / 79

FIGURE 67



6 8 / 7 9

FIGURE 68




6 9 / 7 9

FIGURE 69

1	GAGCATAGGAAAGGCTGACAGGCAGTTATGGAGCAGGACAATGGCACCATCCAGGCTCCA	60
1	MetGluGlnAspAsnGlyThrIleGlnAlaPro	11
61	GGCTTGCCGCCACCACCTGCGTCTACCGTGAGGATTTCAAGCGACTGCTGCTAACCCCG	120
11	GlyLeuProProThrThrCysValTyrArgGluAspPheLysArgLeuLeuLeuThrPro	31
121	GTATACTCGGTGGTGTGGTGGTGGCCCTGCCACTGAACATCTGCGTCATGCCCAGATC	180
31	ValTyrSerValValLeuValValGlyLeuProLeuAsnIleCysValIleAlaGlnIle	51
181	TGCGCATCCCGCCGGACCCCTGACCCGTTCCGCTGTGTACACCCTGAACCTGGCACTGGCG	240
51	CysAlaSerArgArgThrLeuThrArgSerAlaValTyrThrLeuAsnLeuAlaLeuAla	71
241	GACCTGATGTATGCTGTTCACCTACCCCTACTTATCTATAACTAGCCAGGGGACCAC	300
71	AspLeuMetTyrAlaCysSerLeuProLeuLeuIleTyrAsnTyrAlaArgGlyAspHis	91
301	TGGCCCTTCGGAGACCTCGCCTGCGGCTTGTACGCTTCTCTCTATGCCAATCTACAT	360
91	TrpProPheGlyAspLeuAlaCysArgPheValArgPheLeuPheTyrAlaAsnLeuHis	111
361	GGCAGCATCCTGTTCCTCACTGCTAGCTTCCAGCGCTACCTGGGCATCTGCCACCCC	420
111	GlySerIleLeuPheLeuThrCysIleSerPheGlnArgTyrLeuGlyIleCysHisPro	131
421	CTGGCTTCCTGGCACAAGCGTGGAGGTGCGCGTCTGCTGGGTAGTGTGTGGAGTCGTG	480
131	LeuAlaSerTrpHisLysArgGlyGlyArgArgAlaAlaTrpValValCysGlyValVal	151
481	TGGCTGGCTGTGACAGCCCACTGCTGCCCCAGCGAGTCTTTGCTGCCACAGGCATCCAG	540
151	TrpLeuAlaValThrAlaGlnCysLeuProThrAlaValPheAlaAlaThrGlyIleGln	171
541	CGCAACCGCACTGTGTGCTAGCAGCTGAGCCCACTCCTGTCTACTCGCTACCTGCCC	600
171	ArgAsnArgThrValCysTyrAspLeuSerProProIleLeuSerThrArgTyrLeuPro	191
601	TATGGTATGCGCCCTCAGGTCATCGGCTTCTTGCTGCCCTTCATAGCCTTACTGGCTTGT	660
191	TyrGlyMetAlaLeuThrValIleGlyPheLeuLeuProPheIleAlaLeuLeuAlaCys	211
661	TATTGTGCGATGGCCCCGCGCTGTGTGGCCAGGATGGCCCAGCAGGTCTGTGGCCCAA	720
211	TyrCysArgMetAlaArgArgLeuCysArgGlnAspGlyProAlaGlyProValAlaGln	231
721	GAGCGGCGCAGCAAGCGGCTCGTATGGCTGTGGTGGTGGCAGCTGTCTTTGCCATCAGC	780
231	GluArgArgSerLysAlaAlaArgMetAlaValValValAlaAlaValPheAlaIleSer	251
781	TTCTGCGCTTTCCACATCACCAGACAGCCTACTTGGCTGTGCGCTCCACGCCCCGTGTC	840
251	PheLeuProPheHisIleThrLysThrAlaTyrLeuAlaValArgSerThrProGlyVal	271
841	TCTTGCCCTGTGCTGGAGACCTTGGCTGCTGCTACAAAGGCACTCGGCCCTTCGCCAGT	900
271	SerCysProValLeuGluThrPheAlaAlaAlaTyrLysGlyThrArgProPheAlaSer	291
901	GTCAACAGTGTTCTGGACCCATTCTCTTCTACTTCACACAACAGAAGTTCGGCGGCAA	960
291	ValAsnSerValLeuAspProIleLeuPheTyrPheThrGlnGlnLysPheArgArgGln	311
961	CCCCAGATCTCTTACAGAGGCTCACAGCCAAGTGGCAGAGGCAGAGTCTGAGGCCCC	1020
311	ProHisAspLeuLeuGlnArgLeuThrAlaLysTrpGlnArgGlnArgVal***	329

70 / 79

FIGURE 70



71 / 79

FIGURE 71

75+13, CODING	1	EQ-----	10	NGITAG	20	PP-----	30	EVYR--	40	DFRL--	50
P2UR_MOUSE	1	PAADLPWNS	10	TINGWGD	20	GY-----	30	KCFN--	40	DFVVL--	50
P2YR_CHICK	1	TEALISAAL	10	NGITAG	20	LGGWAAGN	30	ATTCSLTKT	40	GFQFY--	50
75+13, CODING	51	SSVLLV--	60	PPNICVLA	70	QV--ASR--	80	LTR--SAVIL	90	HEALALMV	100
P2UR_MOUSE	51	SSGVVCLL	60	CCNVVLY	70	GL--RLK--	80	NA--STTRG	90	HEVSS--	100
P2YR_CHICK	51	SVLLFTTS	60	FG--SSV--	70	M--VFHMP	80	SGI--LAGE	90	HEALALFV	100
75+13, CODING	101	CSLPLLTNN	110	RR--DRNDF	120	LA--RFRRH	130	TA--CHSHL	140	ELTCISFQY	150
P2UR_MOUSE	101	ASLPLLVYY	110	RR--DRNDF	120	TL--CHVRS	130	ETVLYCSIL	140	ELTCISVHC	150
P2YR_CHICK	101	LTFALDFNY	110	FNKTS--	120	SV--MORFI	130	SVVLYGSHL	140	ELTCISVHY	150
75+13, CODING	151	LVICHVIAW	160	HK--G--	170	WV--G--LA	180	IT--G--TA	190	VFAA--RTO	200
P2UR_MOUSE	151	LVLRPEH--	160	--W--A--	170	RR--PA--V	180	LA--QA--V	190	YVWT--ISG	200
P2YR_CHICK	151	TVVIRLKS	160	G--LKK--	170	VY--SSL--	180	SVVIA--II	190	--YSG--	200
75+13, CODING	201	SVV--V--	210	PPI--	220	LP--G--LVI	230	GLL--P--	240	ASTCR--	250
P2UR_MOUSE	201	TV--H--	210	RR--	220	A--SSVMGL	230	LA--V--	240	VWV--	250
P2YR_CHICK	201	SVV--V--	210	SVV--V--	220	--V--	230	MCV--P--	240	GVG--IVK	250
75+13, CODING	251	CRO--G--	260	VAQER--	270	--MAVV--	280	VAIS--	290	IRV--	300
P2UR_MOUSE	251	--L--	260	TGG--P--	270	VTIAL--	280	VAIS--	290	IRV--	300
P2YR_CHICK	251	IY--L--	260	--L--	270	SVLVII--	280	VAIS--	290	IRV--	300
75+13, CODING	301	STP---GV	310	PVETFA--	320	GT--P--	330	SVDP--	340	TO--K--	350
P2UR_MOUSE	301	STP---L	310	HT--N--	320	HT--P--	330	SVDP--	340	TO--K--	350
P2YR_CHICK	301	--P--	310	AFNDKVY	320	QV--P--	330	SVDP--	340	TO--K--	350
75+13, CODING	351	LLQRLTAKW	360	QR--RV--	370	380	390	400
P2UR_MOUSE	351	KKPTEPTP	360	SP--ARR--	370	LGL	380	HRP--RTV	390	LSVSSD	400
P2YR_CHICK	351	RETRKSSRS	360	EEVQS--	370	SEE	380	MTL--ILTEYK	390	ONGOTSL	400
75+13, CODING	401	410	420	430	440	450
P2UR_MOUSE	401	NDIRL.....	410	420	430	440	450
P2YR_CHICK	401	410	420	430	440	450

7 2 / 7 9

FIGURE 72

	9	18	27	36	45	54
5'	GCC ACC AAC GTG TTC ATC CTG TCA CTG	GCC GAT GTG CTG GTG ACA GCC ATC TGC	---	---	---	---
				Ala Asp Val Leu Val Thr Ala Ile Cys		
	63	72	81	90	99	108
CTG CCG GCC AGT CTG CTG GTA GAC ATC ACG GAA TCC TGG CTC TTT GGC CAT GCC						
---	---	---	---	---	---	---
Leu Pro Ala Ser Leu Leu Val Asp Ile Thr Glu Ser Trp Leu Phe Gly His Ala						
	117	126	135	144	153	162
CTC TGC AAG GTC ATC CCC TAT CTA CAG GCC GTG TCC GTG TCA GTG GTC GTG CTG						
---	---	---	---	---	---	---
Leu Cys Lys Val Ile Pro Tyr Leu Gln Ala Val Ser Val Ser Val Val Val Leu						
	171	180	189	198	207	216
ACT CTC AGC TCC ATC GCC CTG GAC CGC TGG TAC GCC ATC TGC CAC CCG CTG TTG						
---	---	---	---	---	---	---
Thr Leu Ser Ser Ile Ala Leu Asp Arg Trp Tyr Ala Ile Cys His Pro Leu Leu						
	225	234	243	252	261	270
TTC AAG AGC ACT GCC CGG CGC GCC CGC GGC TCC ATC CTC GGC ATC TGG GCG GTG						
---	---	---	---	---	---	---
Phe Lys Ser Thr Ala Arg Arg Ala Arg Gly Ser Ile Leu Gly Ile Trp Ala Val						
	279	288	297	306	315	324
TCG CTG GCT GTC ATG GTG CCT CAG GCT GCT GTC ATG GAG TGT AGC AGC GTG CTG						
---	---	---	---	---	---	---
Ser Leu Ala Val Met Val Pro Gln Ala Ala Val Met Glu Cys Ser Ser Val Leu						
	333	342	351	360	369	378
CCC GAG CTG GCC AAC CGC ACC CGC CTC CTG TCT GTC TGT GAT GAG CGC TGG GCA						
---	---	---	---	---	---	---
Pro Glu Leu Ala Asn Arg Thr Arg Leu Leu Ser Val Cys Asp Glu Arg Trp Ala						
	387	396	405	414	423	432
GAC GAC CTG TAC CCC AAG ATC TAC CAC AGC TGC TTC TTC ATT GTC ACC TAC CTG						
---	---	---	---	---	---	---
Asp Asp Leu Tyr Pro Lys Ile Tyr His Ser Cys Phe Phe Ile Val Thr Tyr Leu						
	441	450	459	468	477	486
GCC CCA CTG GGC CTC ATG GCC ATG GCC TAT TTC CAG ATC TTC CGC AAG CTC TGG						
---	---	---	---	---	---	---
Ala Pro Leu Gly Leu Met Ala Met Ala Tyr Phe Gln Ile Phe Arg Lys Leu Trp						
	495	504	513	522	531	540
GGC CGC CAG ATC CCC GGC ACC ACC TCG GCC CTG GTG CGC AAC TGG AAG CGG CCC						

7 3 / 7 9

FIGURE 73

```

-----
Gly Arg Gln Ile Pro Gly Thr Thr Ser Ala Leu Val Arg Asn Trp Lys Arg Pro
      549      558      567      576      585      594
TCA GAC CAG CTG GAC GAC CAG GGC CAG GGC CTG AGC TCA GAG CCC CAG CCC CGG
-----
Ser Asp Gln Leu Asp Asp Gln Gly Gln Gly Leu Ser Ser Glu Pro Gln Pro Arg

      603      612      621      630      639      648
GCC CGC GCC TTC CTG GCC GAG GTG AAA CAG ATG CGA GCC CGG AGG AAG ACG GCC
-----
Ala Arg Ala Phe Leu Ala Glu Val Lys Gln Met Arg Ala Arg Arg Lys Thr Ala

      657      666      675      684      693      702
AAG ATG CTG ATG GTG GTG CTG CTG GTC TTC GCC CTC TGC TAC CTG CCC ATC AGT
-----
Lys Met Leu Met Val Val Leu Leu Val Phe Ala Leu Cys Tyr Leu Pro Ile Ser

      711      720      729      738      747      756
GTC CTC AAC GTC CTC AAG AGG GTC TTC GGG ATG TTC CGC CAA GCC AGC GAC CGA
-----
Val Leu Asn Val Leu Lys Arg Val Phe Gly Met Phe Arg Gln Ala Ser Asp Arg


      765      774      783      792      801      810
GAG GCC ATC TAC GCC TGC TTC ACC TTC TCC CAC TGG CTG GTG TAC GCC AAC AGC
-----
Glu Ala Ile Tyr Ala Cys Phe Thr Phe Ser His Trp Leu Val Tyr Ala Asn Ser

      819      828      837
GCC GCC AAT CCC CTC CTC TAC TCC TTC CTC CCT 3'
-----
Ala Ala

```


7 4 / 7 9

FIGURE 7 4



7 5 / 7 9

FIGURE 75



7 6 / 7 9

FIGURE 76

h3H2-17(5-3)	1	10	20	30	40	50	50
p3H2-17(5')	1	GTGGGCTGG	TGGGCAACAT	CCTGGCTTCC	TGGCACAAGC	GTGGAGGTGG	50
h3H2-17(5-3)	51	60	70	80	90	100	100
p3H2-17(5')	51	CCGTGCTGCT	TGGGTAGTGT	GTGGAGTGTGT	GTGGCTGGCT	GTGACAGGCT	100
h3H2-17(5-3)	101	110	120	130	140	150	150
p3H2-17(5')	101	AGTGGCTGG	CATAGGCATC	TTCGCTGCTC	CAGGCTGCTC	GGGTAGGCTC	150
h3H2-17(5-3)	151	160	170	180	190	200	200
p3H2-17(5')	151	AGTGGCTGG	ATGAGCTCAG	GGTGGCTGCTC	GTGGCTGCTC	ACCTATATGCTC	200
h3H2-17(5-3)	201	210	220	230	240	250	250
p3H2-17(5')	201	AGTGGCTGG	GGTGGCTGCTC	GTGGCTGCTC	GTGGCTGCTC	GTGGCTGCTC	250
h3H2-17(5-3)	251	260	270	280	290	300	300
p3H2-17(5')	251	AGTGGCTGG	GTGGCTGCTC	GTGGCTGCTC	GTGGCTGCTC	GTGGCTGCTC	300
h3H2-17(5-3)	301	310	320	330	340	350	350
p3H2-17(5')	301	CCAGCAGGTC	CTGTGGCCCA	AGAGCGGGCC	AGCAAGGGCG	CTCGTATGGC	350
h3H2-17(5-3)	351	360	370	380	390	400	400
p3H2-17(5')	351	TGTGGTGGTG	GCAGCTGTCT	TTCGCTGCTC	GTGGCTGCTC	CTCTAC	400


77 / 79

FIGURE 77

1	TGACTCCCTGAACATAGGAAACCCACCTGGGCAGCCATGGAATGGGACAATGGCACAGGC	60
1	MetGluTrpAspAsnGlyThrGly	8
61	CAGGCTCTGGGCTTGCCACCCACCACCTGTGTCTACCGGAGAACTTCAAGCAACTGCTG	120
8	GlnAlaLeuGlyLeuProProThrThrCysValTyrArgGluAsnPheLysGlnLeuLeu	28
121	CTGCCACCTGTGTATTCCGGCGGTGCTGGCGGCTGGCCTGCCGCTGAACATCTGTGTCATT	180
28	LeuProProValTyrSerAlaValLeuAlaAlaGlyLeuProLeuAsnIleCysValIle	48
181	ACCCAGATCTGCACGTCCCGCGGGCCCTGACCCGCACGGCCGTGTACACCTAAACCTT	240
48	ThrGlnIleCysThrSerArgArgAlaLeuThrArgThrAlaValTyrThrLeuAsnLeu	68
241	GCTCTGGCTGACCTGCTATATGCCTGCTCCCTGCCCTGCTCATCTACAACCTATGCCCAA	300
68	AlaLeuAlaAspLeuLeuTyrAlaCysSerLeuProLeuLeuIleTyrAsnTyrAlaGln	88
301	GGTGATCACTGGCCCTTTGGCGACTTGGCCTGCCGCTGGTCCGCTTCCTCTTCATGCC	360
88	GlyAspHisTrpProPheGlyAspPheAlaCysArgLeuValArgPheLeuPheTyrAla	108
361	AACCTGCACGGCAGCATCCTCTTCCTCACCTGCATCAGCTTCCAGCGCTACCTGGGCATC	420
108	AsnLeuHisGlySerIleLeuPheLeuThrCysIleSerPheGlnArgTyrLeuGlyIle	128
421	TGCCACCCGCTGGCCCCCTGGCACAACGTTGGGGCGCGCGGGCTGCCTGGCTAGTGTGT	480
128	CysHisProLeuAlaProTrpHisLysArgGlyGlyArgArgAlaAlaTrpLeuValCys	148
481	GTAACCGTGTGGCTGGCCGTGACAACCCAGTGCCTGCCACAGCCATCTTCGCTGCCACA	540
148	ValThrValTrpLeuAlaValThrThrGlnCysLeuProThrAlaIlePheAlaAlaThr	168
541	GGCATCCAGCGTAACCGCACTGTCTGTATGACCTCAGCCCGCCTGCCCTGGCCACCCAC	600
168	GlyIleGlnArgAsnArgThrValCysTyrAspLeuSerProProAlaLeuAlaThrHis	188
601	TATATGCCCTATGGCATGGCTCTCACTGTTCATCGGCTTCTGTGCTGCCCTTTGCTGCCCTG	660
188	TyrMetProTyrGlyMetAlaLeuThrValIleGlyPheLeuLeuProPheAlaAlaLeu	208
661	CTGGCCTGCTACTGTCTCCTGGCCTGCCGCTGTGCCGCCAGGATGGCCCGGCAGGCCT	720
208	LeuAlaCysTyrCysLeuLeuAlaCysArgLeuCysArgGlnAspGlyProAlaGluPro	228
721	GTGGCCCGAGGAGCGGCGTGGCAAGGCGCGCCGATGGCCGTGGTGGTGGCTGTGCCTTT	780
228	ValAlaGlnGluArgArgGlyLysAlaAlaArgMetAlaValValValAlaAlaAlaPhe	248
781	GCCATCAGCTTCTGCCTTTTCACATCACCAGACAGCCTACCTGGCAGTGGGCTCGACG	840
248	AlaIleSerPheLeuProPheHisIleThrLysThrAlaTyrLeuAlaValGlySerThr	268
841	CCGGGCGTCCCTGCACCTGTATTGGAGGCCTTTGCAGCGGCCTACAAAGGCACGCGGCCG	900
268	ProGlyValProCysThrValLeuGluAlaPheAlaAlaAlaTyrLysGlyThrArgPro	288
901	TTTGCCAGTGCCAAACAGCGTGTGGACCCCATCTCTTCTACTTACCCAGAAGAAGTTC	960
288	PheAlaSerAlaAsnSerValLeuAspProIleLeuPheTyrPheThrGlnLysLysPhe	308
961	CGCCGCGGACCACATGAGCTCCTACAGAAACTACAGCCAAATGGCAGAGGCAGGGTCGC	1020
308	ArgArgArgProHisGluLeuLeuGlnLysLeuThrAlaLysTrpGlnArgGlnGlyArg	328
1021	TGA	1023
328	***	329

7 8 / 7 9

FIGURE 78



7 9 / 7 9

FIGURE 79

human prino, mouseFULL3H2	1 1	10 10	20 20	30 30	40 40	50 50
	1	10	20	30	40	50
	1	10	20	30	40	50
human prino, mouseFULL3H2	51 51	60 60	70 70	80 80	90 90	100 100
	51	60	70	80	90	100
	51	60	70	80	90	100
human prino, mouseFULL3H2	101 101	110 110	120 120	130 130	140 140	150 150
	101	110	120	130	140	150
	101	110	120	130	140	150
human prino, mouseFULL3H2	151 151	160 160	170 170	180 180	190 190	200 200
	151	160	170	180	190	200
	151	160	170	180	190	200
human prino, mouseFULL3H2	201 201	210 210	220 220	230 230	240 240	250 250
	201	210	220	230	240	250
	201	210	220	230	240	250
human prino, mouseFULL3H2	251 251	260 260	270 270	280 280	290 290	300 300
	251	260	270	280	290	300
	251	260	270	280	290	300
human prino, mouseFULL3H2	301 301	310 310	320 320	330 330	340 340	350 350
	301	310	320	330	340	350
	301	310	320	330	340	350

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C12N15/12 C07K14/705 C12Q1/68

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 C07K C12Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	MOLECULAR ENDOCRINOLOGY, vol. 5, January 1991 pages 1331-1338, M.T.HARRIGAN ET AL. 'Identification of a gene induced by glucocorticoids in murine T-cells : a potential G protein-coupled receptor'	4-6, 10, 11, 13
Y	see the whole document ---	14-18
Y	WO,A,92 01810 (LERNER MICHAEL R ;LERNER ETHAN A (US)) 6 February 1992 see abstract; claims 1-17 ---	14-18
A	EP,A,0 578 962 (AMERICAN CYANAMID CO) 19 January 1994 see example 2 --- -/--	1-3

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

18 December 1995

Date of mailing of the international search report

05.01.96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+ 31-70) 340-3016

Authorized officer

Gurdjian, D

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	SCIENCE, vol. 244, 1989 pages 569-571, LIBERT F. ET AL. 'Selective amplification and cloning of four new members of the G protein-coupled receptor family' cited in the application see the whole document -----	1-3

INTERNATIONAL SEARCH REPORT

Information on patent family members

Intern al Application No

PCT/JP 95/01599

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9201810	06-02-92	US-A- 5462856	31-10-95
		EP-A- 0539518	05-05-93
		JP-T- 6502757	31-03-94
<hr/>			
EP-A-0578962	19-01-94	AU-B- 4203493	20-01-94
		CA-A- 2100616	18-01-94
		JP-A- 6153954	03-06-94
<hr/>			